

Original Research Article

Effect of combined application of herbicides on weed flora, Yield attributes Yield, and Profitability of chickpea (*Cicer arietinum* L.)

Abstract

Chickpea (*Cicer arietinum* L.) commonly known as gram or Bengal gram belongs to the family Leguminaceae (Fabaceae). It's one of the most important *rabi* season pulse crops grown in India for economic importance and to improve soil fertility. The productivity of chickpea is low despite of having high-yielding varieties and new agronomic practices. One of the causes of poor productivity is the infestation of weeds in the field of chickpea. It is a poor competitor to weeds because of its slow growth rate and limited leaf area development at early stages. To overcome this, a field experiment was conducted during the *rabi* seasons of 2021-22 at the Agronomy Research Farm of the Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (UP). Replicated thrice set of eleven treatments (T₁) Pendimethalin 30 EC @ 1.0 kg a.i./ha PE, (T₂) Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE, (T₃) Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE, (T₄) Propaquizafop 10 EC @ 0.075 kg a.i./ha PoE, (T₅) Quizalfop ethyl 5 EC @ 40 g a.i./ha PoE, (T₆) Pendimethalin 30 EC @ 1.0 kg a.i./ha PE followed by (Fb) Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE, (T₇) Pendimethalin 30 EC @ 1.0 kg a.i./ha PE fb by Propaquizafop 10 EC @ 0.075 kg a.i./ha PoE, (T₈) Pendimethalin 30 EC @ 1.0 kg a.i./ha PE fb by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE, (T₉) Pendimethalin 30 EC @ 1.0 kg a.i./ha pE fb by Quizalfop ethyl 5 EC @ 40 g a.i./ha PoE, (T₁₀) Weed free and (T₁₁) Weedy check, was laid out in randomized block design. Weed management was done as per treatment. Other crop management practices were followed as per the recommendation of the area. Pendimethalin @ 1.0 kg a.i./ha PE fb Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha as PoE during the experiment recorded significantly less weed density, weed dry matter, and higher weed control efficiency, yield attributes (number of pods/plant, number of seeds/pod and test weight), yield (grain yield, stover yield, biological yield, and harvest yield). Economically it recorded higher gross returns, the net return, and B:C ratio was noted and proved to be more remunerative as compared to other herbicide applications used in chickpea crop.

Keywords: Herbicide combination, Chickpea, Weed management, Clodinafoppropargyl

INTRODUCTION

Pulses are an integral part of Indian agriculture. Chickpea (*Cicer arietinum* L.) commonly known as gram or Bengal gram belongs to the family Leguminaceae (Fabaceae). It's one of the most important *rabi* season pulse crops grown in India for economic importance and to improve soil fertility (Nasir, M., & Sidhu, 2012). In Indian pulses, chickpea is the second most important component of the diet after cereals. The net availability of food grains per capita increased day by day from 144.1 kg/year in 1951 to 179.6 kg/year in 2019 despite population growth however the net obtainability of pulses crop has reduced from 25 kg/year in 1961 to 17.5 kg/year in 2019

(Anonymous, 2020). Chickpea is a major pulse in India which contributed about 71% of the world's area (13.57 million hectares) and 67% of the world's production. India ranked first in the area with 9.93 million hectares (34%) and first in production with 9.53 million tones (26%) in the world, followed by Pakistan, Australia, and Iran. In India, during 2019-20, pulses were cultivated over more than 29 million hectares of area and registered the highest ever production of 25.90 mt with a productivity level of 908 kg/hectares. The exponential growth rate in pulse production over the past year has been more than 9 percent. (Anonymous, 2020) The highest production of 6120 kg/ha is observed in Israel followed by Yemen, Canada, and Egypt. India's productivity was 920 kg/ha yields, (Anonymous, 2020). Today, 80% of total pulses production, in India, is realized in six states namely, Madhya Pradesh, Maharashtra, Rajasthan, Andhra Pradesh, Karnataka, and Uttar Pradesh. The productivity of chickpea is low despite of having high-yielding varieties and new agronomic practices. One of the causes of poor productivity is the infestation of weeds in the field of chickpea. It is a poor competitor to weeds because of its slow growth rate and limited leaf area development at early stages (Peerzada *et al.* 2017). Crop yield losses due to non-adoption of weed management practices particularly in irrigated conditions. Poonia and Pithia (2013) reported a 54.7% loss in grain yield due to weed competition in chickpea under rainfed conditions. Major weeds associated with chickpea vary with crops and locations. The Directorate of Weed Research (DWR), Jabalpur, has created a Weed Atlas for major weeds in major crops in 435 districts across 19 states, as well as a weed identification handbook (Naidu 2012). Its findings concluded that weeds have economic significance in specific crops (Rao *et al.* 2014). Weeds for Chickpea are; *Chenopodium album* L., *Avena fatua* L., *Medicago denticulata*, *Chicorium intybus*, *Convolvulus arvensis* L., *Lathyrus aphaca* L. *Lathyrus sativus* L., *Vicia sativa*, *Cyperus rotundus* L., *Phalaris minor*, *Avena ludoviciana*, *Euphorbia geniculata*, and *Melelotus species*. In the total annual loss of agricultural produce from various pests in India, weeds roughly account for 37%, insects for 29%, diseases for 22%, and other pests for 12% (Yaduraju, 2006). Weed emergence with the *rabi* sown chickpea crop creates competition unless controlled timely and effectively. Therefore, it is need of our move from the expensive manual and mechanical weed control to chemical weed control practices (Marwat *et al.*, 2003). The predominant method of weed control by mechanical hoeing and manual weeding over an extensive scale is declining because availability of labor due to shifted from agricultural fields to industries for better and assured wages. Eradication of weed through manual weeding and hoeing is expensive, tedious, and time-consuming, on the other hand use of single herbicides cannot achieve complete weed control due to their selective destruction. Therefore use of herbicide will be more effective if supplemented by manual weeding and hoeing. It is a well-known fact that relying on a single weed control method is ineffective, and an integrated approach may be required. Among several herbicides in the market *viz.* Pendimethalin, Clodinafoppropargyl, Quizalofop-ethyl, Propaquizafop, and Imazethapyr are currently being used for controlling both grassy and broad-leaved weeds but their effects under various climatic conditions are not well defined. It was recently found that the herbicide Clodinafoppropargyl is very effective in controlling weeds in chickpea (Dubey *et al.*, 2018). This herbicide is active against grasses but their effects may differ in different locations depending on soil type, intensity, and weed flora type, among other factors. As a result, it is important to compare the effectiveness of various promising herbicides in terms of chickpea productivity and weed competition in weed-free environments, where weed free environment can be achieved by effective hand weeding. Vaishya *et al.*, (1996) reported that integrated weed management (pre-emergence application of Pendimethalin @ 1.0kg/ha Followed by (*fb*) one hand weeding) was found to be the most economical. Among the chemical weed control treatments, Pendimethalin @ 1.0 kg a.i./ha produced the highest yield, net

monetary returns, and B: C ratio, and was found to be the most effective and economical in controlling weeds and increasing chickpea yield. A suitable herbicide for effective control of mixed weed flora is required for better adoption in this crop by farmers. The introduction of herbicides has made it possible to control a broad spectrum of weeds in pulses effectively at a reasonable cost. Weed management in chickpea is a crucial component of plant protection thus improving the production potential of the crop.

MATERIALS AND METHODS

A field experiment was conducted during 2021-22 *rabi* season at Agronomy Research Farm of the Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (UP) India, which falls in a subtropical climate and situated at 26° 47' North latitude, 82° 12' East longitudes with an altitude of 113 meters above sea level. During the crop growing season of the year 2021-22, the weekly maximum and minimum temperatures, relative humidity, sunshine hours, wind velocity, evaporation ranged between 32.0 °C and 5.7 °C, 87.40 and 74.70 %, 9.0 and 4.7 hours, 5.5 and 2.0 kmph, 6.4 and 2.1 mm respectively. The soil of the experimental site was homogeneous in fertility with uniform textural makeup. The alluvial soils of Indo-Gangetic plains in general are deep, flat, and well drained with low available nitrogen and medium in available phosphorus and potassium (Singh *et al.*, 2015). Chickpea variety RVG 202 was used at seed rate of 80 kg ha⁻¹.

Observation on weed

Weed observations were recorded at 30, 60, and 90 Days after sowing (DAS). The quadrat of 1 m sq. was randomly placed at three places in each lot and then species-wise and total weed counts were recorded. In weed biomass, all the associated weeds were collected, randomly from 0.25 m² quadrat at four places in each plot. The weeds were kept in paper bags and dried in the oven at 60±2°C for 24-25 hours (up to constant weight) and dry weight was recorded at 30, 60, and 90 DAS and expressed as gm⁻². For weeds, the original values were transformed using the square root of X+0.5 transformations and analyzed statistically. Weed control efficiency (%) of treatment expressed in percentage for controlling weeds in comparison to weedy check and based on the given formula.

$$\text{WCE (\%)} = \frac{\text{Dry weight of weeds in control plot} - \text{Dry weight of weeds in treated plot}}{\text{Dry weight of weeds in control plot}} \times 100$$

Observation on crop

Crop yield and yield attributing characters were recorded as the number of pods plant⁻¹ on five randomly selected plants per plot were counted and averaged. The total number of pods from each five sample plants was threshed and the average number of seeds pods⁻¹ was recorded. From a lot of threshed clean seeds of each plot, a random seed sample was taken and one hundred seeds were counted from the samples of each plot, and the weight of seeds was recorded on an electronic balance. Seed yield and Stover yield obtained from each plot were added to obtain biological yield in kilogram from each plot and converted to quintal per hectare. On a double pan balance, the weight of clean seeds received from each plot was recorded. Finally, the seed yield plot⁻¹ was multiplied by the required value to produce ha⁻¹. The residual yield was calculated by subtracting the seed yield from the biological yield of each net plot under each treatment. Using the same conversion factor, the

results were then translated into stover yields ha^{-1} . Harvest Index (%) refers to the ratio of economic yield (seed yield) to the biological (seed + stover) yield under a particular treatment and it was expressed in percentage:

$$\text{Harvest Index} = \frac{\text{Economic Yield}}{\text{Biological Yield}} \times 100$$

Regarding economic analysis of study, the cost of cultivation was calculated treatment-wise, based on prevailing local market prices of different inputs used in the cultivation. Gross returns of grain yield and stover yield were computed in Rs ha^{-1} by using the minimum support prices for grains and prevailing local market price for stover. The gross return in each treatment was obtained by adding the monetary value of grain and stover yield in Rs ha^{-1} . The net return for each treatment was calculated by deducting the cost of cultivation from the respective gross returns. Benefit: The cost ratio in terms of net return per rupee investment was calculated by using the following formula.

$$\text{B:C ratio} = \frac{\text{net return (Rs/ha)}}{\text{Cost of cultivation (Rs/ha)}}$$

Statistical analysis

The data obtained were subjected to statistical analysis as outlined by **Gomez & Gomez (1984)**. The treatment differences were tested by the least significant difference at a 5% level of significance.

RESULTS AND DISCUSSION

Weed indices/ attributes

The data on total weed density (no. m^{-2}) given in Table 1 indicate that the effect of various weed management practices was significant on the density of weeds at 30, 60, and 90 DAS. Weed management practices exerted a significant effect on the density of total weeds. Weedy check recorded a significantly higher density of total weeds (8.96, 11.62, and 12.42) over the rest of the treatments at all stages of crop growth while weed-free treatments registered the least density of total weeds (0.71, 0.71, and 0.71). Among the herbicidal treatments at 30, 60, and 90 DAS the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Clodinafoppropargyl 15 WP @ 0.060 kg a.i./ha PoE recorded minimum density (5.23, 3.83, and 3.97) of weeds which were statistically at par with Pendimethalin EC @ 1.0 kg a.i./ha PE *fb* Imazethapyr 10 % SL @ 0.06 kg a.i./ha PoE. These two treatments reduced the weed density significantly than the rest of the treatments. This might be due to effective weed control by both herbicides at all stages of crop growth whereas weedy check treatment recorded significantly highest density of weeds (80.00, 135.00, and 154.00 m^{-2}) at 30, 60, and 90 DAS respectively. These results were in agreement with the results of **Yadav *et al.* (2017)** and **Jaswal and Menon (2020)**. Data on dry matter of total weeds recorded at 30, 60, and 90, DAS in Table 1 indicated that weedy check recorded maximum weed dry matter of total weeds (4.05, 9.38 and 10.09) at 30, 60 and 90, DAS respectively. While weed-free treatment recorded the lowest dry matter of total weeds at 30, 60, and 90, DAS due to better control of weeds where weeds free condition were maintained by hand weeding. Among the herbicidal treatments, at 30 DAS application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Clodinafop propargyl 15 WP @ 0.060 kg

a.i./ha as PoE recorded significantly lesser dry matter (2.34, 3.12 and 3.26 gm⁻²) which was statistically at par with Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Imazethapyr 10 % SL @ 0.060 kg a.i./ha PoE while significantly lesser than the rest of the treatments. A similar trend was observed at 60 and 90 DAS also during the experimentation. Which might be due to effective control of weeds leading to reduction in dry weight. In the weedy check treatment, the dry matter of weeds was significantly higher because of the high weed population and greater capacity of weeds in utilizing the sunlight, nutrients, moisture, CO₂, space, *etc.*, over the rest of the treatments. In treatment, weed-free for up to 90 days and application of pre-emergence herbicides in combination with PoE application of herbicides recorded the less weed population at a critical stage of crop growth resulting in lower weed dry matter. These results were also reported by **Singh *et al.* (2006)**, **Bhutada and Bhale (2014)**, **Kumar *et al.* (2015)**, and **Kumar *et al.* (2020)**. The data regarding weed control efficiency as influenced by various weed control treatments are given in Table 1. At harvest, the weed control efficiency was highest in the weed-free (100%) treatment which was significantly superior to all other treatments. Among herbicidal treatments the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Clodinafoppropargyl 15 WP @ 0.060 kg a.i./ha noted significantly higher value (90.00%) of weed control efficiency which was found at par with Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Imazethapyr 10 % SL @ 0.060 kg a.i./ha (POE) (89.61) and significantly higher than the rest of the herbicidal treatments. Due to the herbicide's maximal uptake and higher assimilation occurring right after weed emergence, the application of the herbicides Pendimethalin and oxyfluorfen before emergence reduces the density of the weeds and results in high efficiency of weed control. Due to their higher weed control effectiveness, Clodinafoppropargyl and ImazethapyrPoE applications reduced weed intensity and dry weight more than weedy check. These findings were parallel to the conclusions of **Singh and Jain (2017)**, **Rathod *et al.* (2017)**, and **Singh *et al.* (2020)**.

Yield attributes

The data on yield contributing characters *viz.* numbers of pods plant⁻¹, the number of seeds pod⁻¹, and the weight of 1000 seeds (g) are presented in Table 2. It is obvious from the results that the number of pods/plants⁻¹ was affected significantly due to different weed management practices.

Among different weed management practices the maximum number of pods plant⁻¹ (57.00) was recorded with weed-free treatments and the minimum (33.00) was the weedy check. Herbicidal treatments also significantly influenced the number of pods plant⁻¹. The maximum number of pods plant⁻¹ (54.60) was recorded with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE which was at par with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE (53.4) and significantly higher than the rest of the chemical treatments. The number of seeds/pod was also affected significantly due to different weed management practices of the chickpea crop. The maximum number of seeds pod⁻¹ (1.96) was recorded with weed-free treatments and the minimum (1.32) with the weedy check. Among the herbicidal treatments, the maximum number of seeds/pod (1.88) was recorded with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE being at par with the application of

Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE (1.85) and proved significantly higher than the rest of the treatments. Maximum test weight (200.40g) was recorded with weed-free treatments and minimum (178.80g) with the weedy check. Among the herbicidal treatments, the maximum test weight (197.70g) was recorded with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE. All parameters related to yield *viz.*, number of pods/plant, number of seeds/pod (Table 2) were significantly influenced by weed management practices while the test weight failed to reach the level of significance. Significantly higher values for yield attributes (Table 2) were observed with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Clodinafoppropargyl 15 WP @ 0.060 kg a.i./ha PoE. This might be due to effective control of weeds reducing the weed competition leading to the better expression of yield attributes. Similar findings were reported by **Kachhadiya *et al.* (2009)**, **Khope *et al.* (2011)**, and **Singh *et al.* (2019)**.

Yield

Grain yield is a function of the source and sinks relationship, where the source is the various growth parameters and the sink is yield attribute parameters such as dry matter accumulation, nodule number per plant, pod number plant⁻¹, seed number pod⁻¹, and test weight. Different weed management treatments influenced significantly the seed, and stover yield of chickpea and the data are embodied in Table 2. The weed-free treatment recorded maximum seed yield (20.60 q ha⁻¹) and minimum seed yield (13.80 q ha⁻¹) was noted with weedy check. Severe crop weed competition in weedy check might have effected crop growth and recorded low yields as weeds utilize a large amount of moisture and nutrients than crop due to no weed management practice. Among the herbicidal treatments the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE recorded a significantly higher seed yield (19.80 q ha⁻¹) being at par with Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Imazethapyr 10 % SL @ 0.06 kg a.i./ha PoE, while significantly higher than the rest of the herbicidal treatments. This might be due to effective control of weeds resulting in higher yield. Similar findings were reported by **Punia *et al.* (2011)** and **Singh *et al.* (2014)**. Weed management practices showed a significant variation in stover yield of chickpea. Weed-free treatment recorded maximum stover yield (37.43 kg ha⁻¹) and minimum was with the weedy check (26.50 qha⁻¹). Data further reveals that among herbicidal treatments the maximum stover yield (36.27 q ha⁻¹) was recorded with Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE which was statistically at par with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Imazethapyr 10 % SL 0.06 kg a.i./ha PoE while significantly higher than the rest of the herbicidal treatments. These results were in agreement with those of **Poonia *et al.* (2013)**, **Chavada *et al.* (2018)**, and **Singh *et al.* (2020)**. Among weed management practices significantly higher biological yield (58.03 q ha⁻¹) was observed with weed-free treatments and least biological yield (39.85 q ha⁻¹) with weedy check treatment. In the case of herbicidal treatments the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Clodinafoppropargyl 15 WP @ 0.060 kg a.i./ha PoE was statistically at par with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* Imazethapyr 10 % SL @ 0.06 kg a.i./ha PoE but significantly higher than the rest of herbicidal treatments. This might be due to effective control of weeds by avoiding weed competition and resulting in higher biomass production. Similar results were reported by **Kumar *et al.* (2014)** and **Singh *et al.* (2020)**. Among the weed management treatments, the maximum harvest index was recorded with weed-free (35.49 %) and minimum with the weedy check (34.62 %). Among herbicidal treatments the application of Pendimethalin 30 EC @

1.0 kg a.i./ha PE *fb* Clodinafoppropargyl 15 WP @ 0.060 kg a.i./ha PoE recorded a maximum harvest index of 35.38 %. The similar result was obtained by **Chavada et al. (2018)**.

Economics analysis

The economic analysis of the weed control treatments was worked out based on input-output analysis. The cost of cultivation (Rsha⁻¹) incurred on weed control treatments was added to the common cost of treatments and arrived the total cost of cultivation. The data on economic analysis presented in Table 3 showed that the highest cost of cultivation (39024 Rsha⁻¹) was recorded under the treatment weed free and the lowest was with the weedy check (36024 Rsha⁻¹). Among herbicide treatments, the higher cost (38564 Rsha⁻¹) was recorded with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE. In Table 3 the data on gross return computed under different treatments showed that the highest gross return (108803 Rs/ha) was achieved with weed-free treatments and the lowest with the weedy check (72985 Rsha⁻¹). Among herbicidal treatments, the highest gross return (104595 Rsha⁻¹) was recorded with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE. Data on net return computed under different treatments showed that the highest net return (66031 Rsha⁻¹) was obtained with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE. The data on B:C ratio computed under different treatments showed that the highest B:C ratio (1.71 Rs/Rupees investment) was recorded with the application of Pendimethalin 30 EC @ 1.0 kg a.i./ha PE *fb* by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE and lowest with the weedy check (1.02 Rs/Rupees investment). The present finding conforms with the finding of **Dungarwal et al. (2002)**, **Ratnam et al. (2011)**, **Meena et al. (2011)**, and **Dubey et al. (2018)**.

Table:1 Effect of herbicide combination on Weed density, Weed dry weight (g) at 30, 60, 90 DAS, and Weed control efficiency (%) of Chickpea

S. No.	Treatments	Weed density (no m ⁻²)			Weed dry weight (gm ⁻²)			Weed control efficiency (%)
		30 (DAS)	60 (DAS)	90 (DAS)	30 (DAS)	60 (DAS)	90 (DAS)	
T ₁	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE	5.54 (30.30)	6.35 (40.00)	6.67 (44.00)	2.55 (6.06)	5.14 (26.00)	5.43 (29.04)	71.43
T ₂	Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE	7.46 (55.40)	5.95 (35.20)	6.20 (38.00)	3.40 (11.08)	4.82 (22.88)	5.05 (25.08)	75.32
T ₃	Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE	7.83 (61.20)	5.83 (33.60)	6.12 (37.00)	3.56 (12.24)	4.72 (21.84)	4.97 (24.42)	75.97
T ₄	Propaquizafop 10 EC @ 0.075 kg a.i./ha PoE	8.51 (72.00)	7.12 (50.40)	7.43 (55.00)	3.85 (14.40)	5.76 (32.76)	6.06 (36.30)	64.29
T ₅	Quizalfop ethyl 5 EC @ 40	7.66	6.39	6.63	3.48	5.17	5.41	71.56

	g a.i./ha PoE	(58.40)	(40.40)	(43.80)	(11.68)	(26.26)	(28.91)	
T ₆	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE	5.29 (27.60)	3.93 (15.00)	4.06 (16.00)	2.36 (5.10)	3.19 (9.75)	3.31 (10.56)	89.61
T ₇	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Propaquizafop 10 EC @ 0.075 kg a.i./ha PoE	5.42 (29.00)	4.45 (19.31)	4.67 (21.40)	2.51 (5.80)	3.62 (12.63)	3.69 (13.95)	75.13
T ₈	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE	5.23 (27.00)	3.83 (14.20)	3.97 (15.40)	2.34 (5.02)	3.12 (9.23)	3.26 (10.16)	90.00
T ₉	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Quizalfop ethyl 5 EC @ 40 g a.i./ha PoE	5.47 (29.40)	4.51 (20.00)	4.73 (22.00)	2.52 (5.88)	3.67 (13.00)	3.87 (14.52)	76.34
T ₁₀	Weed free	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	100.00
T ₁₁	Weedy check	8.96 (80.00)	11.62 (135.0)	12.42 (154.00)	4.05 (16.00)	9.38 (87.75)	10.09 (101.64)	0.00
	S_{Em}±	0.29	1.00	0.51	0.17	0.61	0.76	0.89
	CD (p = 0.05)	0.88	2.89	1.53	0.53	1.83	2.26	2.52

Table:2 Effect of herbicide combination on Yield attribute, Yields, and Harvest Index (%) of Chickpea

S. No.	Treatments	Yield attribute			Yield (q ha ⁻¹)			Harvest Index (%)
		Pods/plant	Seeds/pod	Test Weight (g)	Grain Yield	Stover Yield	Biological yield	
T ₁	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE	45.03	1.42	180.00	16.50	30.78	47.28	34.89
T ₂	Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE	46.80	1.40	180.60	17.00	31.63	48.63	34.95
T ₃	Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE	47.00	1.38	181.80	17.20	31.44	49.15	34.99
T ₄	Propaquizafop 10 EC @ 0.075 kg a.i./ha PoE	43.60	1.46	179.20	16.00	30.05	46.02	34.82
T ₅	Quizalfop ethyl 5 EC @ 40 g a.i./ha PoE	44.00	1.44	179.80	16.30	30.47	46.77	34.85

T ₆	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE	53.40	1.85	196.00	19.60	35.83	55.43	35.35
T ₇	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Propaquizafop 10 EC @ 0.075 kga.i./ha PoE	47.90	1.45	192.00	16.78	31.07	48.19	35.30
T ₈	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Clodinafoppropargyl 15 WP @ 0.06 kg a.i. /ha PoE	54.60	1.88	197.70	19.80	36.27	55.95	35.38
T ₉	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Quizalfop ethyl 5 EC @ 40 g a.i./ha PoE	48.00	1.47	190.70	16.83	30.66	47.79	35.19
T ₁₀	Weed free	57.00	1.96	200.40	20.60	37.43	58.03	35.49
T ₁₁	Weedy check	33.00	1.32	178.80	13.80	26.05	39.85	34.62
	SEm±	0.65	0.12	9.41	0.87	1.49	1.34	1.25
	CD(p = 0.05)	1.95	0.38	NS	2.58	4.41	4.00	NS

Table:3Effect of herbicide combination on Cost of cultivation, Gross return, Net return, and B: C ratio of Chickpea

S. No.	Treatments	Cost of cultivation (Rs ha ⁻¹)	Gross return (Rs ha ⁻¹)	Net return (Rs ha ⁻¹)	B: C ratio
T ₁	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE	38004	87228	49224	1.29
T ₂	Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE	36905	89863	52958	1.43
T ₃	Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE	36584	90915	54331	1.48
T ₄	Propaquizafop 10 EC @ 0.075 kg a.i./ha PoE	36976	84595	47619	1.28
T ₅	Quizalfop ethyl 5 EC @ 40 g a.i./ha PoE	36944	86177	49233	1.33

T ₆	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Imazethapyr 10% SL @ 0.06 kg a.i./ha PoE	38885	103543	64658	1.66
T ₇	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Propaquizafop 10 EC @ 0.075 kg a.i./ha PoE	38956	100382	61426	1.57
T ₈	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha PoE	38564	104595	66031	1.71
T ₉	Pendimethalin 30 EC @ 1.0 kg a.i./ha PE <i>fb</i> by Quizalfop ethyl 5 EC @ 40 g a.i./ha PoE	38924	99342	60418	1.55
T ₁₀	Weed free	46024	108803	69779	1.51
T ₁₁	Weedy check	36024	72985	36961	1.02

Conclusion

The study on effect of different weed management practices in chickpea, revealed that application of pendimethalin @ 1.0 kg a.i./ha followed by post-emergence Clodinafoppropargyl 15 WP @ 0.06 kg a.i./ha as PoE proved to be the best way to control weed as it has recorded significantly less weed density, weed dry matter, higher weed control efficiency, yield attributes (number of pods plant⁻¹, number of seeds pod⁻¹ and test weight), and culminated in, yield (grain yield, stover yield, biological yield, and harvest index). Economically it also recorded higher gross returns, net return, and benefit cost ratio.

References

- Anonymous (2020). Area and production of principal crops, *Division of Economics and statistics, Krishi Bhawan*, New Delhi.
- Bhutada, P.O. and Bhale, V.M. (2014). Effect of herbicides and cultural practices on nutrient uptake by chickpea and weed. *International Journal Forestry & Crop Improvement* 5(1): 13-15.
- Chavada, J.N., Patel, S.P., Patel, S.B., Panchal, P.P. and Patel, G.N. (2018). Weed management in chickpea (*Cicer arietinum*) under north Gujarat conditions. *International Journal of Science environment and Technology* 6(3): 2278-3687.

- Dubey, S.K., Kumar, A., Singh, D., Partap, T., Chourasia, A. (2018). Effect of Different Weed Control Measures on Performance of Chickpea under Irrigated Condition. *International Journal of Current Microbiology and Applied Sciences* **7**(5): 3103-3111.
- Dungarwal, H.S., Chaplot, P.C. and Nagda, B.L. (2002). Chemical weed control in chickpea. *Indian Journal of Weed Science* **34**(3&4): 208-212.
- Gomez, K. A. and Gomez, A. A. (1984). *Statistical Procedures for Agriculture Research, 2nd edition, John Wiley and Sons, New York.*
- Jaswal, A. and Menon, S. (2020). Review of literature on effect of various herbicidal treatments on chickpea (*Cicer arietinum*. L) Intercropping with lentil (*Lens culinaris* Medik.) *International Journal of Chemical Studies* **8**(5): 1116-1121.
- Kachhadiya, S. P., Savaliya, J. J., Bhalu, V. B., Pansuriya, A. G and Savaliya, S. G. (2009). Evaluation of new herbicides for weed management in chickpea (*Cicer arietinum* L.).
- Khope, D., Kumar, D. and Pannu, R.K. (2011). Evaluation of Post-emergence Herbicides in Chickpea (*Cicer arietinum*). *Indian Journal of Weed Science* **43**(1&2): 92-93.
- Kumar, K.K. Hazr, S.L., Yadav and Singh, S.S. (2015). Weed dynamics and productivity of chickpea (*Cicer arietinum* L.) under pre- and post-emergence application of herbicides. *Indian Journal of Agronomy* **60**(4): 570-575.
- Kumar, N., Kumar, S., Sharma, S. (2020) Efficacy of Different Herbicides on Yield and Nutrient Uptake of Chickpea (*Cicer arietinum* L.). *International Archive of Applied Sciences and Technology* **11**(3): 91-97.
- Kumar, N., Nandal, D.P and Punia., S.S. (2014). Weed management in Chickpea under irrigated condition. *Indian Journal of Weed Science* **46** (3): 300-301.
- Marwat, K. B., Khan, I. A., Gul, H., and Naqibullah, K. (2003). Efficacy of different pre-and post- emergence herbicides for controlling weeds in chickpea. *Pakistan Journal of Weed Science Research* **10**(1/2): 51-54.
- Meena, D. S., Ram, B. Jadon, C and Tetarwal, J. P. (2011). Efficacy of Imazethapyr on weed management in soybean. *Indian Journal of Weed Science* **43**(3&4): 169-171.
- Naidu, (2012). Identification of weeds, *Directorate of Weed Science Research Jabalpur (M.P).*
- Nasir, M., & Sidhu, J. S. (2012). 12 Common Pulses: Chickpea, Lentil, Mungbean, Black Gram, Pigeon Pea and Indian Vetch. *Dry beans and pulses: Production, processing and nutrition*, 283.
- Poonia, T C. and Pithia M. S. (2013). Pre- and post-emergence herbicides for weed management in chickpea *Indian Journal of Weed Science* **45**(3): 223-225.
- Punia, S. S., Samunder Singh and Dharambir Yadav, (2011). Bio-efficacy of Imazathapyr and Chlorimuron-ethyl in clusterbean and the residual effect on succeeding Rabi crop. *Indian Journal of Weed Science* **43**(1&2):48-53.

- Rao, T. K. S., Chauhan, I. S., Fulsoundar, A. B., Gamit, V. V., & Parveen, K. (2014). Improving comfort and welfare to mitigate stress in dairy animals-a review. *Wayamba Journal of Animal Science*, 6, 1070-1084.
- Rathod, P. S., Patil, D. H and Dodamani, B. M., (2017). Evaluation of time and dose of imazethapyr in controlling weeds of chickpea (*Cicer arietinum* L.). *Legume Research- An International Journal* 40(5): 906-910.
- Ratnam, M., Rao, A.S and Reddy, T.Y. (2011). Integrated weed management in chickpea (*Cicer arietinum* L.). *Indian Journal Weed Science* 43:70-72.
- Singh, A. and Jain, N. (2017). Integrated weed management in chickpea. *Indian Journal of Weed Science* 49(1): 93-94.
- Singh, A. K., Chovatia, P. K., Kathiria, R. K. and Savaliya, N. V. (2019). Effect of integrated nutrient management on growth, yield and economics of chickpea (*Cicer arietinum* L.). *International Journal of Chemical Studies* 7(3): 3048-3050.
- Singh, D., Pazhanisamy, S., Kumar, S., Kumar, A. and Reddy, S.A. (2020). Bio- efficacy of different herbicides in broad spectrum weed management for chickpea. *International Journal of Current Microbiology and Applied Sciences* 9(3): 2313-2317.
- Singh, K.R., Shukla, D.N. and Nirmal, D. (2006). Effect of biofertilizers, fertility level and weed management on weed growth and yield of late sown chickpea (*Cicer arietinum* L.). *Indian Journal of Weed Science* 76(9): 561-563.
- Singh, R. P., Verma, S. K., Singh, R. K. and Idnani, L. K., (2014). Influence of sowing dates and weed management on weed growth and nutrients depletion by weeds and uptake by chickpea (*Cicer arietinum* L.) under rainfed condition. *Indian Journal of Agricultural Sciences* 84(4) 468-472.
- Vaishya, R.D., Fayaz, M. and Shrivastava, A.K. (1996). Integrated weed management in chickpea. *Indian Journal of Pulses Research* 9(1): 34-38.
- Yadav, S.L., Chahal, V.P., Yadav, A., Singh, R.K., Mishra, D., Kumar, P. and Kumar, R. (2017). Response of phosphorus doses and weed control methods on nutrient uptake in chickpea (*Cicer arietinum*) under rainfed condition. *Journal of Pharmacognosy and Phytochemistry* 1: 738-742.
- Yaduraju, N. T. Herbicide resistant crops in weed management. In: *The extended Summaries, Golden Jubilee National Symposium on Conservation Agriculture and Environment*. October 26-28, Banaras Hindu University, Varanasi. (2006) 297-98.
- Peerzada, A. M., Ali, H. H., & Chauhan, B. S. (2017). Weed management in sorghum [*Sorghum bicolor* (L.) Moench] using crop competition: a review. *Crop Protection*, 95, 74-80.
- Dubey, S. K., Kumar, A., Singh, D., Partap, T., & Chaurasiya, A. (2018). Effect of different weed control measures on performance of chickpea under irrigated condition. *International Journal of Current Microbiology and Applied Sciences*, 7(5), 3103-3111.

Singh, V. K., Shukla, A. K., Dwivedi, B. S., Singh, M. P., Majumdar, K., Kumar, V., ... & Singh, S. K. (2015). Site-specific nutrient management under rice-based cropping systems in Indo-Gangetic Plains: yield, profit and apparent nutrient balance. *Agricultural Research*, 4(4), 365-377.