

**Induction of Apoptosis in Ehrlich ascites carcinoma cells through an intrinsic pathway by Co(II)-benzoin Thiosemicarbazone Complex [Co(BTSC)<sub>2</sub>]**

**Abstract**

Cancer is a class of disease in which a group of cells display uncontrolled growth, invasion and sometimes metastasis. Cancer recognized as the second leading cause of death globally. Therefore, the discovery and development of new potent and selective anticancer drugs are of high importance in modern cancer research. The objective of this study was to find out the mechanism through which Co(II)-benzoin thiosemicarbazone complex exerts its antitumor activity against Ehrlich ascites carcinoma (EAC) cells bearing Swiss albino mice. Induction of apoptosis in EAC cells was confirmed by observation of nuclear morphology and DNA fragmentation assay. The intrinsic apoptotic pathway induced by Co(BTSC)<sub>2</sub> was evidenced by the mRNA expression levels of several tumor related antiapoptotic genes B-cell lymphoma 2 (bcl-2), B-cell lymphoma extra-large (bcl-xL) and caspase-8, and proapoptotic genes p53 or tumor protein, bcl-2 associated X protein (bax), caspase-9, caspase-3 and poly-ADP ribose polymerase (PARP-1) and *in vitro* effect of caspase inhibitors on EAC cells. Reactive oxygen species (ROS) generation after Co(BTSC)<sub>2</sub> treatment was determined by 2', 7'-dichlorodihydrofluorescein diacetate (DCFH-DA) staining. The results found during this study, confirmed that the induction of apoptosis by Co(BTSC)<sub>2</sub> occurred through an ROS-dependent mitochondria-mediated intrinsic pathway rather than an extrinsic pathway, and was regulated by the bcl-2 protein family. Thus, this study provides evidence to carry out further researches in a way to formulate novel anticancer drugs.

**Keywords**

EAC cells, Cobalt benzoin thiosemicarbazone complex, intrinsic pathway, ROS, Caspase inhibitor.

**Introduction**

Schiff bases are condensation products of aldehyde and ketones with primary amines and were first reported by Hugo Schiff in 1864. These are the compounds containing imine or azomethine (-HC=N-) functional group and are found to be a versatile pharmacophore for design and development of various bioactive lead compounds. Schiff bases as well as schiff base complexes

with transition metals form an important class of the most widely used organic and organometallic compounds and have a wide variety of applications in many fields including analytical, biological and inorganic chemistry. In recent times Schiff bases and Schiff base metal complexes have drawn the attention of many researchers in medicinal and pharmaceutical fields due to a broad spectrum of biological activities like anti-inflammatory, analgesic, antimicrobial, anticonvulsant, antitubercular, antioxidant, anthelmintic, anti HIV, antileprosy, antileukemia, anticancer and so on. Several reports have been published as the successful use of such class of compounds as anticancer agents [1-5]. However, further investigation is necessary for biological studies in order to understand more about their antitumor mechanism(s).

Apoptosis is a process of programmed cell death regulated by cellular signaling pathways and characterized morphologically by cell shrinkage, nuclear fragmentation, chromatin condensation, chromosomal DNA fragmentation and global mRNA decay. Apoptosis can be initiated through one of two pathways, one is intrinsic pathway and other is the extrinsic pathway [6]. The intrinsic (mitochondrial) pathway is induced by cellular stress, infection and DNA damage and results in the activation of pro-apoptotic BCL-2 associated X (BAX) or BCL-2 homologous killer (Bak) proteins oligomerize to create holes or pores on the mitochondrial outer membrane and release cytochrome c, to the cytosol [7]. The another pathway is extrinsic (death receptor) pathway initiated by death ligand such as tumor necrosis factor  $\alpha$  (TNF  $\alpha$ ) or TNF-related apoptosis inducing ligand (TRAIL) binding to cell surface receptors and activate caspase-8, which then activate downstream caspase-3 directly which play essential role in programmed cell death [8]. These two pathways are executed mainly by caspases a family of endoproteases that provide critical links in cell regulatory networks controlling inflammation and cell death with caspase-8 and -9 engaging in the extrinsic and intrinsic pathways, respectively [9-10]. In addition, reactive oxygen species (ROS), a series of oxygen metabolism byproducts, have a role in cell signaling including; apoptosis; gene expression; and the activation of cell signaling cascades [11]. In this present paper we have reported anticancer activities of Cobalt (II) benzoin thiosemicarbazone complex against EAC cells in Swiss albino mice and the mechanism of apoptosis.

## Materials and Methods

## **Chemicals**

All chemicals and reagents used to carry out the research work were of reagent grade.

## **Experimental animal model**


Adult male Swiss albino mice, 5–7 weeks old ( $25 \pm 4$  g body weight), were collected from animal resource branch of the International Centre for Diarrheal Disease Research (ICDDR'B), Mohakhali, Dhaka, Bangladesh. Animals were housed in iron cages containing sterile paddy husk and saw dust as bedding material under hygienic conditions with a maximum of six animals in a cage. They were maintained under controlled conditions (12:12 h light–dark with temperature  $22 \pm 5$  °C). Standard mouse diet (recommended and prepared by International Centre for Diarrheal Disease Research, Bangladesh Mohakhali, Dhaka) and water were given in adequate.

Ethical approval was obtained from institutional animal, medical ethics, biosafety and biosecurity committee for experimentations on animal, human, microbes, and living natural sources from Institute of Biological Sciences, University of Rajshahi, Bangladesh (225/320-IAMEBBC/I B.Sc).

## **Synthesis of the compounds**

### **Synthesis of Co(II)-benzoin thiosemicarbazone complex**

The compound was synthesized according to the method as described in the literature [12-13] For benzoin thiosemicarbazone (BTSC), benzoin and thiosemicarbazide (1:1 molar ratio) were mixed together and refluxed for a period of 3-4 hours and then distilled to half of the total volume. A saturated solution of metal (II) acetate [Co(II)] in ethanol was added to the condensed solution. Within a few minutes crystals of Co(II)-benzoin thiosemicarbazone was obtained. The crystals was washed with ethanol and recrystallized twice, dried in an oven at 50°C and stored in a desiccator.



**Figure- 1:** Synthesis of Co(II)-benzoin thiosemicarbazone complex

### Cell lines

Ehrlich ascites carcinoma (EAC) cells were obtained by the courtesy of Indian Institute of Chemical Biology (IICB), Kolkata, India. The cells were maintained as mammary gland cancer cell in ascites in Swiss albino mice by intraperitoneal inoculation (biweekly) of  $2 \times 10^6$  cells/mouse. For *in vitro* study, EAC cells were cultured in Roswell Park Memorial Institute (RPMI)-1640 medium having glucose, 2 mM L-glutamine in presence of 10% fetal calf serum, and 1% (v/v) penicillin-streptomycin in a humidified atmosphere of 5%  $\text{CO}_2$  at 37 °C.

### *In vitro* Cell growth inhibition

*In vitro* effect of  $\text{Co}(\text{BTSC})_2$  against EAC cell growth inhibition was assayed by MTT colorimetric technique. NAD(P)H-dependent cellular oxidoreductase enzymes may, under defined conditions, reflect the number of viable cells present. These enzymes are capable of

reducing the tetrazolium dye MTT [3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide] to its insoluble formazan, which has a purple color. For this experiment,  $4 \times 10^6$  EAC cells in 200  $\mu\text{L}$  RPMI-1640 media were plated in the 96-well flat bottom culture plate in presence of six different concentrations (5  $\mu\text{g}$ , 10  $\mu\text{g}$ , 20  $\mu\text{g}$ , 40  $\mu\text{g}$ , 80  $\mu\text{g}$  and 120  $\mu\text{g}/\text{mL}$ ) of  $\text{Co}(\text{BTSC})_2$  and incubated for 24 h at 37 °C in  $\text{CO}_2$  incubator. EAC cells treated with dimethyl sulfoxide (DMSO), was used as control. The assays were performed in triplicates to avoid experimental errors. After the incubation period, aliquot were removed carefully, and 180  $\mu\text{L}$  of phosphate buffered saline (PBS) and 20  $\mu\text{L}$  of MTT were added to each well and further incubated for 8 h at 37 °C. The aliquot was removed again and 200  $\mu\text{L}$  of acidic isopropanol was added and incubated again at 37 °C for 1 h. Subsequently, absorbance was taken at 570 nm in microtiter plate reader (Optica Microplate Reader, Mikura Ltd., Horsham, UK). The following equation was used to calculate the cell proliferation inhibition ratio:

$$\text{Proliferation inhibition ratio (\%)} = \frac{(A - B) \times 100}{A}$$

Where A is the  $\text{OD}_{570 \text{ nm}}$  of the cellular homogenate (control) without experimental compounds and B is the  $\text{OD}_{570 \text{ nm}}$  of the cellular homogenate with experimental compounds.

### **DNA fragmentation assay**

DNA fragmentation assay by agarose gel electrophoresis was determined by the method described previously [14]. EAC cells were collected from mice treated with experimental compounds  $\text{Co}(\text{BTSC})_2$  at the dose of 8mg/kg/mice (five consecutive days) and from control mice. The cells were washed with PBS and resuspended again in PBS. The genomic DNA was isolated by using a DNA extraction kit (Promega, Madison, WI, USA) and analyzed by electrophoresis on 1.5% agarose gel containing 0.1 $\mu\text{g}/\text{mL}$  ethidium bromide and visualized under Ultraviolet illuminator.

### **Effect of caspase inhibitors on EAC cells**

In order to confirm the involvement of caspases [15] in the experimental compounds that induced cell death, the untreated EAC cells were collected on day six of tumor inoculation and washed with PBS. Then these cells were incubated in  $\text{CO}_2$  incubator (RPMI-1064 media) with Z-DEVD-FMK (caspase-3 inhibition, 2 $\mu\text{mol}/\text{mL}$ ) and Z-IETD-FMK (caspase-8 inhibitor,

2 $\mu$ mol/mL) for 2h. Then the cells were treated with Co(BTSC)<sub>2</sub> and incubated for another 24 hours at 37<sup>0</sup>C in CO<sub>2</sub> incubator. Finally, the cells were counted using hemocytometer and determined the effects of experimental compounds on cell growth inhibition in presence or absence of specific inhibitor.

### **Measurement of Reactive Oxygen Species (ROS) generation**

The ROS level of the cells was examined using 2', 7'-dichlorodihydrofluorescein diacetate (DCFH-DA; Invitrogen, USA). The cells were collected from control mice and treated mice on day six of tumor inoculation and washed with PBS at 1200rpm for 2minutes. Then the cells were incubated with DCFH-DA 10 $\mu$ L (50 $\mu$ M final concentration) at 37<sup>0</sup>C for 30minutes in the dark. Again the cells were washed twice with PBS and maintained in 1mL culture medium. ROS generation was assessed using a fluorescence microscope at excitation and emission wavelengths of 485nm and 530nm, respectively [16].

### **Reverse transcription polymerase chain reaction**

Total RNA was extracted using TRIzol method from mice receiving Co(BTSC)<sub>2</sub> at the dose of 8mg/kg/day and control EAC bearing mice on day six of tumor implantation. Total RNA was used as a template for reverse transcription using the following protocol: each 20 $\mu$ L reaction contained 125 $\mu$ M deoxynucleotide (dNTP), 100 pmol random hexamer, 100 pmol oligo dT<sub>18</sub> primer, 50 units of MuLV reverse transcriptase ( New England Bio lab, Ipswich, MA, USA), diethylpyrocarbonate (DEPC) treated water and 3 $\mu$ g total RNA. Briefly, RNA and oligo dT<sub>18</sub> primer were incubated at 70<sup>0</sup>C for 15 minutes then immediately placed on ice, after which the other components were added and incubated at 42<sup>0</sup>C for 1h and then at 70<sup>0</sup>C for 15 minutes. Expression of one housekeeping gene (*gapdh*) and eight growth regulatory genes, namely, *bcl-2*, *bax*, *p53*, *bcl-xl*, *parp 1* and Caspase-3,-8,-9 were examined using these cDNA as template for PCR. Reaction mixture were prepared containing 1X of Taq polymerase, 25pmol each of forward and reverse primer, 2.5mM of each dNTP and 0.25U of platinum Tag polymerase (Tiangen, Beijing, China), 2 $\mu$ L cDNA and nuclease-free water to a total volume of 25 $\mu$ L. Primer sequences are listed in Table 8. BioRad (Hercules CA, USA) gradient thermal cycler was used for amplifications. All the PCR products were analyzed in 1.5% agarose gel and Gene Rular 1kb DNA ladder (Fermentus, Pittsburgh, PA, USA) was used as marker. The cycling condition for initial PCR activation step of 3 min at 95 <sup>0</sup>C, followed by 35 cycles of 95 <sup>0</sup>C /1 min. For p53,

Bcl-2 and Caspase-3, -8, -9 genes 55 °C/1 min, 72 °C/1 min and a final extension of 72 °C/1 min. In the case of Bax, GAPDH and Bcl-xL, the annealing temperature was 54 °C.

### **Statistical analysis**

The experimental results have been expressed as the mean  $\pm$  S.E.M. Data have been calculated by one-way ANOVA followed by Dunnett "t" test for the determination of statistical significance using SPSS software of 20 version. The difference was considered to be statistically significant when  $p < 0.05$ .

### **Results and Discussion**

#### ***In vitro* EAC cell growth inhibition with Co(BTSC)<sub>2</sub>**

*In vitro* effect of Co(BTSC)<sub>2</sub> on EAC cell growth inhibition have been investigated by MTT assay. The Co(BTSC)<sub>2</sub> induced EAC cell death is a dose dependent manner (Figure 2). The Co(BTSC)<sub>2</sub> showed 16.2%, 32.3%, 45.7%, 61.62%, 73.9% and 89.72% cell growth inhibition at the concentration of 5,10,20,40,80 and 120  $\mu$ g/mL respectively. The effect decreased with the reduction in concentration of Co(BTSC)<sub>2</sub>.

#### **DNA fragmentation assay**

The activation of the endogenous Ca<sup>2+</sup>/Mg<sup>2+</sup> dependent endonuclease is the most distinctive biochemical hallmark of apoptosis. This activated endonuclease mediated the cleavage of internucleosomes and genera oligonucleotide fragments of about 180-200 base pairs length or their polymers. A DNA band was obtained from agarose gel when genomic DNA from the experimental compound Co(BTSC)<sub>2</sub> treated EAC cells was undergone PCR amplification followed by agarose gel electrophoresis. This DNA showed the characteristic feature of apoptosis induction. On the other hand, control group showed smear-like DNA degradation. Result was shown in figure 3.

#### **Effect of caspases on Co(BTSC)<sub>2</sub>-induced cytotoxicity in EAC cells**

Caspase inhibitors Z-DEVD-FMK (caspase-3 inhibitor) and Z-IETD-FMK (caspase-8 inhibitor) were used to detect the involvement of specific caspases in the apoptotic cell death of EAC cells induced by the treatment of experimental compound Co(BTSC)<sub>2</sub>. In the presence of Co(BTSC)<sub>2</sub> growth inhibition of EAC cells was 78.89% and it decreased to 3.88 and 75.01% in presence of Z-DEVD-FMK (caspase-3 inhibitor) and Z-IETD-FMK (caspase-8 inhibitor) respectively in the culture medium (figure 4).

#### **Altered expressions of cancer-related genes in Co(BTSC)<sub>2</sub> treated EAC cells**

Reverse transcription PCR was used to study the mRNA expression levels of several tumor-related antiapoptotic genes (*bcl-2*, *bcl-xl* and caspase 8) and proapoptotic genes (*p53*, *bax*, *parp-1*, caspase-3 and caspase-9) in control and Co(BTSC)<sub>2</sub>-treated EAC cells (Figure 5). *gapdh* primers were used for an amplification reaction to confirm the suitability of the purified RNAs and the samples were found to be suitable for PCR. The control cells showed high expressions of *bcl-2*, *bcl-xl* and caspase 8 genes, whereas EAC cells treated with experimental compounds showed reduced *bcl-2*, *bcl-xl* and caspase 8 mRNA expressions. In addition, the *p53*, *bax*, *parp-1*, caspase-3 and caspase-9 genes showed increased expressions in treated EAC cells. On the other hand, no expression of these genes was found in control mice.

#### **Determination of intracellular reactive oxygen system in EAC cells**

2, 7-dichlorodihydrofluorescein diacetate (DCFH-DA) is a common fluorescent probe that can be used for detecting several ROS in biological media. The DCFH-DA is firstly hydrolyzed to DCFH by intracellular esterase, which is then oxidized by reactive species and originates a fluorescent compound 2, 7- dichlorofluorescein (DCF), whose fluorescence intensity is proportional to the levels of ROS. Here, DCFH-DA was used to assess the effect of Co(BTSC)<sub>2</sub> on ROS generation in EAC cells after 24 h treatment. As shown in figure. 6, the untreated control cells displayed little green fluorescence, while increasing concentrations of the experimental compound resulted in much stronger signals, with 8 mg/kg (*i. p.*) the compounds producing the strongest intensity.

#### **Conclusion**

In the light of above observations the experimental compounds can be considered as effective antitumor agents and are comparable in efficiency with that of *bleomycin* at least in cell growth inhibition, reduction average tumor weight and enhancement of survival time of tumor bearing mice. These compounds may therefore be considered primarily as potent anticancer agents and the possible mechanism for the anticancer effect is ROS dependent mitochondria-mediated intrinsic pathway rather than an extrinsic pathway and was regulated by the BCL-2 protein family. In order to ascertain these compounds as novel potential anticancer drugs, it is necessary to carry out further experiments against other cancer cell lines with higher test animals and with advanced techniques. These findings definitely give positive support to carry out further researches in a way to formulate novel anticancer drugs.



## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.


## References

- [1]. Zakir HM, Mondol MM, Shahriar SM, Hossain MA, Rahim MA, Dey AK, Amin R, Rahman MZ, Chowdhury AN, Moniruzzaman M, Khanam J, et al. Benzoin Thiosemicarbazone Inhibits Growth and Triggers Apoptosis in Earlich Ascites Carcinoma Cells through an Intrinsic Pathway. *American Journal of Cancer*. 2019. 7(1):1-9.
- [2]. Ali SM, Zakir HM, Shahriar SM, Sarkar MH, Dey AK, Nur HP, Jesmin M, et al. In vivo anticancer activities of Ni (II)-Benzoin thiosemicarbazone complex [Ni (BTSC) 2] against ehrlich ascites carcinoma cells. *Journal of Bio-Science*. 2015. 23:77-88.
- [3]. Zakir HM, Mondol MM, Shahriar SM, Hossain MA, Rahim MA, Dey AK, Amin R, Rahman MZ, Chowdhury AN, Moniruzzaman M, Khanam J, et al. Benzoin Thiosemicarbazone Inhibits Growth and Triggers Apoptosis in Earlich Ascites Carcinoma Cells through an Intrinsic Pathway. *American Journal of Cancer*. 2019. 7(1):1-9.
- [4]. Zakir HM, Ali MM, Jesmin M, et al. Antibacterial activities of benzoin thiosemicarbazone and its complexes with Co (II) and Ni (II), et al. *Asian Journal of Medical and Pharmaceutical Researches*. 2016. 6(4):32-40.
- [5]. Zakir HM, Islam AM, Nurtaj K, Ali MM, Jesmin M, et al. Pesticidal activities of benzoin thiosemicarbazone and its complexes with Co (II) and Ni (II). *International Journal of Biosciences*. 2016. 9(6):92-100.
- [6]. Bai L, Wang S, et al. Targeting apoptosis pathways for new cancer therapeutics. *Annu Rev Med*. 2014. 65:139-155.
- [7]. Tait SW, Green DR, et al. Mitochondria and cell death; outer membrane permeabilization and beyond. *NRMCB*. 2010. 11:621-632.


- [8]. Wallach D, Kang TB, Kovalenko A, et al. The extrinsic cell death pathway and the elanmortel. *Cell death Differ.* 2008. 15:1533-1541.
- [9]. Ola MS, Nawaz M, Ahsan H, et al. Role of Bcl-2 family proteins and caspases in the regulation of apoptosis. *Mol Cell Biochem.* 2011. 351:41-58.
- [10]. McIlwain DR, Berger T, Mak WT, et al. Caspase functions in cell death and disease. *Cold spring harb perspect biol.* 2013.1; 5(4): a008656.
- [11]. Hancock JT, Desikan R, Neill SJ, et al. Role of reactive oxygen species in cell signaling pathways. *Biochem Soc Trans.* 2001.29:345-350.
- [12]. Longman, *Elementary Practical Organic Chemistry Part-2: Qualitative Organic Analysis.* 2<sup>nd</sup> ed. London. 1974; 118.
- [13]. El-Shahawi MS, Al-Jahdali MS, Bashammakh AS, Al-Sibaai AA, Nassef HM, et al. Spectroscopic and electrochemical characterization of some Schiff base metal complexes containing benzoin moiety. *Acta Part A: Mol and Biom Spectroscopy.* 2013. 113:459–465.
- [14]. Islam F, Khatun H, Khatun M, Ali MM, Khanam JA, et al. Growth inhibition and apoptosis of Ehrlich ascites carcinoma cells by the methanol extract of *Eucalyptus camaldulensis*. *Pharm Biol.* 2014. 52:281–290.
- [15]. Yinyuan W, Dianjun W, Xiaodong W, Yinyin W, Fangli R, Donald C, Zhijie C, Baoqing J, et al. Caspase-3 is activated through caspase-8 instead of caspase-9 during H<sub>2</sub>O<sub>2</sub>-induced apoptosis in HeLa Cells. *Cell Physical Biochemistry.* 2011. 27:539-546.
- [16]. Xiao-Dong W, Chen-Yang L, Miao-Miao J, Dong L, Ping W, Xun S, Jun-Da C, Li-Xuan G, Xiao-Peng H, Guo-Qiang L, Jian Z, Chun-Hua W, Zhen-Dan H, et al. Induction of apoptosis in human leukemia cells through an intrinsic pathway by cathachunine, a unique alkaloid isolation from *Catharanthus roseus*. *Phytomedicine.* 2016. 23:641-653.

**Table- 1:** Primer sequences and PCR protocol used for the RT-PCR assay


Sl. No.	Target genes	Primers	Amplification (kb)
1	<i>bcl-2</i> F <i>bcl-2</i> R	5'-GTGGAGGAGCTCTTCAGGGA-3' 5'-AGGCACCCAGGGTGATGCAA-3'	0.200
2	<i>bax</i> F <i>bax</i> R	5'-CGCCCACCAGCTCTGAGCAGA-3' 5'-GCCACGTGGGCGTCCCAAAGT-3'	0.500
3	<i>p<sup>53</sup></i> F <i>p<sup>53</sup></i> R	5'-GCGTCTTAGAGACAGTTGACT-3' 5'-GGATAGGTCGGCGGTTTCATGC-3'	0.550
4	<i>bcl-x</i> F <i>bcl-x</i> R	5'-TTGGACAATGGACTGGTTGA-3' 5'-GTAGAGTGGATGGTCAGTG-3'	0.700
5	<i>caspase-3</i> F <i>caspase-3</i> R	5'-TTAATAAAGGTATCCATGGAGAACACT-3' 5'-TTAGTGATAAAAATAGAGTTCTTTTGT-3'	0.300
6	<i>caspase-8</i> F <i>caspase-8</i> R	5'-CTGCTGGGGATGGCCACTGTG-3' 5'-TCGCCTCGAGGACATCGCTCTC-3'	0.450
7	<i>caspase-9</i> F <i>caspase-9</i> R	5'-ATGGACGAAGCGGATCGG-3' 5'-CCCTGGCCTTATGATGTT-3'	0.400
8	<i>parp-1</i> F <i>parp-1</i> R	5'-AGGCCCTAAAGGCTCAGAAT-3' 5'-CTAGGTTTCTGTGTCTTGAC-3'	0.270
9	<i>gapdh</i> F <i>gapdh</i> R	5'-GTGGAAGGACTCATGACCACAG-3' 5'-CTGGTGCTCAGTGTAGCCCAG-3'	0.350




**Figure-2.** *In vitro* effect of  $\text{Co}(\text{BTSC})_2$  on EAC cell growth inhibition. Data are expressed as the mean of results in 3 repeated experiments. Error bar represent standard error of mean.




**Figure-3:** DNA fragmentation assay in  $\text{Co}(\text{BTSC})_2$ -treated mice




**Figure- 4:** Effect of caspases on Co(BTSC)<sub>2</sub>- induced cytotoxicity in EAC cells. Numbers of mice in each experiment were 6. The results are shown in mean±SEM. Significant value is, \*\*p<0.01.



**Figure-5:** Expressions of different cancer-related genes in Co(BTSC)<sub>2</sub>-treated EAC cells. RNA was extracted from the experimental mice on day 6 and level of mRNA expression of antiapoptotic genes (Bcl-2, Bcl-xL and caspase-8) and proapoptotic genes (p53, Bax, Caspase-3, -9 and PARP-1) were studied. In Ni(BTSC)<sub>2</sub> treated mice, all the proapoptotic genes expression increased remarkably, and the expression of antiapoptotic genes were not observed. In control group, opposite results were observed.





**Figure-6:** Effect of  $\text{Co}(\text{BTSC})_2$  on intracellular ROS generation Effects of  $\text{Co}(\text{BTSC})_2$  on the generation of ROS in EAC cells after 24 h treatment. Observation of cells by fluorescence microscopy (200 $\times$  magnification) after incubation with DCFH-DA. (A) Control cells, (B) 2 mg/kg (*i.p.*), (C) 4 mg/kg (*i.p.*) and (D) 8 mg/kg (*i.p.*).