

Original Research Article

PHYTOCHEMICAL PROFILE AND ANTIOXIDANT POTENTIAL OF A FORMULATED HERBAL PRODUCT FROM *SORGHUM BICOLOR* LINN., *CURCUMA LONGA* LINN., *BRIDELIA FERRUGINEA* BENTH AND HONEY

Abstract

The use of plants in the treatment of human diseases is as old as human itself. Over the years, phytomedicine has gained more attention in both developed and developing countries. This study evaluates the phytochemicals and the antioxidant properties of a formulated herbal product (FHP) made from the leaf sheath of *Sorghum bicolor*, rhizome of *Curcuma longa*, stem bark of *Bridelia ferruginea* and honey. The ethanolic and the aqueous extracts of the different components and the mixture were evaluated. The chemical compositions of the extracts were determined via gas chromatography-mass spectrometry (GC-MS) analysis and High Performance Liquid Chromatography (HP-LC) at the same time, the antioxidant properties were investigated using the ferric reducing antioxidant power (FRAP). The results of the phytochemical screening of FHP and the different components used for its preparation showed the presence of alkaloids, phenols, saponins, tannins and flavonoids irrespective of the extractant used. However, the values were higher in the aqueous extracts than the ethanolic extracts for most of the phytochemicals. For instance, higher values of alkaloids (9.57 ± 0.01 mg/g) and saponins (7.35 ± 0.01 mg/g) were observed in the aqueous extract of FHP while higher values of phenol (0.98 ± 0.01 mg/g), tannins (0.68 ± 0.00 mg/g) and flavonoids (40.54 ± 0.05 mg/g) were observed in the aqueous extract of *Curcuma longa*. Concerning antioxidant potential, the highest FRAP power was observed in the stem bark of *B. ferruginea* (1956.05 ± 15.54 mgAAE/g). This was closely followed by the FRAP value observed in *S. bicolor* (1359.89 ± 27.20 mgAAE/g). A total of 8 compounds was identified using High performance Liquid chromatography. These compounds include Gallic acid, Catechin, Ferrulic acid, Rutin, Quercetin, P-coumaric acid, Kaempferol and Apigenin. Most of these chemicals are known to possess health beneficial and therapeutic effects. It is conceivable therefore that FHP might exert significant therapeutic effects when used to treat specific debilitating medical conditions.

Key words: Formulated herbal product, phytochemicals, antioxidant property, organic compounds

Introduction

“Many plants referred to as medicinal plants are used as the raw material for many herbal formulations and supplements. The use of herbal medicines has been on the rise in recent years

due to their low price” [1]. About 80% of the people living in developed countries use traditional medicine [2]. Plants are used by humans as phytomedicine because of their bioactive constituents [3] “The active constituents of some plants such as flavonoids, polyphenols, alkaloids, terpenoids, saponins, enzymes, and tannins are widely used for prevention and treatment of many diseases. Nowadays, there is a great interest in drugs from natural origin with minimal side effects as against synthetic ones” [4, 5, 6]

Sorghum (*Sorghum bicolor* Linn.) is a grain cultivated across the world as food crop most especially in America and Africa. Evidence has shown that sorghum content phenolic acids, polyphenols, luteolinidin and apigenidin which are good antioxidants [7]. “This contributes to its lipid peroxidation inhibitory properties observed during mashing and boiling. The leaf sheath has a different chemical composition than the leaf blade” [8]. “The *S. bicolor* leaf sheaths have a strong chemoprotective potential and inhibit proliferation of gastrointestinal cancer cell lines, and these effects are partially independent of their antioxidant range” [9]. “The anticancer properties of sorghum are due in part to the high content of 3-deoxyanthocyanidins” [9, 10].” In addition to the high content of anti-inflammatory phenolic compounds, *S. bicolor* leaf contains several groups of bioactive compounds with the capacity to induce proinflammatory immune responses. Water-soluble beta-glucans found in sorghum are biologically active and are capable of initiating macrophage activation” [11]. According to Ali *et al.* [12], the therapeutic roles reported on sorghum are as a result of anti-inflammatory, anti-carcinogenic [9, 10] antibiotic, antifungal, antiviral, hepatoprotective, anti-ulcer, anti-neoplastic, cholesterol-lowering and digestibility showing properties. Such pharmaceutical functions are allied to the phytochemical contents of the plant such as phytosterols, policosanols, saponins, carotenoids and phenolic compounds, including flavonoids, tannins, and anthocyanins [13] “Different important flavonoids and phenolic acids identified in the leaves extract of sorghum include gallic acid, catechin, chlorogenic acid, caffeic acid, ellagic acid, rutin, isoquercitrin, quercitrin, quercetin, and kaempferol” [14].

“*Curcuma longa* Linn. (Turmeric) is a tropical plant of South, South-eastern Asia, and primarily found grown in tropical regions of Bangladesh, China, Thailand, Cambodia, Malaysia, Indonesia, Phillipines and Nigeria” [15]. “*C. longa* has healthy influence on digestive system and enhances the mucin secretion in the digestive tract. Several actions of *C. longa* have been identified. It has antibacterial, antihelminthic, anticancer, antiparasitic, antiseptic, anti-oxidative [16], anti-inflammatory, anti-rheumatic, anti-tumour, anti-phlegmatic, antiviral activities”. Moreover, “it has astringent, aromatic and blood purifying ability. It can also clear skin blemishes and neutralize free radicals. Furthermore, it has hepatoprotective and nephroprotective properties against toxic agents-induced hepato-renal failure” [17, 18]. Also, “it has protective activity against cytotoxic, teratogenic, and neurodegenerative diseases [5], anticoagulant and possesses anti-HIV activity to combat AIDS” [3].

“*Bridelia ferruginea* Benth. (Euphorbiaceae) is a plant that is commonly found in Savannah regions” [19]. “Its bark extract has been used for the coagulation of milk and lime juice for the

formulation of a traditional gargle “Egun Efu”. It is reported of having potential for water treatment and chemo preventive potential” [20]. Several phenolic compounds isolated from *B. ferruginea* stem bark were found to display radical scavenging and xanthine oxidase inhibition activities, therefore supporting the application of *B. ferruginea* in traditional medicine [6] as an anthelmintic, antiamebic, antianaemic, antibacterial, anticonvulsant, antidiabetic, antidiarrhoeal, anti-inflammatory [21], antimicrobial [22], antiviral, hypoglycemic and for abdominal pain, cardiovascular, gynecological and sexual diseases [23] or anti-cancer [24]. Other reported activities of the bark extract include trypanocidal, molluscidal [25].

“Honey is a natural sweet viscous fluid produced by honeybees from the pollen and nectar of flowering plants or from the nectar of blossoms which honeybees collect and transform by combining with their salivary secretions and deposit, dehydrate and store in the honeycomb to ripen” [26]. Some honey substances are essential for human life such as sugars (the major sugar present in all the types of honey is fructose), proteins, vitamins, organic acids and minerals [27]. Honey has been reported to have antibacterial activity against diarrhoeagenic bacteria [28, 29] and immunostimulatory potential [30].

According to Alfa *et al.* [31] “traditionally used medicinal plants of ethno-pharmacological relevance can be a substantial source of drugs and thus are worthy of investigation for potential biomedicine development. Therefore, this study is aimed toward better understanding of the pharmacotherapy property of formulated herbal mixture from *Sorghum bicolor* Linn., *Curcuma longa* Linn., *Bridelia ferruginea* Benth and honey”.

MATERIALS AND METHODS

Sample Collection

The plants materials (*Sorghum bicolor*, *Curcuma longa* and *Bridelia ferruginea*) were bought from Oja-Oba in Ado Ekiti, Nigeria while honey was purchased from Pelumi Pure Natural Honey Farm in Ifaki Ekiti, Nigeria.

Plant identification was done at the herbarium at the School of Agriculture, Federal University of Technology, Akure, Nigeria.

Preparation of the Herbal Product

The plant materials were rinsed under running water and air dried at room temperature ($28 \pm 2^\circ\text{C}$) to constant weight before pulverizing into powder with electric blender (Hex Mokappi). Mixing ratio was determined according to Akande *et al.* [32], Haddad *et al.* [33] and Olarewaju *et al.* [34] as follows: *Sorghum bicolor*, (59.2%), *Curcuma longa* (18.5%), honey (19.2%), *Bridelia ferruginea* (3.1%).

Crude Extraction

Each powdered plant materials and honey were separately extracted with hot water and cold 70% ethanol. The filtrates were evaporated to dryness at 45°C using evaporating dish in a water bath. The crude extracts were then scrapped and stored in a sterile container in refrigerator [35].

Determination of Total Phenols

Total Phenolic compound was determined according to Tayyebe *et al.* [36] with a slight modification. An aliquot of 500 µL of the extract was transferred to a test tube, 500µL of the Folin-Ciocalteu solution was added, then 1 mL of sodium carbonate solution and 8 ml of distilled water were also added. The samples remain at 28±2°C for 30 mins. Absorbance was measured at 760nm spectrophotometrically. These assays were performed in duplicates and the total phenolic content was expressed as milligrams of tannic acid equivalents per gram of sample (mg TAE/g).

Determination of Tannins

“Casein (500 mg) was weighed and transferred into a 25 ml Erlenmeyer flask, 5 ml of the extract and 5 ml of distilled water were added. After 2h, the extracts were filtered into a 10 mL volumetric flasks and its volume was adjusted with distilled water. Then 500 µL of the extract was transferred to a test tube, 500µL of the Folin-Ciocalteu solution was added, then 1 mL of sodium carbonate solution and 8 ml of distilled water were also added. The samples remain at room temperature for 30 mins. Absorbance was measured at 760nm spectrophotometrically. These assays were performed in duplicates and total tannins content is expressed as milligrams of tannic acid equivalents per gram of sample. (mg TAE/g)” [36].

Determination of Flavonoids

“An aliquot of 500 µL of the extract was transferred to a test tube, 500 µL of acetic acid solution, 2 mL of pyridine solution, 1 ml of aluminium chloride solution and 6 ml of 80% methanol were also added. The samples remain at 28±2°C for 30 mins. Absorbance was measured at 420nm spectrophotometrically. The test was carried out in duplicates and the flavonoid content was expressed as milligrams of quercetin equivalents per gram of sample (mgQE/g) on dry weight basis using standard curve” [37].

Determination of Saponins

Two grams (2g) of the extract were weighed into a conical flask and 20ml of 20% aqueous ethanol was added. It was then heated over the water bath for 4 h with continuous stirring at 55°C. This mixture was then filtered and the residue was re-extracted with another 200ml of 20% ethanol. These combined extracts were reduced to 40ml over water bath at about 90°C. These concentrates were transferred into a 250ml separator funnel and 20ml of diethyl ether was added and shaken vigorously. The aqueous layer was recovered while the ether layer was discarded. The purification process was repeated and 60ml of n-butanol was added. The combined n-butanol extracts were washed twice with 10ml of 5% aqueous sodium chloride. The remaining

solution was heated in the water bath. After evaporation the samples were dried in the oven to a constant weight to give the saponins which was then calculated according to Obadoni and Ochuko [38].

Determination of Alkaloids

“Five grames (5g) of the sample w weighed into 250 ml beaker and 200ml of 10% acetic acid in ethanol was added, covered and allowed to stand for 4 h. This was then filtered and the extract was concentrated on a water bath to one-quarter of the original volume. Concentrated ammonium hydroxide was added dropwise to the extract until precipitation was completed. The whole solution was allowed to settle and the residue was washed with dilute ammonium hydroxide and filtered. The residue which is the alkaloids was dried and weighed” [39].

Ferric Reducing Antioxidant Power (FRAP) Assay

The reducing capacity of the sample extract and standards were determined using the method of Boulanouar *et al.* [27]. At different concentrations (50–250 $\mu\text{g/mL}$), 2.5 mL of samples or standard were mixed with PBS (2.5 mL, 0.2 M, pH 6.6) and potassium ferricyanide (2.5 mL, 1.0%). The obtained mixture was incubated at 50°C during 20 mins, TCA (2.5 mL, 10%) was added to the mixture. Afterward, 2.5 mL of this solution was mixed with distilled water (2.5 mL) and FeCl_3 (0.5 mL, 0.1%). The ferric/ferrous transformation was investigated, and the absorbance values were measured at 700 nm in a spectrophotometer.

GC-MS Analysis of FHP

A GC 6890N instrument from Agilent (Palo Alto, CA, USA) coupled with a mass detector (MS5973; Agilent) was used to analysis the sample. Experimental conditions of the GC-MS system were as follows: a DB 5MS column (30 m \times 0.25 mm, 0.25 μm film thickness) was used and the flow rate of the mobile phase (He) was set at 1 mL/min. In the GC part, temperature was kept at 35°C for 8 min and then increased to 60°C at 6°C/min intervals followed by 4°C/min to 160°C and 20°C/min to 200°C/min and was kept at 200°C for 1 min. Organic compounds were identified in Wiley’s NIST Mass Spectral Library if the obtained comparison scores were higher than 95%. Otherwise, fragmentation peaks of the compounds were evaluated, and the compounds were identified using the memory background for the identification of the compounds that appeared in GC-MS chromatograms. Contents of individual compounds in the sample were given in percent of the total compound in the sample. This was the standard way used to quantify the organic compounds present in FHP.

Determination of Drug probative and Drug-Likeness

The drug-likeness and properties of the suspected bioactive compounds of FHP was evaluated using online program Osiris Server [40].

Results

The present investigation revealed the presence of medicinally bioactive compounds in FHP. The phytochemical characteristics of FHP and the individual components of the product are presented in Tables 1 and 2. Alkaloids, phenols, saponins, tannins and flavonoids were found in the FHP and the different components used for the preparation of FHP irrespective of the extractant used. The highest values of alkaloids ($9.57 \pm 0.01 \text{ mg/g}$) and saponins ($7.35 \pm 0.01 \text{ mg/g}$) were observed in the aqueous extract of FHP while the highest values of phenol ($0.98 \pm 0.01 \text{ mg/g}$), tannins ($0.68 \pm 0.00 \text{ mg/g}$) and flavonoids ($40.54 \pm 0.05 \text{ mg/g}$) were observed in the aqueous extract of *C. longa*.

The antioxidant activities of the materials investigated revealed that *B. ferruginea* ($1956.05 \pm 15.54 \text{ mgAAE/g}$) has the highest Ferric Reducing Antioxidant capacity followed by *S. bicolor* ($1359.89 \pm 27.2 \text{ mgAAE/g}$) while FHP had the lowest (Table 3). The results of the GC-MS and HPLC spectra of the aqueous extraction of FHP are shown in Figures 1, 2 and 3 respectively while Tables 4 and 5 on the other hand show the drug properties and some selected bioactive compounds in the GC-MS and HPLC screening of FHP.

The chromatograph of the GCMS figure 1 shows the peaks that are representing twenty seven (27) compounds. Some peaks depict the same compound with different retention time and thus their area percentage compositions are additive. The most abundant of the compounds were Hexadecanoic acid, methyl ester / Pentadecanoic acid, 14-methyl-, methyl ester with percentage area (11.69%;) and retention-time (RT=37.083) followed by Naphthalene / 1H-Indene, 1-methylene- / Naphthalene with percentage area and retention-time (10.35%; RT=16.228), Benzene, (1-methyl-1-propenyl)-, (E)-/ Benzene, 2-ethenyl-1,3-dimethyl- (9.60%; RT=14.98), Phthalic acid, di(oct-3-yl) ester / Phthalic acid, hept-2-yl isohexyl ester Phthalic acid, cyclohexyl 2-pentyl ester (8.21%; RT=40.861), Benzene, 1-ethyl-2,4-dimethyl- / o-Cymene/ Benzene, 2-ethyl-1,4-dimethyl- (6.61%; RT=13.97), 9-Octadecenoic acid (Z)-, methyl ester/ 9-Octadecenoic acid, methyl ester (E)- / 11-Octadecenoic acid, methyl ester (5.20%; RT=38.359), 1-Phenyl-1-butene/ (E)-1-Phenyl-1-butene/ Benzene, 1-ethenyl-4-ethyl- (5.52%; RT=14.589), 1H-Indene, 2,3-dihydro-1,6-dimethyl-/ Benzene, 1-methyl-2-(1-methyl-2-propenyl)- (4.53%; RT=16.459), 2-Ethyl-2,3-dihydro-1H-indene/ Bicyclo[4.2.1]nona-2,4,7-triene, 7-ethyl-/ 1H-Indene, 1-ethyl-2,3-dihydro- (4.49%; RT=18.085), 1H-Indene, 2,3-dihydro-4,7-dimethyl-/ 2,2-Dimethylindene, 2,3-dihydro-/ 1H-Indene, 2,3-dihydro-1,6-dimethyl- (4.24%; RT=18.661), 1H-Indene, 2,3-dihydro-1,2-dimethyl-/ Naphthalene, 1,2,3,4-tetrahydro-5-methyl- (3.14%; RT=19.105), Benzene, 1-ethyl-2,3-dimethyl-/ o-Cymene / Benzene, 1-ethyl-2,4-dimethyl- (3.00%; RT=12.36), Benzene, 1-ethyl-4-(1-methylethyl)/ 1,5,6,7-Tetramethylbicyclo[3.2.0]hepta-2,6-diene/ 3,4-Dimethylcumene (2.54%; RT=17.354), Benzene, 1,2,4,5-tetramethyl- / Benzene, 1,2,3,4-tetramethyl- (2.54v%; RT=13.81), Benzene, 2-ethenyl-1,3,5-trimethyl / 2-Propenal, 3-(4-methylphenyl)- / Benzene, (2-methyl-1-butenyl)- (2.51%; RT=19.712), Benzene, 1-ethyl-2,4-dimethyl- / Benzene, 2-ethyl-1,3-dimethyl-/ Benzene, 1-ethyl-2,4-dimethyl- (2.49%; RT=12.62), Benzene, (1,2-dimethyl-1-propenyl)/ 1H-Indene, 2,3-dihydro-1,2-dimethyl-/ 1H-Indene, 2,3-dihydro-1,6-dimethyl- (2.43%;

RT=16.784), Benzene, 4-ethyl-1,2-dimethyl- / Benzene,1-ethyl-2,4-dimethyl- / Benzene, 1-ethyl-3,5-dimethyl- (1.56%; RT=11.62), Benzene, 1,2,4,5-tetramethyl- / Benzene,1,2,3,4-tetramethyl- / 1,3-Cyclopentadiene, 1,2,3,4-tetramethyl-5-methylene- (1.38%; RT=13.40), Cyclotetracosane / 1-Nonadecene / Trifluoroacetoxy hexadecane (1.28%; RT=37.63), Benzene, 1-methyl-4-(1-methylpropyl)- / Benzene, (1,1-dimethylpropyl)- / Bis[3,4-dimethylbenzyl]sulfone (1.06%; RT=16.071), Benzene, 1-ethenyl-4-ethyl- / Benzene, (2-methyl-1-propenyl)- / Indan, 1-methyl- (0.72%, RT=12.54), Naphthalene, 2-methyl- / Naphthalene, 1-methyl- (0.85%; RT=20.656), Cetene / 8-Heptadecene, 1-chloro- -Octadecanol (0.49%; RT=35.325), Benzene, 1-ethyl-3-methyl- / Benzene,1-ethyl-4-methyl- / Benzene,1-ethyl-3-methyl- (0.46%; RT=10.02), while the least abundant of the compounds identified in GCMS assay was Mesitylene/ Benzene, 1,2,3-trimethyl- (0.40%; RT=8.89)

Also, the HPLC assay of flavonoid compounds (figure 2) shows presences of P-coumaric acid with percentage area (46.21%) and retention-time (18.79), Gallic acid (14.73%; RT=4.51), Ferrulic acid (14.56%; RT=20.95), Catechin (6.25%; RT= 13.22), Apigenin (2.39% RT=30.19), Quercetin (2.92%; RT= 36.69), Kampferol (2.72 %; RT= 38.44), Rutin (2.03%; RT= 23.31), While compounds of the 3 peaks with percentage area and retention-time (0.28%; RT=19.96, 0.18%; RT=22.27, 0.17%; RT=32.05) were not found in the library

The HPLC assay of Alkaloid compounds (figure 3) shows Alpha-Tumerone (30.51%/ RT=3.70) as the most abundant compounds followed by Curcumin (29.74%; RT= 17.23) Borneol (10.3%; RT=7.97), Choline (9.74%; RT= 5.88), Quercetin (9.17%; RT=15.50), Demethoxycurcumin (1.99%; RT=19.40), Alpha-Phelland (1.91%; RT=10.50), Myrtenal (1.63%; RT=20.50), Cycloartenol (1.57%; RT=21.42), Longifoline (1.29%; RT=9.12), Epicatechin (1.05%; RT=7.233), Verbenone (1.07%; RT=11.30)

Discussion

The presence of phytochemicals in the individual components used for the preparation of FHP in this study agrees with the results of previous studies in which a myriad of phytochemicals were observed in *Sorghum bicolor* [32, 11], *C. longa* [41, 42, 3], *B. ferruginea* [23, 35, 20, 6] and honey [43].

The phytochemical profiles of the aqueous extract of FHP as compared to the ethanol extracted FHP showed higher values of all the phytochemicals investigated. This shows that aqueous extraction using hot water facilitated the release of the phytochemicals in FHP better than the use of ethanol and also in the release of the phytochemicals in the different components used for its preparation established that hot water is a better extractant. However, this is in contrast with the report of Arawande *et al.* [42] who observed high extractive value using ethanol and low using water. The differences in extractive value may be associated with the fact that the present study makes use of hot water. According to Chaves *et al.* [44], increase of water temperature produces dramatic changes in its physical-chemical properties therefore enhancing the solubility of the

analytes, breaking matrix-analyte interactions achieving a higher diffusion rate. Tena [45] also stated that with moderately elevated temperature, the polarity of water could be reduced considerably and can act as if ethanol or acetone were being employed. Specifically, the higher extractive value of water as compared to ethanol was observed in the yield of flavonoids and alkaloids in FHP and turmeric in this study. Highest phenolic content in *C. longa* as observed in this present study disagrees with the report of Mahomoodally *et al.* [6] who reported high phenolic content in stem bark of *B. ferruginea*.

According to Ogbonnia *et al.* [20], the antioxidant power of plant material is associated with the extractive value of flavonoids. It is employed to manage various disease states with different oxidative stress aetiologies. The antioxidant activities as shown in Table 3 revealed that *B. ferruginea* (1956.05±15.54mgAAE/g) has the highest Ferric Reducing Antioxidant capacity followed by *S. bicolor* (1359.89±27.2 mgAAE/g). This agrees with the previous study [46] while the observation that *S. bicolor* also has very high FRAC also corroborates the report of Kathleen *et al.* (2013) who stated that the antioxidant protection capacity of the leaf sheaths from *S. bicolor* is many-fold higher than that reported for cereal grains and vegetables

The most abundant compound identified in GC-MS analysis were Hexadecanoic acid, methyl ester/ Pentadecanoic acid, 14-methyl-, methyl ester and 1H-Indene, 1-methylene this was similar to the report of El-Damhougy *et al.*, [47] who also noted the compounds as the most abundant compound in *Callyspongia crassa*. Hexadecanoic acid, methyl ester and Pentadecanoic acid, 14-methyl-, methyl ester have been reported as possessing antioxidant, antifungal, hypocholesterolemic, nematocidal, antibiotic pesticide and lubricant activities, anticancer and hemolytic 5-alpha reductase inhibitors properties [47]. While 1H-Indene, 1-methylene- a fatty acid methyl ester is a natural product often found in *Streptomyces malaysiense*, *Streptomyces* antioxidants and possesses antifungal antibacterial and antifungal activity [48].

Phthalic acid, di(oct-3-yl) ester and Phthalic acid, hept-2-yl isohexyl ester have been reported of demonstration allelopathic, antimicrobial, insecticidal and Antifouling effect [49; 50]. o-Cymene and Benzene, 1-ethyl-2,4-dimethyl- have been reported as antibiotic, anti-anxiety treatment, anti-inflammatory, and potentially as a cancer treatment, as well as being a flavouring agent [51;52].

Also, 9-Octadecenoic acid (Z)-, methyl ester, 9-Octadecenoic acid, methyl ester, (E)-/ 11-Octadecenoic acid, methyl ester have been reported as Anti-inflammatory, antibacterial, antiandrogenic cancer preventive, dermatitogenic hypocholesterolemic, 5-alpha reductase inhibitor, anemiagenic, insectifuge [53; 54.55].

The Most dominate compound identified in the HPLC assay includes; P-coumaric acid, Gallic acid, which have been reported of possessing antioxidant, anti-cancer, antimicrobial, promoted the recovery of hyperlipidemia antiviral, anti-inflammatory, antiplatelet aggregation, anxiolytic, antipyretic, analgesic, and anti-arthritis activities [56; 57; 58; 59; 60].

Catechin has also been shown to effectively inhibit influenza A (H1N1) virus infection and also has anti-hepatitis B virus activity [48], antibacterial, antifungal antioxidant, antidiabetic [61]

Alpha-Tumerone, Curcumin, Borneol, Ferrulic acid Kaempferol and Apigenin have been reported as anti-inflammatory, antioxidant, anticancer, antimutagenic, antimicrobial, antiobesity, hypolipidemic, cardioprotective, and neuroprotective effects [11; 62].

The identification of curcumin Gallic acid and Catechin in this present result is similar to Subramanian *et al.*, [63] and Mahomoodally *et al.*, [6] report who also identified curcumin in *Curcuma longa* and Gallic acid and Catechin from *bridelia* respectively. Curcumin is a responsible for the yellow pigment in *cucuma lunga*, it is use as additive in food [65] and has been reported of possesses antiviral, antioxidant, anti-carcinogenic, wound healing, anti-inflammatory, anti-diabetic, anti-stress and properties [63]

Furthermore, rutin one of the compounds identified in this present study has also been reported to have strong inhibitory effect on the SARS-CoV-2 main protease (Mpro) [65] while ferrulic acid and quercetin have been reported to have a strong ant-viral potential [66, 67].

Table 1: Quantitative phytochemical profile of ethanolic extract of FHP and individual components

Sample Codes	ALKALOIDS (mg/g)	PHENOLS (mg/g)	SAPONINS (mg/g)	TANNINS (mg/g)	FLAVONOIDS (mg/g)
A (<i>B. ferruginea</i>)	5.09±0.03	0.79±0.01	2.44±0.11	0.30±0.02	7.57±0.05
B(<i>S. bicolor</i>)	0.35±0.01	0.74±0.01	0.58± 0.04	0.63±0.02	26.56±0.27
C (<i>C. longa</i>)	0.32±0.01	0.76±0.01	0.04±0.02	0.63±0.03	33.15±0.02
D(Honey)	0.596±0.02	0.26±0.02	5.94±0.02	0.26±0.03	0.34±0.02
FHP	0.32±0.01	0.26±0.03	0.02±0.004	0.14±0.04	3.17±0.45

Table presents mean and mean deviation

Table 2: Quantitative phytochemical profile of aqueous extract of FHP and individual components

Sample Codes	Alkaloid (mg/g)	Phenols (mg/g)	Saponins (mg/g)	Tannins (mg/g)	Flavonoids (mg/g)
A (<i>B. ferruginea</i>)	5.26±0.00	0.80±0.00	1.95±0.00	0.32±0.00	7.92±0.00
B(<i>S. bicolor</i>)	0.48±0.00	0.86±0.00	0.90±0.02	0.53±0.00	28.16±0.04
C (<i>C. longa</i>)	0.43±0.00	0.98±0.01	0.22±0.00	0.68±0.00	40.54±0.05
D (Honey)	0.485±0.01	0.33±0.03	0.11±0.01	0.18±0.01	2.935±10.2
FHP	9.57±0.01	0.59±0.01	7.35±0.01	0.21±0.01	23.78±0.08

Table presents mean and mean deviation

Table 3: Ferric Reducing Antioxidant Power (FRAP) of FHP and individual components

Sample	FRAP (mgAAE/g))
A (<i>B. ferruginea</i>)	1956.05±15.54
B(<i>S. bicolor</i>)	1359.89±27.20
C (<i>C. longa</i>)	485.03±13.41
FHP	359.89±3.88

Key: AAE means Ascorbic acid equivalent




Fig. 1: Gas Chromatograph of FHP




Figure 2: High Performance Liquid Chromatograph of flavonoid compound of FHP




Figure 3: High Performance Liquid Chromatograph of Alkaloids compound of FHP

Table 4: Drug properties of the identified compounds from GC-MS of FHP

S/N	Compound name	Drug likeness	Mutagenic	Tumorigenic	Irritability	Reproductive effect	Drug Score
1	O-Cymene	-2.5088	None	none	none	None	0.47
2	Sulfone	-0.165	None	none	none	None	0.50
3	1H-Indene	3.9929	None	none	none	None	0.49
4	Benzene, 1-ethyl-3-methyl-	-2.8	None	None	None	None	0.48
5	Benzene, 1-ethenyl-4-ethyl-	-4.99	None	None	Medium	None	0.35
6	Benzene, (2-methyl-1-propenyl)-	-3.18	None	None	None	None	0.45
7	Indan, 1-methyl-	-1.82	None	None	None	None	0.50
8	(E)-1-Phenyl-1-butene	-6.1	None	None	None	None	0.44
9	2-Ethyl-2,3-dihydro-1H-indene	-0.79	None	None	None	None	0.57
10	Bicyclo[4.2.1]nona-2,4,7-triene	-0.84	None	None	None	None	0.62

Table 5: Drug properties of the identified compounds in HPLC of the Flavonoid of FHP

Compound name	Drug likeness	Mutagenic	Tumorigenic	Irritability	Reproductive effect	Drug Score
Gallic acid	0.12	High	None	None	High	0.27
Catechin	1.92	None	None	None	None	0.87
P-coumaric acid	0.58	None	None	None	High	0.47
Ferrulic acid	1.12	High	High	None	High	0.18
Rutin	3.31	None	None	None	None	0.57
Apigenin	1.21	High	None	None	None	0.47
Quercetin	1.6	High	High	None	None	0.3
Kampferol	0.9	High	None	None	None	0.48
Hexadecanoic acid, methyl ester	-35.36	None	None	None	None	0.24
o-Cymene	-2.42	None	None	None	None	0.47
Benzene, 1-ethyl-2,4-dimethyl-	-5.26	Moderate	Moderate	Moderate	High	0.17
Borneol	-3.53	None	None	None	None	0.49

CONCLUSION

This study has shown that the formulated herbal product prepared by our team has bioactive constituents such as Catechin, P-coumaric acid, Ferrulic acid, Rutin, Apigenin, Quercetin and

Gallic acid. The implication of this is that the formulated herbal product has good medicinal potential.

Significance of the study

The search for new drugs is on the increase because most of the available drugs are becoming non-effective as a result of overuse and drug resistance by most pathogenic microorganisms. Different plants have long been exploited in folklore medicine therefore researching into plants as source of discovery of new drugs has gained so more attention not only in the developing countries but also in the developed countries. This present study focuses on screening a formulated herbal product (FHP) made from the leaf sheath of *Sorghum bicolor*, rhizome of *Curcuma longa*, stem bark of *Bridelia ferruginea* and honey for phytochemicals and antioxidants. The results of the phytochemical screening of FHP revealed the presence of alkaloids, phenols, saponins, tannins, flavonoids and some organic compounds which are known to possess health and therapeutic effects such as Gallic acid, Catechin, Ferrulic acid, Rutin, Quercetin, P-coumaric acid, Kampferol and Apigenin. It is conceivable therefore that FHP might exert significant therapeutic effects when used to treat specific debilitating medical conditions.

REFERENCES

1. Sawant RS. and Godghate AG. Qualitative Phytochemical Screening of Rhizomes of *Curcuma longa* Linn. *International Journal of Science, Environment and Technology*. 2013; 2(4):634 – 641
2. Pawar MA. Phytochemical and Physicochemical Investigation of *Curcuma longa* Linn. Rhizome. *International Journal of Chemical and Physical Sciences*. 2015; 4 (special issue) NCSC.
3. Sayantani, C. and Ramachandra, T.V. Phytochemical and Pharmacological Importance of Turmeric (*Curcuma longa*): A Review. *Research & Reviews: A Journal of Pharmacology*. 2019; 9 (1):16-23 ISSN: 2230-9861, ISSN: 2349-1299
4. Neamat-Allah ANF, El-Murr A. and Abd El-Hakim Y. Dietary supplementation with low molecular weight sodium alginate improves growth, haematology, immune reactions and resistance against *Aeromonas hydrophila* in *Clarias gariepinus*. *Aquac. Res*. 2019a; 50: 1547–1556. <https://doi.org/10.1111/are.14031>
5. Amany A M A, Nasr A M, Nasr E.D, Heba I A. and Ahmed N F. Effect of the aqueous root extract of *Curcuma longa* L. (turmeric) against thermally oxidized oil-induced hematological, biochemical and histopathological alterations. *Comparative clinical Pathology*. 2020; <https://doi.org/10.1007/s00580-020-03108-w> Springer-Verlag London Ltd., part of Springer Nature
6. Mahomoodally MF., Jugreet S, Sinan K.I, Zengin G Ak G, Ceylan R, Jek'O J, Cziáky Z, Angelini P, Angeles FG, et al. Pharmacological Potential and Chemical Characterization of

- Bridelia ferruginea* Benth. - A Native Tropical African Medicinal Plant. *Antibiotics*. 2021; 10: 223. <https://doi.org/10.3390/antibiotics10020223>
7. Oyinbo CA, Robert FO, Avwioro OG. and Igbigbi PS. Jobelyn suppresses hippocampal neuronal apoptosis and necrosis in experimental alcohol-induced brain stress. *Pathophysiology*. 2018; 25(4): 317-325.
 8. Adetokunbo OT, Adedoyin OD, Irete OA, Adeniyi A, Olufemi AO, Akinsegun AA, Bodunrin, IO, Samira BL.M. An open-label, randomized, parallel-group comparative study of the efficacy of *Sorghum bicolor* extract in preoperative anaemia. *Nutrition*. 2017; 33: 113-117.
 9. Phi-Hung N, Bing TZ, Jeong HL, Young HK, Byung SM, Mi HW. Isolation of benzoic and cinnamic acid derivatives from the grains of *Sorghum bicolor* and their inhibition of lipopolysaccharide-induced nitric oxide production in RAW 264.7 cells. *Food Chemistry*. 2015; 168: 512-519.
 10. Norma JSL, Guadalupe LP, Rocío CV, Marcela GM, Eduardo MS J, Marina E B, Gustavo AG, Maribel RS. The Extrusion Process as an Alternative for Improving the Biological Potential of *Sorghum* Bran: Phenolic Compounds and Antiradical and Anti-Inflammatory Capacity. *Evidence-Based Complementary and Alternative Medicine*. 2016; 1-8.
 11. Kathleen FB, Joni LB, Boxin O, Ademola O, Olajuwon O. and Gitte S J. West African *Sorghum bicolor* Leaf Sheaths Have Anti-Inflammatory and Immune-Modulating Properties *In Vitro*. *Journal of Medicinal Food*. 2013; 16 (3): 230–238
 12. Ali KS, Elsheikh AM, Ahmed HM, Hassan HA, Hamza NB, Osman MG. and Daffalla HM. Phytochemical screening and antibacterial activities of *Sorghum bicolor* leaves derived from *in vitro* culture. *GSC Biological and Pharmaceutical Sciences*. 2020; 10(1): 65-72.
 13. Salazar-López N.J, González-Aguilar G, Rouzaud-Sández O. and Robles-Sánchez M. Technologies applied to sorghum (*Sorghum bicolor* L. Moench): changes in phenolic compounds and antioxidant capacity. *Food Science Technology*. 2018; 38(3): 369-382.
 14. Borokini FB. Identification and quantification of polyphenols in *Sorghum bicolor* (L) Moench leaves extract using reverse-phase HPLC-DAD. *Clinical Experimental Pharmacology*. 2017; 7: 2.
 15. Abdulsalam AT, Adeniran AT, Adamu B S, Olaifa G O, Rita MS. and Memunetu, U. Phytochemical Properties, Proximate and Mineral Composition of *Curcuma longa* Linn. and *Zingiber officinale* Rosc.: A Comparative Study. *Journal of Scientific Research & Reports*. 2017; 13(4): 1-7, DOI: 10.9734/JSRR/2017/32623
 16. Hewlings SJ. and Kalman DS Curcumin: A Review of Its'Effects on Human Health. *Foods*. 2017; 6:6. <https://doi.org/10.3390/foods6100092>
 17. Farkhondeh T, Samarghandian S, Azimi-Nezhad M, Shahri AMP. Protective effects of curcumin against toxic agents-induced renal failure: a review. *Cardiovasc Hematol Disord Drug Targets*. 2018; <https://doi.org/10.2174/1871529x18666180905160830>
 18. Karamalakova YD, Nikolova GD, Georgiev TK., Gadjeva VG. and Tolekova AN. Hepatoprotective properties of *Curcuma longa* L. extract in bleomycin-induced chronic hepatotoxicity. *Drug Discovery Therapy*. 2019; 13: 9–16. <https://doi.org/10.5582/ddt.2018.01081>
 19. Ekanem, JT, Kolawole OM. and Abbah OC. Trypanocidal Potential of Ethanolic extracts of *Bridelia ferruginea* Benth bark in *Rattus norvegicus*. *African Journal of Biochemical Research*. 2008; 2(2):045-050
 20. Ogbonnia SO, Ginikachukwu O, Ezemenahi SI, Okeke AI and Ota D. Preliminary Phytochemistry, Antioxidant activities and GC/MS of the most abundant compounds of different

- solvents fractions of the Plant *Bridelia ferruginea* Benth used locally in the management of Diabetes. *Journal of Pharmacognosy and Phytochemistry*. 2021; 10(3): 154-164
21. Olajide, OA, Makinde J M and Awe SO. Effect of aqueous extract of *Bridelia ferruginea* stem bark on carrageenan induced edema and granuloma tissue formation on rats and mice, *Journal of Ethnopharmacology*. 1999; 66(1):113-117.
22. Ndukwe KC, Okeke I. N, Lamikanra A, Adesina SK. and Aboderin, O. Antibacterial Activity of Aqueous Extracts of selected chewing sticks. *J. Contemp. Dent. Pract.* 2005; 6(3): 086-094
23. Alfred M. Ethnopharmacology and Therapeutic Value of *Bridelia micrantha* (Hochst.) Baill. in Tropical Africa: A Comprehensive Review. *Molecules* 2017; 22: 1493; doi:10.3390/molecules22091493
24. Feng S, Cheng S, Yuan Z, Leitch M. and Xu CC. Valorization of bark for chemicals and materials: A review. *Renew. Sustain. Energy Rev.* 2013; 26: 560–578
25. Adeoye A.O, Abaeli A.M. and Owowumi, CJ. Antimicrobial activity of *Bridelia ferruginea*, In: Book of abstracts of the symposium on drug production from natural products Ile-Ife, 1988; 24
26. Rebiai A, Lanez T. and Belfar M L. Total polyphenol contents, radical scavenging and cyclic voltammetry of Algerian propolis. *Int J Pharm Sci.* 2014; 6(1):395-400
27. Boulanouar B, İlginç KT, Aytaç Güder, Ömür GÇ, Sevim ÇY, Sanaa, K. B, Mosad A. G. Chemical Composition and Biological Activities of Honeybee Products from Algeria. *J Appl Biotechnol Rep.* 2020; 7(2):93-103
28. Adebolu, T T. Effects of natural honey on local isolates of diarrhoea causing bacteria in Southwestern Nigeria. *African Journal of Biotechnology* 2005; 4(10): 1172 – 1174.
29. John-Isa J F, Adebolu, TT. and Oyetayo V O. Antibacterial Effects of Honey in Nigeria on Selected Diarrhoeagenic Bacteria. *South Asian Journal of Research in Microbiology* 2019a; 3(2): 1-11.
30. John-Isa J F, Adebolu, T T. and Oyetayo, V.O. Immunomodulatory Effects of Honey in Wistar Rats Infected with *Salmonella typhimurium*. *Asian Journal of Immunology* 2019b; 2(1): 1-8.
31. Alfa AA., Ayodeji OA, Teru GAD, Tijani KB. Studies on the Phytochemical Compounds in the Ethanolic Leaf Extract (ELE), Ethanolic Bark Extract (EBE) and Ethanolic Root Extract (ERE) of *Bridelia ferruginea* Benth (Euphorbiaceae). *Asian J. Biochem. Genet. Mol. Biol.* 2019; 2: 1–8.
32. Akande IS, Oseni AA. and Biobaku O.A. Effects of aqueous extract of *Sorghum bicolor* on hepatic, histological and haematological indices in rats. *Journal of Cell and Animal Biology* 2010; 4(9): 137-142, Available online at <http://www.academicjournals.org/JCAB> ISSN 1996-0867 ©2010 Academic Journals
33. Haddad E R A.Madeha N A. and Suad, MA. Bees' Honey Protects the Liver of Male Rats against Melamine Toxicity. *BioMed Research International* 2013; Volume 2013, Article ID 786051, 8 pages. <https://doi.org/10.1155/2013/786051>.
34. Olarewaju OI, Oloyede OI, Ojo OA. and Onikanni SA. Effects of aqueous extract of *Bridelia ferruginea* stem bark on some haematological parameters of albino rats. *Innovations in Pharmaceuticals and Pharmacotherapy*. 2013; 1 (2): 70-75.
35. Mela I L, Stanley CO, Vincent OO. and John A. Phytochemical screening and *in vitro* valuation of antibacterial activity of aqueous and ethanolic extracts of root and stem bark of *Bridelia ferruginea* Benth. (Euphorbiaceae). *Journal of Medicinal Plants Research* 2020; 14(1): 54-61, DOI: 10.5897/JMPR2019.6799

36. Tayyebe H, Sayyed AE, Sayyed S.M. and Mahdi, M. Total Phenolic Contents and Antioxidant Activities of Different Extracts and Fractions from the Aerial Parts of *Artemisia biennis* Willd. *Iran Journal of Pharmaceutical Research* 2014, 13(2): 551–559.
37. Elba LCA, Valériumc TNAC, Joabe GM, Allan JCC. and Tadeu, J.SPS. Standard Operating Procedures (SOP) for the Spectrophotometric Determination of Phenolic Compounds Contained in Plant Samples, Latest Research into Quality Control, Isin Akyar, Intech Open, 2012; DOI: 10.5772/51686. Available from: <https://www.intechopen.com/chapters/38925>. Assessed on 16/10/2021
38. Obadoni BO. and Ochuko PO. Phytochemical Studies and Comparative Efficacy of the Crude Extracts of Some Homeostatic Plants in Edo and Delta State of Nigeria. *Global Journal of Pure and Applied Science* 2001; 8: 203-208.
39. Edeoga HO, Okwu D E. and Mbaebie BO. Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnology* 2005; 4(7): 685-688. Available online at <http://www.academicjournals.org/AJB>
40. Thomas T. Osiris Compound property explorer. 2017; Available at <https://pubs.acs.org/doi/pdf/10.1021/jm00105a033>
41. Harshal P, Mugdha K, Nilesh M, Pravin J and Kavita M. Phytochemical Evaluation and Curcumin Content Determination of Turmeric Rhizomes Collected from Bhandara District of Maharashtra (India). *Medicinal Chemistry* 2014, 4(8): 588-591
42. Arawande J O, Akinnusotu A, & Alademeyin J.O. Extractive Value and Phytochemical Screening of Ginger (*Zingiber officinale*) and Turmeric (*Curcuma longa*) Using Different Solvents. *International Journal of Traditional and Natural Medicine* 2018, 8(1): 13-22 ISSN: 2167-1141.
43. Mbotto C I, Eja M E, Adegoke, A. A, Iwatt GD, Asikong BE, Takon I, Udo SM and Akeh M. Phytochemical properties and antimicrobial activities of combined effect of extracts of the leaves of *Garcinia kola*, *Vernonia amygdalina* and honey on some medically important microorganisms. *African Journal of Microbiology Research* 2009, 3(9): 557-559
44. Chaves J O, de Souza M C, da Silva L C, Lachos-Perez D, Torres-Mayanga P C, Machado A P da F, Forster-Carneiro T, Vázquez-Espinosa M, González-de-Peredo A V, Barbero G F, & Rostagno M A. Extraction of Flavonoids from Natural Sources Using Modern Techniques. *Frontiers in Chemistry*, <https://doi.org/10.3389/fchem.2020.507887> Available at: <https://www.frontiersin.org/articles/10.3389/fchem.2020.507887/full>
- 45 Tena MT. Pressurized liquid extraction reference module in chemistry. *Mol. Sci. Chem. Eng.* 2017, doi: 10.1016/B978-0-12-409547-2.14407-5
46. Omotade I O. and Seun OB. *In vitro* antioxidant activity of ethanolic extract of *Bridelia ferruginea* (stem bark). *Academic Research International*, 2012, 2(3), ISSN-L: 2223-9553, ISSN: 2223-9944
47. El-Damhougy K. A1, Mansour A.E. Bashar, Hussein A. E., Hassan A.H. I., and Fekry M.A.S. GC-MS Analysis of bioactive components of *Callyspongia crassa* (PORIFERA) from Gulf of Aqaba red sea (Egypt). *Al Azhar Bulletin of Science* Vol. 9th., Conf., March 2017, p. 111-11
48. National Center for Biotechnology Information (2022). PubChem Compound Summary for CID 21205, Methyl 14-methylpentadecanoate. Retrieved September 14, 2022 from <https://pubchem.ncbi.nlm.nih.gov/compound/Methyl-14-methylpentadecanoate>.

49. Huang, L.; Zhu, X.; Zhou, S.; Cheng, Z.; Shi, K.; Zhang, C.; Shao, H. Phthalic Acid Esters: Natural Sources and Biological Activities. *Toxins* 2021, 13, 495. <https://doi.org/10.3390/toxins13070495>
50. Ingole, S. N. (2016). Phytochemical analysis of leaf extract of *Ocimum americanum* L. (Lamiaceae) by GCMS method. 37, 76–87. www.worldscientificnews.com
51. Marchese A., Arciola C. R., Barbieri, R., Silva A. S., Nabavi S. F., Tsetegho Sokeng A. J., Izadi, M. Jafari N. J., Suntar I., Daglia M., & Nabavi S. M. Update on Monoterpenes as Antimicrobial Agents: A Particular Focus on p-Cymene. *Materials*, 2017; 10(8), 947. <https://doi.org/10.3390/ma10080947>
52. Elshafie, H. S., & Camele, I. (2017). An Overview of the Biological Effects of Some Mediterranean Essential Oils on Human Health. *BioMed Research International*, 2017, 1–14. <https://doi.org/10.1155/2017/9268468>
53. Salisu T. F , Okpuzor JE. and Jaja SI. Identification, characterization and quantification of chemical compounds in selected edible wild leafy vegetable. *Ife Journal of Science* 2019; vol. 21, no. 1. <https://dx.doi.org/10.4314/ijfs.v21i1.19>
54. Adeoye-Isijola MO, Olajuyigbe OO, Jonathan SG, Coopoosamy RM. Bioactive compounds in ethanol extract of *Lentinus squarrosulus* Mont-a Nigerian medicinal macrofungus. 2018; *African J Tradit Complement Altern Med* 15:42–50
55. Krishnamoorthy, K., & Subramaniam, P. (2014). Phytochemical Profiling of Leaf, Stem, and Tuber Parts of *Solena amplexicaulis* (Lam.) Gandhi Using GC-MS. *International Scholarly Research Notices*, 2014; 567409. <https://doi.org/10.1155/2014/567409>
56. Yang K, Zhang L, Liao P, Xiao Z, Zhang F, Sindaye D, Xin Z, Tan C, Deng J, Yin Y and Deng B. Impact of Gallic Acid on Gut Health: Focus on the Gut Microbiome, Immune Response, and Mechanisms of Action. *Front. Immunol.* 2020; 11:580208. doi: 10.3389/fimmu.2020.580208
57. Kahkeshani N, Farzaei F, Fotouhi M, Alavi SS, Bahramsoltani R, Naseri R, Momtaz S, Abbasabadi Z, Rahimi R, Farzaei MH, Bishayee A. Pharmacological effects of gallic acid in health and diseases: A mechanistic review. *Iran J Basic Med Sci.* 2019; 22(3):225-237. doi: 10.22038/ijbms.2019.32806.7897. PMID: 31156781; PMCID: PMC6528712.
58. Pei K, Ou J, Huang J, Ou S. p-Coumaric acid and its conjugates: dietary sources, pharmacokinetic properties and biological activities. 2016; *J Sci Food Agric.* 96(9):2952-62. doi: 10.1002/jsfa.7578. Epub 2016 Jan 25. PMID: 26692250.
59. Srinivasan M, Sudheer AR, Menon VP. Ferulic Acid: therapeutic potential through its antioxidant property. *J Clin Biochem Nutr.* 2007; 40(2):92-100. doi: 10.3164/jcfn.40.92. PMID: 18188410; PMCID: PMC2127228
60. Kang Y, Limeng Z, Pinfeng L, Zaili X, Fan Z, Daniel S, Zhongquan X, Chengquan Tan, Jinping D, Yulong Y. and Baichuan, D. Impact of Gallic Acid on Gut Health: Focus on the Gut Microbiome, Immune Response, and Mechanisms of Action. *Frontier in Immunology.* 2020, Vol. 11:580208. doi: 10.3389/fimmu.2020.580208
61. Chacko SM, Thambi PT, Kuttan R, Nishigaki I. Beneficial effects of green tea: a literature review. *Chin Med.* 2010; 6;5:13. doi: 10.1186/1749-8546-5-13. PMID: 20370896; PMCID: PMC2855614.
62. Singletary, Keith PhD. Turmeric: Potential Health Benefits. *Nutrition Today:* 1/2 2020 - Volume 55 - Issue 1 - p 45-56 doi: 10.1097/NT.0000000000000392

63. Subramanian A, Samuel T., Selvaraj D., Shameem R. and Grace M. GC-MS analysis of bioactive compounds of *Curcuma longa* Linnaeus (Zingiberaceae) rhizome extract. *Journal of Pharmacognosy and Phytochemistry* 2019; 8(6): 49-52
64. Rad J, Rayess YE, Rizk AA, Sadaka C, Zgheib R, Zam W, Sestito S, Rapposelli S, Neffe-Skocińska K, Zielińska D, Salehi B, Setzer WN, Dosoky NS, Taheri Y, El Beyrouthy M, Martorell M, Ostrander EA, Suleria HAR, Cho WC, Maroyi A and Martins N. Turmeric and Its Major Compound Curcumin on Health: Bioactive Effects and Safety Profiles for Food, Pharmaceutical, Biotechnological and Medicinal Applications. *Front. Pharmacol.* 2020; 11:01021. doi: 10.3389/fphar.2020.01021
65. You H, Huang C, Chen C, Chang C, Liao P. and Huang S. Anti-pandemic influenza A (H1N1) virus potential of catechin and gallic acid. *J Chin Med Assoc.* 2018, 81:458–68. doi: 10.1016/j.jcma.2017.11.00
66. Pawan, K.A., Chandan A. and Gerald, B. L.. Rutin: A Potential Antiviral for Repurposing as a SARS-CoV-2 Main Protease (Mpro) Inhibitor. *Natural Product Communications* 2021, 16(4): 1–12.
67. Xinyi Y, Xiaoli Z, Haiyan J, Junxiao D, Panpan L, Zhengtao J, Xian L, Yibo W, Chuqiao W, Jingya Z, Yanan W, Yangcheng Z, He Y. and Huanzhang, Z. Quercetin synergistically reactivates human immunodeficiency virus type 1 latency by activating nuclear factor κ B. *Molecular Medicine Reports* 2017, 17(2): 2501-2508. <https://doi.org/10.3892/mmr.2017.8188>