

Original Research Article

Castor seed oil: a promising biomitecicide for sustainable management of the red coffee mite (*Oligonychus ilicis*) (McGregor, 1917) (Prostigmata: Tetranychidae)

ABSTRACT

The coffee red mite [*Oligonychus ilicis* (McGregor, 1917)] (Prostigmata: Tetranychidae) is one of the main causes of conilon coffee (*Coffea canephora* Pierre & Froehn) crop damage due to its phytophagous habit. Nowadays, environment and human harmful synthetic pesticides are used to control this pest. In this context, plant-derived bioactive compounds have been studied as a sustainable alternative for the pest mite management in crops. The aim of this study was to evaluate the castor (*Ricinus communis*) seed oil action on eggs, larvae, nymph (protonymph and deutonymph) and adults of *O. ilicis*. Coffee leaf discs (4 cm in diameter) containing 12 individuals of *O. ilicis* were sprayed with castor seed oil at concentrations of 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0% (v/v) using airbrush. The ovicidal and mortality activity of this extract were evaluated against *O. ilicis*. The effective concentration of castor oil to cause 50% inhibition of *O. ilicis* larvae hatching (LC 50) was 1.26% (v/v). *Oligonychus ilicis* treated with this oil at a concentration of 3.0% presented a larvae hatching percentage of 29.3%, lower than that obtained for the control treatment (79.1%). Castor seed oil 3.0% (v/v) was highly toxic to nymphs and adults of *O. ilicis* with mortality of 96 and 88%, respectively. Castor seed oil was effective for larvae hatching inhibition, and nymphs and adults mortality of *O. ilicis*, being promising for the coffee red mite sustainable biocontrol.

Keywords: *Coffea canephora*, alternative control, mitecicide, biopesticide

1. INTRODUCTION

World coffee production exceeded 144.7 million bags in 2015, of which 41.5% were Conilon coffee produced in emerging countries, with Brazil being the second largest producer of this species in the world [1]. The coffee red mite, *Oligonychus ilicis* (McGregor, 1917) (Prostigmata: Tetranychidae), is one of the main phytophagous mites of the coffee tree (*Coffea* spp.), being one of the most important pests for the conilon coffee crop in Brazil [2]. Coffee tree red mite attack is favored by prolonged drought periods, affecting the entire crop if not controlled at the beginning of the infestation [3]. *Oligonychus ilicis* perforate the leaves cells and suck part of the cellular content to feed, causing reduction of a photosynthetic area of the leaves reduction, leaving them with a tanned appearance and covered with webs [2, 4].

Chemicals belonging to the aldicarb, bifenthrin, chlorpyrifos, pyrethroids and organophosphate groups are used to reduce mite damage in crops [5, 6, 7]. However, the use of these chemicals leads to the selection of red mite populations resistant to these products, pest resurgence, secondary pest outbreaks, and compromised performance of natural enemies [8, 9]. In addition, these pesticides are toxic to humans, being related to the development of diseases and effects such as tachycardia, hypertension, mydriasis, nausea, and vomiting occur, but more serious sequelae including respiratory failure, seizures and even death [10].

Alternative methods, such as the use of repellent biocompounds or insecticides/miticides derived from plants, have been developed for the management of mites and insects pests [11]. *Jatropha curcas* L. oil at 1.0% (m/v) and leaf and stem extracts from this plant at 3.0% (m/v) caused mortality of *Tetranychus urticae* among 82.6 and 88.5% [12]. *Argemone mexicana* flower extract effectively acted on the mortality of *Culex quinquefasciatus* Say (Diptera: Culicidae) with LC 50 and LC 90 of 18.61 and 39.86 ppm after 24 hours, and 9.47 and 21.76 ppm after 48 hours, respectively [13].

Castor (*Ricinus communis*) has been described for its insecticidal/miticidal capacity, presenting ricinine in its seed, a toxic alkaloid for mites and insects [14]. Castor seed oil was effective for controlling infestations of Mexican weevil *Zabrotes subfasciatus* (Boh.) and *Diaphania nitidalis* Cramer, 1782 (Lepidoptera: Crambidae) [15] (Mushobozy, Nganilevanu, Ruheza, & Swella, 2009). Although castor seed oil has been described as a promising biopesticide [15, 14, 16], there are no studies of this oil in the control of *O. ilicis*, the red coffee mite.

The objective of this work was to evaluate the ~~miticide~~ ~~miticidal~~ effect of castor seed oil on the control of different stages of development of the coffee red mite.

2. MATERIAL AND METHODS

The experiment was carried out at the Instituto Federal de Educação, Ciência e Tecnologia do Espírito Santo – Campus Itapina (IFES – Campus Itapina), at the Laboratory of Agricultural Entomology and Acarology, in the first half of 2018. The experimental units were kept in acclimatized chambers at a temperature of 25 ± 1 °C, relative humidity $70 \pm 10\%$ and a 12-hour photophase.

2.1. *Oligonychus ilicis* cultivation

Oligonychus ilicis adults were collected in the field, in a conventional conilon (*Coffea canephora*) coffee plantation, located in the Santo Antônio do Mutum heritage site at Sítio Renascer, Colatina, Espírito Santo State, Brazil.

The adopted creation technique was adapted from Reis et al. (1997) [17]. The coffee leaves were collected from chemical-free crops located at the Instituto Federal do Espírito Santo, Campus Itapina and washed with sodium dichloroisocyanurate and distilled water. Afterward, they were dried and placed on cotton in a Petri dish (14.0 x 1.5 cm), with the ~~leaves~~ ~~leaf~~ edges covered with moistened cotton to maintain the turgidity of the leaf and prevent the escape of mites.

After the procedure, the mites collected in the field were transferred to the plates and kept in acclimatized chambers (25 ± 1 °C, $70 \pm 10\%$ RH, and 12h photophase). Twenty adult females and 5 males of *O. ilicis* were removed from the breeding plates and transferred to Petri dishes as described above and kept for up to 24 hours to obtain eggs of the same age. This procedure was repeated for each development stage tested, with the same aim of age standardization. For each treatment, in each repetition, 12 individuals were left per plate.

2.2. Plant extracts production

Dry castor seeds were collected in an area free from chemical products at the Instituto Federal de Educação Ciência e Tecnologia do Espírito Santo (IFES) - Campus Itapina for vegetable oil production. In the laboratory, the castor seeds were peeled and then cold-pressed in a hydraulic press to extract the oil. For dilution, 3.0 mL of castor seed oil were added to 100 mL of distilled water and Tween® 80 adhesive spread (0.05%). Then, the mixture was stirred for 24 hours at room temperature on a magnetic stirrer. After this period, the material remained at rest for 20 minutes for decantation. After decantation, simple filtration was carried out in a cotton funnel to obtain a solid-free castor seed bean solution. The castor seed oil concentrations were 0.0, 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0 % (v/v) as described above.

2.3. Bioassays

Castor seed oil was tested at concentrations of 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0% (v/v) on eggs, larvae, nymphs (protonymphs and deutonymphs), and adults obtained as described above. Eight replicates per treatment were performed, with 12 individuals per repetition. Each experimental unit consisted of a Petri dish (10.0 x 1.2 cm) lined with cotton inside. On top of the cotton, 4 cm diameter coffee leaf discs were added, with cotton moistened around them to maintain the turgidity of the leaf and prevent the escape of mites.

The spraying was carried out using an Alpha 2 model airbrush, connected to a calibrated compressor with a constant pressure of 1.3 psi and 2mL of castor seed oil solution, each formulated for each repetition. Distilled water and the adhesive spreader Tween® 80 (0.05% v/v) mixture was used as a control. The ~~miticide-~~miticidal effect for eggs was evaluated from the 5th to the 7th day after application of the extract and for the other phases after 24, 48 and 72 hours after application.

The design used was completely randomized. Mortality data for each stage of development were corrected by Abbott's formula (1925) [18], and later submitted to variance and regression analysis tests by the SISVAR 5.6 program ($p \leq 0.05$). From the equations obtained, lethal concentration for killing 50% of the individuals or eggs (LC50) was calculated at each stage of development of *O. ilicis*. The differences in mortality in relation to time, for each concentration, in each stage, were submitted to analysis of variance and means were compared by the Tukey test ($p \leq 0.05$). The grouping of the toxicological effect was performed according to the model by Hassan et al. (1994) [19], when considering the percentage values of larvae hatching and corrected mortality, as follows: innocuous <25%; slightly toxic 25-50%; moderately toxic 51-75% and highly toxic >75%.

3. RESULTS AND DISCUSSION

3.1. Ovicidal effect of castor seed oil against *O. ilicis* eggs

The hatching percentage of *O. ilicis* eggs was inversely proportional to the concentration of castor seed oil (Figure 1).




Figure 1 - Larvae hatch percentage (A), larvae (B), nymphs (C), and adults (D) mortality of *Oligonychus ilicis* under different concentrations of castor seed oil (*Ricinus communis*), corrected by Abbott (1925) formula [18].

Treatments of *O. ilicis* eggs with castor seed oil at concentrations of 0.5, 1.0, 1.5, 2.0, and 2.5 and 3% (v/v) showed hatching of 74.2, 67.6, 53.1, 41.7, and 29.3%, respectively, while the control treatment showed 79.1% hatching (Figure 2).




Figure 2 - Larvae hatch percentage (A), larvae (B), nymphs (C), and adults (D) mortality of *Oligonychus ilicis* at 24, 48, and 72 hours after the application of castor (*Ricinus communis*) seed oil.

The castor seed oil effective concentration to cause 50% of larvae hatching of *O. ilicis* (LC 50) was 1.26 % (v/v). These results agree with those of *Tetranychus urticae* eggs sprayed with Azamax® (insecticide/natural miticide) showing viability inversely proportional to the concentrations of this product [20]. The increase in the Castor seed oil concentration inhibiting the hatching of larvae from *O. ilicis* eggs is justified by the presence of substances such as rhinoleic and oleic in this oil, described as toxic to oocytes of *Rhipicephalus sanguineus* (Latreille) and mosquito larvae, respectively [21].

Castor seed oil at 2.5 and 3.0% (v/v) was highly toxic to *O. ilicis* eggs, with corrected larvae hatching of 23.7 and 10.6%, respectively (Table 1).

Table 1: Castor oil toxicity at different stages of development of the coffee red mite (*Oligonychus ilicis*), with toxicological class of each concentration [19].

Development stage	Concentration	Larval Mortality	Hatch Average (%)	and Larval Hatch Mean and Corrected Mortality (%)	Toxicity
Larvae hatch	0	79.1		-	-
	0.5	74.2		67.4	Mild
	1.0	67.5		50.0	Mild
	1.5	53.1		40.8	Moderate
	2.0	41.7		26.3	Moderate
	2.5	39.6		23.7	High
	3.0	29.3		10.6	High
Larvae	0	4.6		-	-
	0.5	22.5		18.7	Innocuous
	1.0	40.4		37.6	Mild
	1.5	58.6		56.6	Moderate
	2.0	61.6		59.7	Moderate
	2.5	77.0		75.9	High
	3.0	75.2		74.0	Moderate
Nymphs	0	10.5		-	-
	0.5	39.6		32.5	Mild
	1.0	58.1		53.2	Moderate
	1.5	74.1		71.1	Moderate
	2.0	90.1		88.9	High
	2.5	89.4		87.8	High

	3.0	96.9	96.5	High
	0	9.3	-	-
	0.5	41.6	35.6	Mild
	1.0	57.6	53.2	Moderate
Adults	1.5	60.7	56.7	Moderate
	2.0	70.5	67.5	Moderate
	2.5	84.8	83.3	High
	3.0	88.1	86.9	High

The high toxicity of castor oil against *O. ilicis* eggs can be explained by its high concentration of ricinoleic acid, described as inhibiting the development of tick oocytes of the species *Rhipicephalus sanguineus* [22]. In addition, *Tetranychus* eggs present a respiration mechanism through embryonic stigmas connected to the chorion and a specialized region of the intermediate membrane containing perforations [23]. These perforations are a likely way of penetration of bioactive compounds, justifying the high ovicidal activity of *R. communis* in eggs of this mite.

Mites parasitizing rabbits fed on feed containing castor oil have been reported to have oocytes with massive numbers of vacuoles, resulting from the toxic effects of ingested ricinoleic acid esters [22], agreeing with the high toxicity of this oil against *O. ilicis* eggs. Possibly the reduction in the number of larvae hatched *O. ilicis* with increasing concentrations of castor oil is related to the characteristics of embryonic development of this mite.

3.2. Larvae, nymphs and adults *O. ilicis* mortality

The *O. ilicis* larvae, nymph and adult stages mortality was directly proportional to the castor seed oil concentration (Figure 1), in agreement with the report of *Jatropha curcas* L. (*Euphrobiaceae*) extracts applied on the striped mite, *Tetranychus urticae* Koch (Acari: *Tetranychidae*), causing dose-dependent mortality [12]. The high toxicity of castor oil against *O. ilicis* can be explained by the high ricinoleic acid concentration in this oil, described as toxic to *Anopheles culicifacies* larvae, the vector of malaria [24].

Castor oil at 1.5, 2.0 and 3.0% (v/v) for nymphs, and 2.5 and 3.0%, for adults, caused higher *O. ilicis* mortality in the first 24 hours after application of the product (Figure 2), agreeing with the report of high mortality of *Varroa destructor* in the first 24 hours of treatment with bifenthrin, fenpyroximate, fenazaquin, tebufenpyrad, chlorfenapyr and abamectin miticides [25].

The effective castor oil concentrations to cause 50% mortality (LC 50) of larvae, nymphs and adults of *O. ilicis* were 1.58, 0.88 and 1.07% (v v⁻¹), respectively, lower than that observed for newly emerged female *Hypothenemus hampei* (3.49% v v⁻¹) [26], confirming the high lethality of this oil against *O. ilicis*.

Castor seed oil was highly toxic to larvae, nymphs and adults of *O. ilicis* at a concentration of 2.5% (v/v), for the first, 2.0, 2.5 and 3.0%, for the second, and 2.5 and 3.0%, for the third, respectively (Table). Higher *O. ilicis* mortality at higher castor oil concentrations is justified by

greater cuticular penetration due to overload of the *O. ilicis* metabolic detoxification system [27]. The fact that castor oil was applied simultaneously on *O. ilicis* and its food (leaf disc) suggests that this mite was poisoned by three ways: ingestion, contact and respiratory tract, being the mode of action by contact faster than by ingestion [12, 28, 29].

Castor seed oil caused high mortality of eggs, nymphs and adults of *O. ilicis*, being a promising biocompound for the sustainable management of the red coffee mite.

4. CONCLUSION

The plant-based castor seed oil is ecologically correct and has a natural pesticide action. From this study, it can be concluded that castor seed oil is promising for the red coffee mite sustainable control due to its high efficiency against eggs, larvae, nymphs and adults of *O. ilicis* in laboratory assay.

Field tests applying castor seed oil for the *O. ilicis* control should be carried out in future works to confirm the efficiency of this oil for the red coffee mite management.

REFERENCES

1. Covre MA, Partelli FL, Bonomo R, Braun H, Ronchi CP. Vegetative growth of Conilon coffee plants under two water conditions in the Atlantic region of Bahia State, Brazil. *Acta Scientiarum. Agronomy*. 2016; 38 (4), 535-545. DOI: <https://doi.org/10.4025/actasciagron.v38i4.30627>
2. Silveira EC, Reis PR, Siqueira MF, Toledo MA, Liska GR, Cirillo MA. Functional response of *Euseius concordis* feeding on *Oligonychus ilicis* (Acari: Phytoseiidae, Tetranychidae). *Experimental and Applied Acarology*. 2020; 80:215–226. DOI: 10.1007/s10493-019-00454-5
3. Lopez L, Liburd OE. Injury to southern highbush blueberries by southern red mites and management using various miticides. *Insects*, 2020;11, 233. DOI:10.3390/insects11040233
4. Andrade DJ, Rocha CM, Matos STS, Zanardi OZ. Oxymatrine-based biomiteicide as a management tool against *Oligonychus ilicis* (McGregor) (Acari: Tetranychidae) in coffee. *Crop Protection*. 2020; 134, 105182. DOI: 10.1016/j.cropro.2020.105182
5. Hu J, Wang J, Yu Y, Rao W, Chen F, Wang C, Fan G. Cross-Resistance Pattern and Genetic Studies in Spirotetramat-Resistant Citrus Red Mite, *Panonychus citri* (Acari: Tetranychidae). *Agriculture*. 2022; 12(5):737. <https://doi.org/10.3390/agriculture12050737>
6. Herron GA, Rophail J. Potential insecticides for control of *Oligonychus ilicis* (McGregor) (Acari: Tetranychidae), a new threat to Australian horticulture. *Australian Journal of Entomology*. 2000;39(2), 86-88. DOI: 10.1046/j.1440-6055.2000.00153.x
7. Venzon M, Rosado MC, Fadini MAM, Ciociola Jr AI, Pallini A. The potential of Neem Azal for the control of coffee leaf pests. *Crop Protection*. 2005; 24(3), 213-219. DOI: 10.1016/j.cropro.2004.07.008
8. Cordeiro EMG, Moura ILT, Fadini MAM, Guedes RNC. Beyond selectivity: Are behavioral avoidance and hormesis likely causes of pyrethroid-induced outbreaks of the southern red mite *Oligonychus ilicis*? *Chemosphere*; 2013; 1111-1116. DOI: 10.1016/j.chemosphere.2013.06.030
9. Lima DB, Melo JWS, Gondim MGC, Guedes RNC, Oliveira JEM, Pallini A. Acaricide-impaired functional predation response of the phytoseiid mite *Neoseiulus baraki* to the coconut mite *Aceria guerreronis*. *Ecotoxicology*. 2015; 24:1124–1130. DOI: 10.1007/s10646-015-1459-z

10. Mohamed F, Gawarammana I, Robertson TA, Roberts MS, Palangasinghe C, Zawahir S, Jayamanne S, Kandasamy J, Eddleston M, Buckley NA, Dawson AH, Roberts DM. Acute Human Self-Poisoning with Imidacloprid Compound: A Neonicotinoid Insecticide. *PLoS ONE*. 2009; 4(4): e5127. DOI: 10.1371/journal.pone.0005127
11. Liang Z, Abdelshafy AM, Luo Z, Belwal T, Lin X, Xu Y, Wang L, Yang M, Qi M, Dong Y, Li L. Occurrence, detection, and dissipation of pesticide residue in plant-derived foodstuff: A state-of-the-art review. *Food Chemistry*. 2022; 132494.
12. Holtz AM, Rondelli VM, de Paulo HH, Franzin ML, Botti JMC, dos Santos VV, Pratisoli, D. Toxicity of physic nut extracts to two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae). *Magistra*. 2016; 28(1), 74-80. DOI: 0102-5333
13. Granados-Echegoyen CA, Chan-Bacab MJ, Ortega-Morales BO, Vásquez-López A, Lagunez-Rivera L, Diego-Nava F, Gaylarde C. *Argemone mexicana* (Papaverales: Papaveraceae) as an alternative for mosquito control: First report of larvicidal activity of flower extract. *Journal of Medical Entomology*. 2019; 56 (1), 261-267. doi: 10.1093/jme/tjy159.
14. Flores-Macías A, Vela-Correa G, Rodríguez-Gamiño M, Akhtar Y, Figueroa-Brito R, Pérez-Moreno V, Ramos-López MA. Effect of potassium nitrate on the production of ricinine by *Ricinus communis* and its insecticidal activity against *Spodoptera frugiperda*. *Revista fitotecnia Mexicana*. 2016; 39(1), 41-47. DOI: 0187-7380
15. Jadhav KB, Sawant MG, Satvekar T, Nagarkar JM. Biopesticide formulations of karanj and castor oil using soapnut. *Journal of Dispersion Science and Technology*. 2022; 43(3), 330-335. DOI: 10.1080/01932691.2020.1841001
16. Fragoso DFM, Túler AC, Pratisoli D, Carvalho JR, Valbon WR, de Queiroz VT, Pinheiro PF, Costa, A V, Bueno RCOF. Biological activity of plant extracts on the small tomato borer *Neoleucinodes elegantalis*, an important pest in the Neotropical region. *Crop Protection*. 2021; 145, 105606. DOI: <https://doi.org/10.1016/j.cropro.2021.105606>
17. Reis PR, Alves EB, Sousa EO. Biologia do ácaro-vermelho do cafeeiro *Oligonychus ilicis* (McGregor, 1917). *Ciência e Agrotecnologia*. 1997; 21(3), 260-266.
18. Abbott WS. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*. 1925 18(2), 265-267. DOI: 10.3109/13880209.2012.674950
19. Hassan SA, Bigler F, Bogenschütz H, Boller E, Brun J, Calis JNM, Vogt H. Results of the sixth joint pesticide testing programme of the IOBC/WPRS-working group «pesticides and beneficial organisms». *Entomophaga*. 1994; 39(1), 107-119. DOI:10.1007/BF02373500
20. Fernandes MHA, Souza ID, Oliveira AC, Paz HRR, Menezes KO, Oliveira JDM. Ação do extrato de nim (*Azadirachta indica*) sobre mortalidade de ovos de *Tetranychus urticae* Koch (Acari: Tetranychidae). In Embrapa Semiárido-Resumo em anais de congresso (ALICE). In: SIMPÓSIO DE CONTROLE BIOLÓGICO. Bonito. Faça bonito: use controle biológico: anais. Brasília, DF: Embrapa, 2013, 13.
21. Alonso EC, Santos DYAC. *Ricinus communis* and *Jatropha curcas* (Euphorbiaceae) Seed Oil Toxicity Against *Atta sexdens rubropilosa* (Hymenoptera: Formicidae). *Journal of Economic Entomology*. 2013; 106, 2, 742–746, <https://doi.org/10.1603/EC12035>
22. Arnosti A, Brienza PD, Furquim KCS, Chierice GO, Bechara GH, Calligaris IB, Camargo-Mathias MI. Effects of ricinoleic acid esters from castor oil of *Ricinus communis* on the vitello genesis of *Rhipicephalus sanguineus* (Latreille, 1806) (Acari: Ixodidae) ticks. *Experimental Parasitology*. 2011; 127(2), 575–580. DOI: 10.1016/j.exppara.2010.10.006

23. Gonçalves ME, Oliveira JV, Barros R, Torres JB. Effect of plant extracts on immature stages and adults females of *Mononychellus tanajoa* (Bondar) (Acari: Tetranychidae). *Neotropical Entomology*. 2001; 30(2), 305-309. DOI: 10.1590/S1519-566X2001000200015
24. Sogan N, Kapoor N, Kala S, Patanjali PK, Nagpal BN. Larvicidal activity of castor oil Nanoemulsion against malaria vector *Anopheles culicifacies*. *International Journal of Mosquito Research*. 2018; 5(3), 01–06. DOI: 2348-5906
25. Bahreini R, Nasr M, Docherty C, de Herdt O, Muirhead S, Feindel D. Evaluation of potential miticide toxicity to *Varroa destructor* and honey bees, *Apis mellifera*, under laboratory conditions. *Scientific reports*. 2020; 10(1), 1-14. DOI: 10.1038/s41598-020-78561-2
26. Calestino FN, Pratisoli D, Machado LC, Santos Junior HJG, de Queiroz VT, Mardgan L. Control of coffee berry borer, *Hypothenemus hampei* (Ferrari) (Coleoptera: Curculionidae: Scolytinae) with botanical insecticides and mineral oils. 2016; 38(1), 1-8. DOI: <https://doi.org/10.4025/actasciagron.v38i1.27430>
27. Van Leeuwen T, Vontas J, Tsagkarakou A, Dermauw W, Tirry L. Acaricide resistance mechanisms in the two-spotted spider mite *Tetranychus urticae* and other important Acari: a review. *Insect biochemistry and molecular biology*. 2010; 40(8), 563-572. DOI: 10.1016/j.ibmb.2010.05.008
28. Isman MB. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annual Review of Entomology*. 2006; 51, 45-66. DOI: 10.1146/annurev.ento.51.110104.151146
29. Ghani SBA, Al-Azzazy MM, Alhewairini SS, Al-Deghairi MA. The miticidal activity of silver nanoparticles towards date palm mite (*Oligonychus afrasiaticus* (McGregor)): efficacy, selectivity, and risk assessment. *Brazilian Journal of Biology*. 2024; 84, e261262. DOI: 10.1590/1519-6984.261262