

1
2
3
4
5
6
7
8
9

Variance of Arrowroot (*Maranta arundinacea*) Starch Granule Morphology Among Five Different Provinces in Sri Lanka

ABSTRACT

Arrowroot (*Maranta arundinacea*) is an underutilized tuber crop in Sri Lanka that produces a gluten-free, easily digestible starch. This research aimed to determine the variance of arrowroot starch granular morphology among the plants grown in five different provinces (Western, North-Western, Southern, Sabaragamuwa, Uva). Arrowroot starch granules were observed using the light microscope and scanning electron microscope. Oval, irregular globular and spherical shapes were the predominant granule shapes for arrowroot. The mean percentage of oval shaped granules ranged between 48.46 % - 59.34 %. The length and width of the granules were not significantly different among the five provinces. The length of the starch granules ranged between 42.91 - 45.86 μm while the width ranged between 30.81 – 32.32 μm . Arrowroot flour samples from five different provinces in Sri Lanka were not significantly different with regard to the starch granular morphology and therefore, arrowroot flour can be utilized in the local food industry without concerning their geographical locations.

Keywords: arrowroot, granular shape, granular morphology, Maranta arundinacea, starch

1. INTRODUCTION

The most widely available and cost-effective commodity product is starch, which is the second-largest source of biomass after cellulose. Glucose results from photosynthesis and is stored in chloroplasts of the storage organs of plants like tubers, roots, and rhizomes in the form of granules [1,2]. Starch is utilized in the food industry as a thickening agent, water retentive agent, gelling agent, and colloidal stabilizer to improve texture [3]. Baked products, sauces, soups, confectionaries, ice cream, sugar syrups, snacks, baby meals, and pharmaceuticals are produced from starch [4]. The semi-crystalline granular structure of starch is one of the main factors contributing to its economic competitiveness [5].

Since starch is made in granular form, extraction of starch through wet milling is more prevalent [6]. Amylose and amylopectin are the two main structural components of starch. Amylopectin 80–90% and amylose 10–20% are the most common compositions in starch. Physicochemical properties of starch, such as retrogradation, viscosity, and gel stability, are affected by the differences between these two molecules [1]. When combined with iodine solution, starch turns into blue colour. The linkage between glucose residues in starch is 1-4, while the linkage at branch points is 1-6 [7]. While amylose is amorphous, amylopectin in the granule has a semi-crystalline structure [6]. The hilum (the center of growth) is connected to the surface of the granules by a series of concentric layers known as growth rings of increasing diameter. The circumscribed growth rings themselves comprise shifting crystalline and amorphous zones of varying densities [8]. Depending on the type of plant, the amylose:amylopectin ratio in starch granule characteristics varies. Different botanical sources of starch have distinctive forms, sizes, and morphologies [9].

Starch granules range from submicron to more than one hundred microns. Granules of starch can be formed into spherical, disks, ovals, polygons, domes, elongated rods, and complex starches. The amount and kind of starch both play a significant role in the final texture of the food product [10]. The swelling power, solubility, and digestibility of the starch are particularly affected by the size and form of the granules [11,12]. A larger size of

starch granules results in a decreased enzymatic susceptibility [13]. Recently, it has been discovered that the size of the starch granule is a significant factor in both dietary and industrial applications of starches [14]. Industrialists are interested in the introduction of novel starch sources since it has the potential to affect the global market. Starches from roots and tubers have received significant attention recently due to their prospective applications [15,16,17].

Arrowroot (*Maranta arundinacea*, Family: Marantaceae) is an underutilized tuber crop in Sri Lanka which is locally known as “Hulankeeriya” Aerukka” and is used in folk medicine. Arrowroot tuber starch has emerged as a viable dietary remedy for individuals on special diets who have celiac disorders [18]. Moreover, for the purpose of production of films with good functional properties, arrowroot starch has unique physicochemical characteristics, including a high amylose content (16–27 %) [19,20]. Arrowroot starch has the advantage of excellent digestibility [21] and gelling ability [22,23]. Due to its high digestibility, arrowroot is used to feed children, the elderly, and individuals with digestive issues [3].

In order to determine whether arrowroot flour from any growing region may be used in the future for food applications, the aims of this study were to assess the granular morphology of arrowroot flour and investigate variations among flour samples from five different provinces in Sri Lanka with a view to enhancing the utilization of arrowroot plant in Sri Lanka.

2. MATERIAL AND METHODS

2.1 Sample Collection

In total, sixteen locations were selected in Sri Lanka, representing five provinces: Gampaha, Divulapitiya, Horana, Mahara (Western province), Kegalle, Deraniyagala, Rambukkana (Sabaragamuwa province), Thihagoda, Matara, Galle (Southern province), Hakgala, Welimada, Badulla (Uva province), Makandura, Kuliyaipitiya, Kurunegala (North-Western province). Arrowroot tubers were harvested at correct maturity and the flour was prepared by wet extraction method [20]. Prior to further examination, samples were sifted through a 425 μm sieve, packed in airtight containers, and placed in the freezer (-18 °C).

2.2. Determination of Starch Granular Morphology

A 1:1 (v/v) ratio of distilled water to glycerin was used to create the starch suspension. After staining a sample of starch suspension with a 1.0% iodine solution, a thin smear was made on a glass slide and covered with a cover slip. A light microscope (OPTICA Microscope B-290, Italy) was used to measure the size of the arrowroot starch granules and the OPTICA Pro View digital camera program N-plan was used as the imaging software [5]. Dry arrowroot starch powder was deposited in a thin layer on adhesive metallic support and then sputter coated with gold. The metalized samples were examined using a scanning electron microscope (SU 6600, HI-2108-003, Japan) operated at 5 kV [24].

2.3. Experimental Design and Statistical Analysis

Measurements were done for arrowroot flour samples from five different provinces as triplicates. Data analysis was done using analysis of variance (ANOVA) at 0.05 confidence level from MINITAB statistical software (version 19).

3. RESULTS AND DISCUSSION

The variations in length (μm) and width (μm) of arrowroot starch granules from five different provinces are presented in Table 1. Spherical shape average-sized starch granules were selected to take the measurements. According to the measurements taken on arrowroot starch samples, the length and width of the granules from five different provinces of Sri Lanka were not statistically significant ($P < 0.05$). The mean length of starch granules ranged from 42.91 - 45.86 μm while the width ranged between 30.81 – 32.32 μm . Previous studies conducted in Brazil have recorded that the average granule size in arrowroot starch was 24.97 μm , while the range of sizes was 20 μm - 40 μm [3,25] while in another study the average starch granule size was given as 56.60 μm [26]. The size of oval shape arrowroot starch granules varied from 8.6 - 42 μm as reported by [27,28]. A study conducted in Sri Lanka has indicated the average granule diameter as 23.7 μm [29]. Arrowroot starch granule dimensions of the current study are compatible with the results of past research studies. However, starch granule sizes may vary due to the difference in their botanical origins [30]. Nevertheless, there was no significant variation in arrowroot starch granule sizes from different geographical locations within Sri Lanka.

Table 1. Variation in arrowroot starch granule size from five different provinces in Sri Lanka

Province	Length (μm)	Width(μm)
North-Western province	43.25 \pm 4.68 ^a	30.81 \pm 3.15 ^a
Sabaragamuwa province	45.86 \pm 5.43 ^a	32.32 \pm 2.54 ^a
Southern province	42.91 \pm 4.50 ^a	31.97 \pm 3.01 ^a
Uva province	45.10 \pm 4.30 ^a	31.44 \pm 3.76 ^a
Western province	44.99 \pm 5.06 ^a	31.34 \pm 4.16 ^a

Values are Mean \pm SD

The same superscript letter in each column represents values not significantly different from each other at $p = 0.05$.

Based on the granular size, the following classes for the size of starch granules have been identified: Very small (<5 μm), Small (5-10 μm), medium (10-25 μm), and large (>25 μm) [12]. According to this categorization, arrowroot starch granules can be categorized as 'large' because their size falls between 30 - 46 μm . Most cereal starch granules are smaller than those from tuber and legume starches [5]. Size distribution varies depending on the stage of maturity of the plant and the form of tuber [25,31]. In the current study, the arrowroot tubers used were at the harvesting stage for the market.

Granule shape is a highly important characteristic to identify the uniqueness of the starch. According to published reports, the granules of arrowroot starch were diverse in shape and size, varying from elongated, spheroid, and oval in shape [3, 26, 27, 28, 32]. Granules of starch have a smooth surface without any indications of fissures [20]. Three identical starch granule shapes were identified in the current study, namely oval, spherical, and irregular globular shapes. Oval shape was the most predominant shape among all samples. Most of the starch granules were oval or spherical with a considerable roundness.




Fig. 1. (a) Light micrograph (x40) of different shapes of arrowroot starch granules, (b) Light micrograph (x100) of different shapes of arrowroot starch granules, (c) Scanning electron microscope image (x500) of arrowroot starch granules, (d) Scanning electron microscope image (x150) of starch granules (A – spherical shape, B – Oval shape, C- Irregular globular shape)

The majority of starch granules, 48.46 % - 59.34 % had oval shapes, which were more common than spherical and irregular globular shapes. The second most common shape was spherical, followed by irregular globular shape. The irregular globular shape is a middle shape between the spherical and oval shapes. Arrowroot from the Western province contained a relatively higher percentage of oval shape starch granules. Nonetheless, the results indicate that a high consistency is present in the granule size and shape of starch from different geographical regions in Sri Lanka.

The size of starch granules and starch digestibility were found to be negatively correlated in some previous studies [33,34]. Due to their higher specific surface area for enzyme interaction and response, smaller starch granules are more receptive to enzymatic hydrolysis [35]. In addition to granule size, starch digestibility is also influenced by its shape. The specific surface area of the granule is substantially impacted by the shape of the granule, which ranges from spherical to polyhedral [33].

The quality of starch is significantly influenced by the size of the starch granules. Smaller granules have worse swelling characteristics than larger ones, which causes them to gelatinize more slowly. They also have poor storing and flavor properties. The starch content of starch grains reduces as their average size grows but their starch content increases. The size and shape of the starch grains affect the temperature of gelatinization, the quantity of bound moisture, the viscosity of the starch paste, the ratio of the starch fractions, the color of the iodine sample, and other physicochemical features. [36]. Because arrowroot is having large starch granules, it gelatinizes quickly and forms a viscous gel. Therefore, arrowroot flour can be used as a thickener or stabilizer in the food industry.

4. CONCLUSION

There are three predominant starch granular shapes for arrowroot: oval, spherical, and irregular globular shape. Among them, the oval shape is the most prevalent shape (48.46 % - 59.34 %) among all the collected flour samples while the spherical shape is the second most prevalent (23.48 % - 29.82 %). The length and width of arrowroot starch granules did not significantly differ among flour samples. The length of granules ranges from 42.91 - 45.86 μm while the width of ranged from 30.81 - 32.32 μm . Arrowroot flour samples from the five provinces were not significantly different in terms of the starch granular morphology and the flour could be used in the local food industry irrespective of the growing location.

ACKNOWLEDGEMENTS

The authors wish to acknowledge National Research Council Grant 20 – 074. And also authors are grateful to laboratory staff members of the department of Biosystems Engineering, Faculty of Agriculture and Plantation Management, Wayamba University of Sri Lanka. Mrs. R.M.A.I.A.N. Rajakaruna, Mr. R.D.R.K. Nawarathne from Department of Bio-systems Engineering, Faculty of Agriculture and Plantation Management, Wayamba University of Sri Lanka are acknowledged for the provided technical support for the research.

1

2

3

COMPETING INTERESTS

4

5

This research was funded by National Research Council Grant 20-074.

6

7

AUTHORS' CONTRIBUTIONS

8

9

Author M.K.S. Malki conducted research work, data collection and analysis and drafting of the manuscript; Authors J.A.A.C. Wijesinghe, R.H.M.K. Ratnayake and G.C. Thilakarathna contributed by developing the concept, research design, scientific advice, correction of the manuscript. All authors read and approved the final manuscript.

10

11

12

13

14

15

REFERENCES

16

17

[1] Vegh KR. Starch bearing crops as food sources, Cultivated plants primarily as food sources. 2002;1. ISBN: 978-1-84826-100-6.

18

19

[2] Qiang L. Chemistry, physical properties and applications, Food carbohydrates, (Steve W .Cui), CRC press. 2005. ISBN:978-0-8-493-1574-9, DOI 10.1201/9780203485286.ch7.2005.

20

- 21 [3] Souza DC, Silva RJ, Guerra TS, Silva LFL, Resende LV, Pereira J. Characterization of
22 arrowroot starch in different agronomic managements. Rev. Ceres, Vicoso, 2019;66(5): 323-
23 332.
- 24 [4] Burrell MM. Starch: the need for improved quality and quantity – an overview. Journal of
25 Experimental Botany. 2003; 54:451-456.
- 26 [5] Wijesinghe JAAC, Wickramasinghe I, Saranandha KH, Variance of Granular Morphology
27 in Kithul (*Caryota urens*) Flour among Five Different Growing Areas in Sri Lanka,
28 International Journal of Multidisciplinary Studies (IJMS).2014;2(1):74-78.
- 29 [6] Jane J. Current understanding on starch granule structures. J. Appl. Glycosci, 2006;53:
30 205-213.
- 31 [7] Lentfer CJ, Therin M, Torrence R. Starch grains and environmenta reconstruction: a
32 modern test case from west new Britain, Papua New Guinea. Archaeol. Sci. 2002; 29:687-
33 698.
- 34 [8] Copeland L, Blazek J, Salman H, Tang MC. Form and functionality of starch, Food
35 hydrocolloids.2009;23:1527-1534.
- 36 [9] Jane J, Kasemsuwan T, Leas S, Zobel H, Robyt JF. Anthology of starch granule
37 morphology by scanning electron microscopy. Starch/Starke, 1994;46:121-129.
- 38 [10] Billaderis CG. The structure and interaction of starch with food constituents, Can J
39 physiol pharmacol. 1999;69:60-78.
- 40 [11] Svihus B, Uhlen AK, Harstad OM. Effect of starch granule structure, associated
41 components and processing on nutritive value of cereal starch: A review, Elsevier; Animal
42 Feed Science and Technology. 2005;122(306):303-320.
- 43 [12] Adejomo A, Aderibigbe AF, Layokun SK. Cassava Starch: Production, Physicochemical
44 Properties and Hydrolysatation- A Review. Advances in Food and Energy Security. 2011;2.
- 45 [13] Franco CML. and Ciacco CF. Factors that Affect the Enzymatic Degradation of Natural
46 Starch Granules. Effect of the Size of the Granules. Starch/Starke, 1992;44:422-426.
- 47 [14] Campbell MR, Li J, Berke TG, Glover DV. Variation of Starch Granule Size in Tropical
48 Maize Germ Plasm. Cereal Chem. American Association of Cereal Chemists, Inc.
49 1996;73(5):536-538.
- 50 [15] Takeiti C, Fakhouri F, Ormenese R, Steel C, Collares F. Freeze-thaw stability of gels
51 prepared from starches of non-conventional sources. Starch/Starke, 2007;59:156-160.
- 52 [16] Nwokocha LM, Williams PA. Some properties of white and yellow plantain (*Musa*
53 *paradisica*, *normalis*) starches. Carbohyd. Polym. 2009;76:133-138.
- 54 [17] Jyoti PS, and Konwar BK. Physicochemical properties of starch from aroids of north
55 east India. Int. J. Food Prop. 2011.DOI: 10.1080/ 10942912.2010.491929.
- 56 [18] Madineni NM, Sheema F, Ragu SS, Ramasamy R, Manisha G. Morphological,
57 Structural, and functional properties of Maranta (*Maranta arundinacea* L) Starch. Food
58 science and Biotechnology, 2012;21:747-752.
- 59 [19] Moorthy SN. Physicochemical and functional properties of tropical tuber starches: A
60 review. Starch/Starke, 2002;54:559-592.
- 61 [20] Nogueira GF, Fakhouri FM, Oliveira RA. Extraction and characterization of arrowroot
62 (*Maranta arundinacea*) starch and its application in edible films, Carbohydrate polymers,
63 2018;186:64-72.
- 64 [21] Villas-Baos F, and Franco CML. Effect of bacterial β -amylase and fungal α -amylase on
65 the digestability and structural characteristics of potato and arrowroot starches. Food
66 Hydrocolloids.2016; 52:795-803.
- 67 [22] Hoover R. Composition, molecular structure, properties and modification of pulse
68 starches: A review. Carbohydrate polymers, 2001;45(3):253-267.
- 69 [23] Charles AL, Cato K, Huang TC, Chang YH, Ciou JY, Chang JS, et al. Functional
70 properties of arrowroot starch in cassava and sweet potato composite starches. Food
71 Hydrocolloids, 2016; 53:187-191.

- 72 [24] Horovitz O, Cioica N, Jumate N, Pojar-Fenesan M, Balea A, Liteanu N, Mocanu A,
73 Tomodia-Cotisel M. SEM Characterization of starch granules. *Studia ubb*
74 *Chemia*,2011;1:212-219.
- 75 [25] Leonal M. Analise de forma e tamanho de granulos de amidos de diferentes fonts
76 botanicas. *Ciencia e Tecnologia de Alimentos*. 2007;27:579-588.
- 77 [26] Valencia GA, Moraes ICF, Lourenco RV, Habitante, Bittante AMQB, Sobral PJA.
78 Physicochemical properties of Maranta (*Maranta arundinacea* L.) starch, *International*
79 *journal of food properties*. 2014; 18: 1990-2001.
- 80 [27] Erdman MD. Starch from Arrowroot (*Maranta arundinacea*) grown at Tifon, Georgia,
81 *Cereal Chemistry*.1986; 63 (39): 277-279.
- 82 [28] Peroni FHG, Rocha TS, Franco CML. Some structural and physicochemical
83 characteristics of tuber and root starches, *Food. Sci. tech. Int*. 2006;12(6):505-513.
- 84 [29] Wickramasinghe HAM, Takigawa S, Matsuura-Endo C, Yamauchi H, Noda T.
85 Comparative analysis of starch properties of different root and tuber crops of Sri Lanka,
86 *Food Chemistry*. 2009;112(1):98-103.
- 87 [30] Lindeboom N, Chang PR, Tyler RT. Analytical, Biochemical an Physicochemical
88 Aspects of Starch Granule Size, with Emphasis on Small Granule Starches: A Review.
89 *Starch/Stärke*, 2004; 56:89-99.
- 90 [31] Singh N, Singh J, Kaur L, Sodhi NS, Gill BS. Morphological, thermal and rheological
91 properties of starches from different botanical sources. *Food Chemistry*.2003; 81:219-231.
- 92 [32] Perez E, Larez M, Gonzalez Z. Some characteristics of sagu (*Canna edulis*) and zulu
93 (*Maranta* sp) rhizomes. *Journal of Agricultural and Food chemistry*. 1997;45:2546-2549.
- 94 [33] Singh J, Dartois A, Kaur L. Starch digestibility in food matrix: a review. *Trends Food Sci*
95 *Technol*. 2010;21:168–180. doi: 10.1016/j.tifs.2009.12.001.
- 96 [34] Asare EK, Jaiswal S, Maley J, Båga M, Sammynaiken R, Rossnagel BG, Chibbar RN.
97 Barley grain constituents, starch composition, and structure affect starch in vitro enzymatic
98 hydrolysis. *J Agric Food Chem*. 2011;59:4743–4754. doi: 10.1021/jf200054e.
- 99 [35] Tester RF, Karkalas J, Qi X. Starch—composition, fine structure and architecture. *J*
100 *Cereal Sci*. 2004;39:151–165. doi: 10.1016/j.jcs.2003.12.001.
- 101 [36] Zabolotets A, Litvyak V, Yermakou A, Ospankulova G. Morphological characteristics of
102 starch granules of Eastern and Central European potato varieties (*Solanum Tuberosum*).
103 *Food Technology*. 2018;8(1):18-33.

104
105
106
107