

## **Original Research Article**

### **Effects of melatonin priming on seed germination of wheat under salt stress**

**ABSTRACT :** Melatonin (N-acetyl-5-methoxytryptamine) has emerged as a new growth regulator in plants due to its positive role in alleviation of a biotic stresses. The experiment was conducted at the Wheat Cultivation Department Laboratory, Gezira Scheme, Sudan. To investigate the effects of melatonin priming in alleviating the negative effect of salt stress on seed germination in two cultivars of winter wheat (*Triticum aestivum* L. cv Elnilein and Imam). Six concentrations were used: control, salt (300 mM NaCl) and four concentrations of melatonin (M) 10, 100, 500 and 1000  $\mu$ M with salt (300 mM NaCl) add to each concentration: CT, ST, M1+ST, M2+ST, M3+ST, M4+ST. The treatments were arranged in a Randomized Complete Block Design with three replications. The parameters studied included: seed germination rate (%), germination index, mean germination time (d), lengths (cm) and dry weight (mg) of coleoptile and radicle, starch, soluble sugar, and sucrose contents, amylase activities, activities of the antioxidant enzymes including superoxide dismutase (SOD), peroxidase (POD), ascorbate peroxidase (APX) and catalase (CAT), reactive oxygen species (ROS) including malondialdehyde (MDA), Hydrogen peroxide ( $H_2O_2$ ) contents and superoxide ( $O_2^-$ ) production rate, abscisic acid (ABA). The parameters of seedling establishment were: Shoot height, root length, fresh and dry weight of shoot and root. The treatment M3+ST (500  $\mu$ M Melatonin + 300 mM NaCl) had the highest seed germination rate, germination index, lengths and dry weights of coleoptile and radicle,

soluble sugar, sucrose contents, amylase and antioxidant enzymes activities, and ABA content, seedling shoot and root length, and fresh and dry weight of shoot and root in Imam and Elnilein cultivars, respectively .

The study revealed that the melatonin priming effectively alleviates salt stress during seed germination and seedling establishment in winter wheat.

**Key words :** Salt stress, Melatonin, Germination, Antioxidant system, Winter wheat (*Triticum aestivum* L.)

## 1. Introduction

Amongst the various a biotic stresses that plants convene, heat and salt stress are mainly imperative in agriculture since both stresses can lead to considerable reduce in crop yield and are predictable to raise in incidence in the near future because of a consequence of climate change (Wassmann et al., 2009). Presently about 6% of total land area or 20% agricultural total lands are influenced by salt (Ibrahim et al., 2016). Seed germination and seedling establishment are the two critical stages for the crops establishment (Hubbard et al., 2012), and harmfully affected by salt stress (Hampson and Simpson 1990). Salinity causes reduced emergence and impaired growth presumably due to salt-induced osmotic effects, specific ion toxicity, nutrient imbalance, oxidative damage, and alterations in endogenous levels of hormones (Munns and Tester 2008; Ashraf 2009; Fahad et al., 2015). In this admiration (Jamil et al., 2005), (Yadav et al., 2011), (Rouhi et al., 2011) and (Ansari and Sharif-Zadeh 2012) affirmed that increasing salt content reduces the germination percentage and increases time of germination.

Salinity can influence the seeds germination by the toxic effect of sodium and chloride ions on viability of embryo (Jahromi et al., 2008; Daszkowska-Golec 2011). The toxic effects include disruption to the

structure of enzymes and other macromolecules, damage to cell organelles and the plasma membrane, the disruption of respiration, photosynthesis and protein synthesis (Panda and Khan 2009; Daszkowska-Golec 2011; Matraszek et al., 2015). malondialdehyde (MDA) accumulation increases through the increase in salt stress in different crops (Khan and Weber 2006). MDA content is a signal of membrane injury at the cellular levels in salt stress. Hence, the concentration of MDA can serve as an essential oxidative stress sign (Gong et al., 2008). Oxidative stress is the term utilized to define the imbalance between production of ROS and scavenging or detoxification (Gamir et al., 2014). Plants have developed the antioxidant defense system to reduce the harmful impact of ROS like antioxidant enzymes include superoxide dismutase (SOD), peroxidase (POD) catalase (CAT), glutathione reductase (GR) and ascorbate peroxidase (APX) but salt stimulate ROS production exceeds capacity of antioxidant in the cells, oxidative stress arises (Erdal and Demirtas 2010; Genisel et al., 2013).

Melatonin (*N*-acetyl-5-methoxytryptamine) has comprehensively been known as an important hormone of animal occupied in many biological processes (Escames et al., 2012; Calvo et al., 2013). However, its role in animals the same like a hormone has been well-characterized, its information in plants kingdom is still blurry (Arnao and Hernández-Ruiz 2014). Melatonin is considered as a growth regulator of plant resulting of its particular physiological role the same as growth regulation of the shoots, root and explants, germination of seed, rhizogenesis and remain senescence in leaf (Arnao and Hernández-Ruiz 2014). The melatonin mechanisms mediated tolerance of stress occupy the antioxidant support biosynthesis, the start of connected enzymes and the directly ROS scavenging subsequent the plants exposure to harm environments (Li et al., 2012; Tan et al., 2012; Park, Lee et al. 2013; Zhang et al., 2013;

Bajwa et al., 2014). Previous study has reported that while applied exogenously melatonin alleviates oxidative injury through ROS scavenging and stimulating antioxidant system that are in different environmental stresses (Dubbels et al., 1995; Lei et al., 2004; Tan et al., 2007; Paredes et al., 2009; Posmyk et al., 2009; Zhao et al., 2011; Szafrńska et al., 2012; Tan et al., 2012; Bajwa et al., 2014).

In view of the above information, the aim of this study focused on determining the effect of melatonin pretreatment on the seed germination, physiological, and biochemical change happening in winter wheat gemmating seeds exposed to salt stress. The research will contribute to explaining melatonin's multifaceted effects on stress-induced oxidative injure.

## **2. Materials and methods**

### **2.1 Experimental design**

The study was performed in the wheat cultivation department Laboratory, Gezira scheme, Gezira state, Sudan. Two cultivars of winter wheat (*Triticum aestivum* L. cv Elnilein and Imam) uniform seeds were chosen and surface-sterilized with 2.5% sodium hypochlorite for 10 min and washed seven times with sterile distilled water according to the protocol of (Zheng et al., 2009). Two treatments were placed with soaking the seeds in distilled water as control and 10, 100, 500, and 1000  $\mu$ M of melatonin solution for 20 h, respectively. The concentrations of melatonin used here were established to be optimal in the preliminary experiments. The soaked 50 seeds of each cultivar were arranged on a filter paper in petri dishes with a diameter of 9 cm, and 300 mM of NaCl was then added in to each dish. The dishes were incubated at  $22 \pm 0.5$  °C for 7 days in to Chamber. At the two-leaf stage (ca. 20 days after germination), wheat plants were transplanted to plastic containers (45 cm

in length, 35 cm in width and 18 cm in height). The density was 30 plants in each container. The Hoagland nutrient solution was used for hydroponic culture. The growth temperatures were 25 °C/20 °C (day/night), six treatments were used are: Control , 300 mM of salt (NaCl) , 10 µM of melatonin+300 mM NaCl , 100 µM of melatonin+300 mM NaCl, 500 µM of melatonin+300 mM NaCl , 1000 µM of melatonin+300 mM NaCl, arranged as: CT, ST, M1+ST, M2+ST, M3+ST, M4+ST. The experiment was a Randomized Complete Block Design with three replications. Germinating seeds of each cultivar were sampled at days 2, 3, 4, 5 and 7 after treatment for biochemical and physiological measurements, and another batch was sampled at day 7 after treatment for determinations of dry weight, seed germination rate, and ABA content. At the four-leaf stage, wheat seedlings were harvested for measurements of seedling shoot height, root length, fresh and dry weights of shoot and root.

## **2.2 Germination rate (GR), mean germination time (MGT), germination index (GI), dry weight of germinating seeds**

GR is the proportion of germinated seeds in each Petri-dish during 7 days after germination. MGT was determined following the method of (Ellis and Roberts 1978), i.e.  $MGT = \sum (f \times i) / \sum f$ . where, f is the number of recently germinated seeds on Germinating Day i. Subsequently, long MGT indicates slow germination. Germinated seeds percentage on each day was measured. The method of (Wang et al., 2004) was used to calculate the GI as  $GI = \sum (G_i / T_i)$ , where  $G_i$  is the germination percentage at the ith day, and  $T_i$  is day of germination test. High GI demonstrates high quality of seed and performance (Wang et al., 2004).

Radicles, coleoptile and the seed residues of germinating seeds of two cultivars were selected and killed at 105 °C for 2 h, after that dried at 80 °C awaiting stable weight.

### **2.3 Contents of starch and soluble sugars**

Germinating seed Oven- dried powder (1 g) was mixed by 10 ml 0.33 mM of HCl and then heated at 100 °C for 10 min to remove starch. Half ml of ZnSO<sub>4</sub> (30 %, w/v) was added to the extraction to remove proteins. After that, 0.5 ml K<sub>3</sub> [Fe (CN)<sub>6</sub>] (30 %, w/v) was added and mixed, and then the mixture volume was attuned to 20 ml with distilled water. After completely shaken and filtrated, the concentration of starch was determined like the amount of glucose produced at 20–25 °C using the polarimetric analysis through an automatic recording polarimeter (WZZ-2B, Shanghai, China) according to (Xie et al., 2003). Moreover, to extract total soluble sugars, 0.1 g Oven- dried germinating seed powder sample was homogenized in (80 %) ethanol at 80 °C for 10 min and centrifugation at 600g. The supernatant was collected for content of soluble sugar and sucrose measurement using the method of anthrone (Fales 1951).

### **2.4 Amylase extraction and activity assess**

Activity of amylase and extraction were estimated fallowing the method of (Kishorekumar et al., 2007). The concentration of soluble protein in germinating seeds was estimated fallowing the method of (Bradford 1976) by using bovine serum albumin (BSA) like standard. Activities of both  $\alpha$ - and  $\beta$ -amylase were articulated as unit mg<sup>-1</sup> protein, and one unit is equivalent to release of one mg maltose from starch per minute by the amylases.

## **2.5 Activities of antioxidant enzymes and O<sub>2</sub>- production rate, H<sub>2</sub>O<sub>2</sub>, MDA concentrations**

The MDA and H<sub>2</sub>O<sub>2</sub> content measurements and antioxidant enzymes activities in germinating seed were measured by the method of (Tan et al., 2008). In brief, 0.5 g samples of germinating seed were sliced and homogenized in a mortar and pestle with 5 ml of ice-cold extraction buffer (50 mM potassium phosphate buffer, pH 7.0, 0.4 % polyvinylpyrrolidone (PVP)). The homogenate was centrifuged at 10,000xg for 30 min and the supernatant was used like crude extract. All procedures were done at 4 °C. MDA content was calculated followed the method of (Du and Bramlage 1992). In brief, 20 % trichloroacetic acid containing 0.5 % 2-thiobarbituric acid was added to 0.2 ml of the suspension and heated at 95°C for 25 min after that cooled down in a cool water bath and centrifuged at 3,000xg for 5 min. Concentration of MDA was calculated at 532 nm and corrected by subtracting the absorbance at 600 nm. The concentrations of H<sub>2</sub>O<sub>2</sub> was carried out by the method of (Moloi and van der Westhuizen 2006). Briefly, 1 ml suspension was added to 0.1 ml 20 % titanium tetrachloride (TiCl<sub>4</sub>) and mixed with shaking. 0.2 ml of 25 % ammonia was added to the mixture. The reaction solution was after that centrifuged at 5,930xg for 5 min, and the precipitate was washed three times with cold acetone to eradicate the color. Lastly, the precipitate was dissolved in 3 ml 1 M H<sub>2</sub>SO<sub>4</sub>, and filtered prior to measurement of the absorbance at 415 nm. O<sub>2</sub><sup>-</sup> production rate was measured using the method of (Zheng et al., 2009). The activities of SOD, and CAT were calculated according to (Tan et al., 2008). For SOD activity measurement, 3 ml reaction mixture contained 130 mM methionine, 750 mM NBT, 100 mM EDTA and 50 ml of enzyme extract in 50 mM phosphate buffer (pH 7.8). The reaction was started with 20 mM riboflavin through exposing the cuvette to a 15 W circular white light tube for 10 min. One unit of

SOD activity was identified as the amount of enzyme per mg of protein sample causing 50 % inhibition of the rate of NBT reduction at 560 nm (Yang et al., 2007). CAT activity was estimated as described by (Tan et al., 2008). The activity of POD was determined according to the changes in absorbance at 470 nm due to guaiacol oxidation. Ascorbate peroxidase (APX) was measured using the method of (Nakano and Asada 1981).

## **2.6 Determination of endogenous hormone concentration**

The content of abscisic acid (ABA) was analyzed by an indirect ELISA technique. Briefly, the samples of 0.1 g dry seeds was homogenized in liquid nitrogen and removed in cold 80% (v/v) methanol by butylated hydroxytoluene (1 mM) overnight at 4 °C. The extracts were then collected after centrifugation at 10,000 x g (4 °C) for 20 min, and after that the extracts were passed throughout a C<sub>18</sub> Sep-Pak cartridge (Waters, Milford, MA) and dried in N<sub>2</sub>. Then the residues were melted in PBS (0.01 M, pH 7.4). To estimate the content of ABA using the method described by (You-Ming et al., 2001). Three replicates were used for each treatment.

## **2.7 Statistical analysis**

All data were subjected to the one-way analysis of variance (ANOVA) to determine the significant differences between the treatments using the software of SPSS (Ver. 10.0 SPSS, Chicago, IL, USA).

# **3. Results**

## **3.1 Seed germination parameters**

Seed germination rate was significantly reduced under salt stress. During 7 days of germination, seed germination rate (GR) and germination index (GI) were considerably enhanced whereas mean germination time (MGT) was reduced by melatonin priming treatment



when compared with salt stressed-seed in both cultivars, with values being higher in Imam than that in the Elnilein (Table 1). Also, lengths and dry weights of coleoptile and radicle were clearly increased in both cultivars while seed residues dry weight was significantly ( $P < 0.05$ ) reduced with application of melatonin in both cultivars, as compared with salt stress alone. Therefore, this increase was more prominent in Imam than that in the Elnilein (Table 1). However, seed pretreatment with melatonin application it seemed that enhanced seed germination in wheat under salt stress.

**Table 1** Effect of melatonin priming on seed germination rate (GR), germination index (GI), mean germination time (MGT), length of radicle and coleoptile, and dry weight of radicle, coleoptile and whole seed after 7 days germination under salt stress in wheat.

Cultivar	Treatments	GR(%)	GI	MGT(d)	Length (cm)		Dry weight (mg seed <sup>-1</sup> )		
					Radicle	Coleoptile	Radicle	Coleoptile	Seed residue
Elnilein	CT	95.36±0.28a	92.78±0.05a	1.54±0.03d	9.38±0.05a	8.27±0.03a	13.18±0.02a	10.26±0.05a	7.13±0.02f
	ST	64.39±0.26e	21.49±0.36f	4.62±0.22a	0.9±0.01f	0.82±0.01f	1.39±0.05f	1.12±0.01f	32.36±0.04a
	M1+ST	76.28±0.23d	26.13±0.27e	3.59±0.13b	1.59±0.02e	1.21±0.02e	2.84±0.02e	2.03±0.01e	25.71±0.15b
	M2+ST	88.47±0.41b	51.47±0.23c	1.87±0.13d	5.23±0.02c	4.78±0.14c	6.35±0.04c	5.01±0.04c	12.45±0.16d
	M3+ST	94.23±0.20a	74.19±0.18b	1.73±0.01d	8.04±0.01b	7.14±0.04b	12.03±0.02b	8.63±0.28b	8.49±0.09e
	M4+ST	81.63±0.46c	46.68±0.33d	2.42±0.04c	3.12±0.01d	3.29±0.03d	4.15±0.01d	3.23±0.01d	20.18±0.04c
Imam	CT	97.83±0.40a	95.32±0.17a	1.47±0.02c	10.89±0.09a	9.43±0.23a	15.72±0.06a	11.79±0.09a	9.27±0.05f
	ST	67.82±0.37f	25.46±0.30f	3.48±0.08a	0.95±0.01f	0.91±0.01f	1.98±0.01f	1.35±0.01f	36.55±0.27a
	M1+ST	78.53±0.31e	29.38±0.49e	2.92±0.06b	1.85±0.02e	1.98±0.05e	3.94±0.02e	3.28±0.03e	29.18±0.36b
	M2+ST	91.72±0.35c	69.37±0.90c	1.69±0.02c	7.33±0.03c	5.83±0.05c	9.47±0.18c	7.37±0.19c	13.95±0.03d
	M3+ST	95.57±0.27b	78.67±0.65b	1.58±0.02c	8.63±0.22b	8.65±0.27b	13.69±0.14b	9.72±0.29b	10.87±0.31e
	M4+ST	82.68±0.67d	55.03±0.63d	2.12±0.07b	5.63±0.08d	4.61±0.04d	7.44±0.04d	5.04±0.01d	23.59±0.04c

Different small letters in the same column refer to significant difference between treatments at  $P < 0.05$  level


### 3.2 Starch, soluble sugar, and sucrose contents

During seed germination progress starch content decreased, but total soluble sugar and sucrose contents were increased in germinating seeds of wheat in both cultivars (Fig. 1). Moreover, in both cultivars, starch


content of seed was lower under melatonin priming compared to salt stress during 7 days of germination (Fig. 1). In compare, the contents of total soluble sugar and sucrose were higher in the control and melatonin than that under salt stress in both cultivars. However, total soluble sugar and sucrose contents were significantly ( $P < 0.05$ ) higher in Elnilein than those in Imam and oppositely in case of starch content (Fig. 1). These results indicated that melatonin priming improved the resistance of wheat seed to salt stress.

### **3.3 Activities of $\alpha$ - and $\beta$ - amylase**

Under control the  $\alpha$ -amylase activity in both wheat cultivars were improved in seeds germinating than that in the level of salt stress during 7 days of germination (Fig. 2). Nevertheless, melatonin priming significantly ( $P < 0.05$ ) enhanced  $\alpha$ -amylase activity, when contrasted with salt stress alone. Furthermore, in both cultivars  $\beta$ -amylase activity decreased under salt stress. On other hand, melatonin pretreatment significantly ( $P < 0.05$ ) increased the activity of  $\beta$ -amylase compared with the salt stress alone (Fig. 2). Conversely, under control and melatonin mutually activities of  $\alpha$ - and  $\beta$ - in germinated seeds in both cultivars were higher than those under salt stress level, and they increased more in Elnilein than that in Imam.




**Fig. 1** Effect of melatonin priming on changes in contents of starch, soluble sugar, and sucrose in wheat seeds during 7 days germination under salt stress. Data are means  $\pm$ SD (n=3). Different small letters at each day indicate significant difference between treatments at  $P < 0.05$ .



**Fig. 2** Effect of melatonin priming on changes in activities of  $\alpha$ - and  $\beta$ - amylase in wheat seeds during 7 day's germination under salt stress. Data are means  $\pm$ SD (n=3). Different small letters at each day indicate significant difference between treatments at  $P < 0.05$ .


### **3.4 O<sub>2</sub><sup>-</sup> production rate, concentrations of H<sub>2</sub>O<sub>2</sub> and MDA, and activities of antioxidant enzymes**

Compared to the control, salt stress significantly ( $P < 0.05$ ) increased the concentration of MDA and H<sub>2</sub>O<sub>2</sub> in wheat germinating seed in both cultivars (Fig.3). In contrast, exogenous of melatonin treatment significantly ( $P < 0.05$ ) decreased the concentration of MDA and H<sub>2</sub>O<sub>2</sub> in both cultivars compared to salt stress alone. In both cultivars the production rate of O<sub>2</sub><sup>-</sup> demonstrates the same pattern similar to H<sub>2</sub>O<sub>2</sub> and MDA accumulation during the germinating seeds (Fig. 3). Reversely, salt stress alone significantly ( $P < 0.05$ ) reduced in the activities of SOD, POD, CAT and APX compared to control in both cultivars (Fig.4). In contrast, pretreatment of melatonin remarkably enhanced ( $P < 0.05$ ) SOD, CAT, POD, and APX activities in both cultivars and they increased more in Elnilein than that in Imam. These results confirmed that priming of melatonin was responsible for the improvement of adaptive responses of wheat seed germination against salt stress.



**Fig.3** Effect of melatonin priming on changes in contents of malondialdehyde (MDA), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), and O<sub>2</sub><sup>-</sup> production rate in wheat seeds during 7 days germination under salt stress. Data are means  $\pm$ SD (n=3). Different small letters at each day indicate significant difference between treatments at P < 0.05





**Fig. 4** Effect of melatonin priming on changes in activities of superoxide dismutase (SOD), catalase (CAT), peroxidase (POD), and ascorbate peroxidase (APX) in wheat seeds during 7 days germination under salt stress. Data are means  $\pm$ SD (n=3). Different small letters at each day indicate significant difference between treatments at  $P < 0.05$


### **3.5 ABA concentration**

After set of treatments the concentrations of ABA in germinating seeds of wheat was much higher under salt stress when compared with control in both cultivars (Fig. 5). On the other hand, under salinity stress melatonin priming significantly ( $P < 0.05$ ) enhanced germination seeds ABA content in both cultivars (Fig. 5). In general, melatonin priming had much higher levels of ABA in wheat germinating seed as compared to both control and salt stress alone. The protective effect of melatonin on ABA concentration was greater in Imam than that in Elnilein ( $P < 0.05$ ).

### **3.6 Seedling growth parameters**

After germination, wheat plants were still growing at the same temperature conditions as during their germination period till the 4-leaf stage, to further investigate effects of melatonin priming on seedling establishment under salt stress (Table 2). Salt stress induced by 300 mM NaCl significantly inhibited the growth of wheat seedlings, This attribute was demonstrated by comparing the decrease in the shoot height, root length, fresh and dry weight of shoot and root, in both cultivars. However, melatonins priming significantly ( $P < 0.05$ ) alleviate this response. The positive effect of melatonin priming had more profound in Imam than that in the Elnilein.





**Fig. 5** Effect of melatonin priming on changes in content of ABA in wheat seeds during 7 days germination under salt stress. Data are means  $\pm$ SD (n=3). Different small letters indicate significant difference between treatments at  $P < 0.05$

**Table 2** Effect of melatonin priming on seedling shoot height and root length, fresh and dry weight of shoot and root of four- leaf-stage seedling grown under salt stress in wheat.

Variety	Treatments	Shoot height (cm)	Root length (cm)	Fresh weight (mg plant <sup>-1</sup> )		Dry weight (mg plant <sup>-1</sup> )	
				Shoot	Root	Shoot	Root
Elnilein	CT	23.56 $\pm$ 1.70a	26.82 $\pm$ 1.84a	23.97 $\pm$ 0.54a	16.85 $\pm$ 0.33a	6.97 $\pm$ 0.44a	3.96 $\pm$ 0.37a
	ST	11.98 $\pm$ 0.66f	13.27 $\pm$ 0.68f	9.81 $\pm$ 0.41f	6.19 $\pm$ 0.41f	2.38 $\pm$ 0.42f	1.03 $\pm$ 0.42d
	M1+ST	15.11 $\pm$ 1.45e	17.93 $\pm$ 1.54e	13.21 $\pm$ 0.39e	8.88 $\pm$ 0.32e	3.92 $\pm$ 0.40e	1.97 $\pm$ 0.42d
	M2+ST	19.67 $\pm$ 0.50c	20.19 $\pm$ 0.28c	17.66 $\pm$ 0.36c	13.44 $\pm$ 0.63c	4.98 $\pm$ 0.57c	2.89 $\pm$ 0.53b
	M3+ST	22.44 $\pm$ 0.36b	23.74 $\pm$ 0.84b	20.13 $\pm$ 0.31b	15.67 $\pm$ 0.23b	5.11 $\pm$ 0.31b	3.58 $\pm$ 0.13b
	M4+ST	17.22 $\pm$ 0.36d	18.87 $\pm$ 0.84d	15.23 $\pm$ 0.31d	10.16 $\pm$ 0.45d	4.09 $\pm$ 0.38d	2.01 $\pm$ 0.35c
Imam	CT	25.61 $\pm$ 1.70a	28.87 $\pm$ 1.81a	26.96 $\pm$ 0.49a	19.79 $\pm$ 0.28a	8.96 $\pm$ 0.49a	5.89 $\pm$ 0.48a
	ST	13.18 $\pm$ 1.66f	15.59 $\pm$ 0.81f	12.14 $\pm$ 0.19f	7.92 $\pm$ 0.45f	3.56 $\pm$ 0.49f	1.98 $\pm$ 0.46e
	M1+ST	16.96 $\pm$ 0.78e	18.24 $\pm$ 1.52e	16.15 $\pm$ 0.56e	10.53 $\pm$ 0.16e	4.32 $\pm$ 0.57e	2.67 $\pm$ 0.46e
	M2+ST	20.73 $\pm$ 0.50c	22.43 $\pm$ 0.42c	22.22 $\pm$ 0.19c	15.91 $\pm$ 0.62c	6.12 $\pm$ 0.43c	4.78 $\pm$ 0.63c
	M3+ST	23.47 $\pm$ 0.36b	25.91 $\pm$ 0.82b	24.42 $\pm$ 0.12b	17.19 $\pm$ 0.49b	7.43 $\pm$ 0.18b	5.17 $\pm$ 0.19b
	M4+ST	18.23 $\pm$ 0.36d	20.34 $\pm$ 0.82d	19.21 $\pm$ 0.27d	13.39 $\pm$ 0.39d	5.83 $\pm$ 0.47d	3.32 $\pm$ 0.39d

Different small letters in the same column refer to significant difference between treatments at  $P < 0.05$  level

#### **4. Discussion.**

It is extremely feasible that growth inhibition in saline surroundings was caused with a reduce in the supply of available energy of biochemical essential for synthesis reactions, most important to an inhibition growth of root and coleoptiles (Geniselet al., 2015). An increasing body of confirmation has recommended that melatonin acting a significant function in responses of plant to environmental stimulus, including biotic and a biotic stresses ( Zhang et al., 2013; Bajwa et al., 2014; Lee et al., 2014). In this study, salt stress clearly reduced seed germination, growth of root and coleoptile in contrast by seedlings of control in both wheat cultivars (Table. 1). But the melatonin priming improved these parameters when compared with salt stress. These results be in agreement with the results of more studies, in which mentioned that salt stress decreased seed germination, growth of early seedling in germinating seeds of barley, bean, and maize (Huang and Redmann 1995; Erdal 2012; Erdal et al., 2012). Melatonin pretreatment was establish effectual to defend plant cell from oxidative harm (Zhang et al., 2015). In agreement with this, our results revealed that melatonin pretreatment significantly enhancing germination rate and morphogenesis of radicle and coleoptile under salt stress. This was directly related with the rapid seed starch humiliation during with speedy increase in contents of soluble sugars and sucrose in germinating seeds under salt stress (Fig.1).

Throughout seed germination, different hydrolytic enzymes are created and implicated in reserve mobilization (Zhang et al., 2005). In harmony with this, our results discovered that salt stress caused extremely inhibition on amylase activity while melatonin use considerably alleviated this inhibition improved the activities of  $\alpha$ -amylase and  $\beta$ -amylase in both cultivars (Fig. 2). Hence, melatonin priming assists the

starch change into sugars (Fig. 1), which enhancement wheat seed germination under salt stress.

Salt stress stimulates more ROS production, which create injure to macromolecules like, proteins, lipids and nucleic acids (Cakmak et al. 2010; Erdal and Dumlupinar 2010). (Mittova et al., 2003; Petrusa et al., 2008) mentioned that salt stress increased the concentration of  $H_2O_2$  and MDA in the tomato mitochondria. This might be caused by a salt-induced raise in the  $O_2$  production rate (Møller 2001). In agreement with the previous researches mentioned by (Erdal 2012). Our study appeared that salt stress quickly increased the concentration of MDA,  $H_2O_2$ , and  $O_2$ , in wheat germinating seed. But, there was a significant reduces in these factors upon the exogenous of melatonin as contrasted by salt-stressed wheat germinating seeds in both cultivars (Fig. 3). A similarly stress role of anti-oxidative of melatonin priming was also seeing in salt-stressed barley plants (Li et al., 2008). Consequently, melatonin pretreatment might be an effectual applies to defend plants and germinating seeds from oxidative harm caused by salt stress. It is strongly documented that the antioxidant enzymes like SOD, CAT, POD, and APX cooperate an important function in ROS scavenging in salt stressed plants (Tseng et al., 2007; Ashraf and Ali 2008; Tuna et al., 2008; Ashraf 2009). Previous researches have showed that exogenous melatonin defends contents against oxidative stress in reed under heat stress (Song et al., 2006) and in wheat under drought. This associated with increased the activities of SOD, CAT, POD and APX. In agreement with these results, we experiential that activities of these enzymes in the mitochondria of wheat seed were significantly improved by melatonin pretreatment application in the current study (Fig. 4), and which may have provided to the alleviated oxidative stress in the mitochondria of wheat germinating seeds and thus enhanced germination rate under salt stress.

Abscisic acid (ABA) is essential in the role of  $\alpha$ -amylase through germination (Bethke et al., 1997). (Larkindale and Knight 2002) mentioned that ABA plays a role in alleviating oxidative damage under a biotic stress. Absciscic acid (ABA) has been proposed to act as a mediator in plant responses to a range of stresses, including drought and salt stress (Ashraf and Foolad 2005). Among different phytohormones, abscisic acid (ABA), is the central controller of a biotic stress resistance in plants and organizes group of functions (Finkelstein 2013; Wani and Kumar 2015), allowing plants to cope with various stresses. ABA content considerably increases in plants under stress conditions like drought, extreme temperature, and high salinity, stimulating stress-tolerance effects that help plants adapt and survive under these stressful conditions (Ng et al., 2014), and is necessary in the plants for stress tolerance (Hamayun et al., 2010). (UMEZAWA et al., 2001; Xu et al. 2011) establish that the level of ABA in soybean seed germination increased significantly under salt stress. It is thus feasible that ABA improves salt tolerance in soybean. Our results also demonstrated that the salt stress increased the ABA content of both cultivars when compared to the control germinating seed. Furthermore, melatonin pretreatment significantly increased seed ABA contents as compared to the salt stress and control during germination under salt stress (Fig.5). Therefore, we can deduce that increasing ABA concentration due to seeds priming with melatonin may be one of the mechanisms that enabled wheat seeds to tolerate salt stress.

(Parida and Das 2005) mentioned that during the onset and development of salt stress all seedlings growth parameter are affected and reduced when compared with control plants. In this study, we found that values for almost all of the parameters of seedling growth measured were significantly ( $P < 0.05$ ) decreased in plants under salt stress as contrasted with the control. Nevertheless, melatonin priming application

considerably alleviated those inhibitory causes (Table 2). Various studies have established that melatonin can perform as a potential modulator of plant growth and improvement in a dose-dependent way (Hernandez-Ruiz, Cano et al. 2004; Afreen, Zobayed et al. 2006; Arnao and Hernández-Ruiz 2007; Hernández-Ruiz and Arnao 2008). This could be well exemplified by the increased heights and weights of both shoot and root of seedlings grown under salt stress. This of course was firstly ascribed to the significance of improved seed germination by melatonin pre-treatment.

### **Conclusions**

The study clearly demonstrated that the deleterious effects of salt stress on seed germination and early seedling growth of wheat cultivars were considerably alleviated by melatonin priming. On one hand, the exogenous application of melatonin improved soluble sugar, sucrose contents,  $\alpha$ - and  $\beta$  activities under salt stress. Also enhanced ABA content in germinating seed. Conversely, melatonin could effectively scavenge ROS and improved antioxidant enzymes activities. These findings provide a novel use of melatonin in wheat during seed germination and seedling establishment under salt stress. We expect that the positive effect of melatonin in mitigation salt stress present new opportunities for its employ in agriculture. We also finding that the higher value for most parameters measured were obtained at 500  $\mu$ M of melatonin level and it is used for our next experiment of wheat seedling performance under salt stress.

## References

- Afreen, F., S. Zobayed, et al. (2006). "Melatonin in Glycyrrhiza uralensis: response of plant roots to spectral quality of light and UV-B radiation." Journal of pineal research **41**(2): 108-115.
- Ansari, O. and F. Sharif-Zadeh (2012). "Osmo and hydro priming improvement germination characteristics and enzyme activity of Mountain Rye (*Secale montanum*) seeds under drought stress." Journal of Stress Physiology & Biochemistry **8**(4).
- Arnao, M. B. and J. Hernández-Ruiz (2007). "Melatonin promotes adventitious and lateral root regeneration in etiolated hypocotyls of *Lupinus albus* L." Journal of pineal research **42**(2): 147-152.
- Arnao, M. B. and J. Hernández-Ruiz (2014). "Melatonin: plant growth regulator and/or biostimulator during stress?" Trends in plant science **19**(12): 789-797.
- Ashraf, M. (2009). "Biotechnological approach of improving plant salt tolerance using antioxidants as markers." Biotechnology advances **27**(1): 84-93.
- Ashraf, M. and Q. Ali (2008). "Relative membrane permeability and activities of some antioxidant enzymes as the key determinants of salt tolerance in canola (*Brassica napus* L.)." Environmental and Experimental Botany **63**(1): 266-273.
- Ashraf, M. and M. Foolad (2005). "Pre-sowing seed treatment—A shotgun approach to improve germination, plant growth, and crop yield under saline and non-saline conditions." Advances in Agronomy **88**: 223-271.
- Bajwa, V. S., M. R. Shukla, et al. (2014). "Role of melatonin in alleviating cold stress in *Arabidopsis thaliana*." Journal of pineal research **56**(3): 238-245.

- Bethke, P. C., R. Schuurink, et al. (1997). "Hormonal signalling in cereal aleurone." Journal of Experimental Botany **48**(7): 1337-1356.
- Bradford, M. M. (1976). "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding." Analytical biochemistry **72**(1-2): 248-254.
- Cakmak, T., R. Dumlupinar, et al. (2010). "Chilling resistance of *Phaseolus vulgaris* and *Brassica oleracea* under a high-intensity electric field." Zeitschrift für Naturforschung C **65**(5-6): 380-386.
- Calvo, J. R., C. Gonzalez-Yanes, et al. (2013). "The role of melatonin in the cells of the innate immunity: a review." Journal of pineal research **55**(2): 103-120.
- Daszkowska-Golec, A. (2011). "Arabidopsis seed germination under abiotic stress as a concert of action of phytohormones." Omics: a journal of integrative biology **15**(11): 763-774.
- Du, Z. and W. J. Bramlage (1992). "Modified thiobarbituric acid assay for measuring lipid oxidation in sugar-rich plant tissue extracts." Journal of Agricultural and Food Chemistry **40**(9): 1566-1570.
- Dubbels, R., R. Reiter, et al. (1995). "Melatonin in edible plants identified by radioimmunoassay and by high performance liquid chromatography-mass spectrometry." Journal of pineal research **18**(1): 28-31.
- Ellis, R. and E. Roberts (1978). "Towards a rational basis for testing seed quality." Proceedings-Easter School in Agricultural Science, University of Nottingham.
- Erdal, S. (2012). "Alleviation of salt stress in wheat seedlings by mammalian sex hormones." Journal of the Science of Food and Agriculture **92**(7): 1411-1416.

- Erdal, S. (2012). "Androsterone-induced molecular and physiological changes in maize seedlings in response to chilling stress." Plant Physiology and Biochemistry **57**: 1-7.
- Erdal, S. (2012). "Exogenous mammalian sex hormones mitigate inhibition in growth by enhancing antioxidant activity and synthesis reactions in germinating maize seeds under salt stress." Journal of the Science of Food and Agriculture **92**(4): 839-843.
- Erdal, S. and A. Demirtas (2010). "Effects of cement flue dust from a cement factory on stress parameters and diversity of aquatic plants." Toxicology and industrial health **26**(6): 339-343.
- Erdal, S. and R. Dumlupinar (2010). "Progesterone and  $\beta$ -estradiol stimulate seed germination in chickpea by causing important changes in biochemical parameters." Zeitschrift für Naturforschung C **65**(3-4): 239-244.
- Erdal, S., M. Genisel, et al. (2012). "Effects of progesterone application on antioxidant enzyme activities and  $K^+/Na^+$  ratio in bean seeds exposed to salt stress." Toxicology and industrial health: 0748233711430975.
- Escames, G., G. Ozturk, et al. (2012). "Exercise and melatonin in humans: reciprocal benefits." Journal of pineal research **52**(1): 1-11.
- Fahad, S., S. Hussain, et al. (2015). "Phytohormones and plant responses to salinity stress: a review." Plant Growth Regulation **75**(2): 391-404.
- Fales, F. W. (1951). "The assimilation and degradation of carbohydrates by yeast cells." Journal of Biological Chemistry **193**(1): 113-124.
- Finkelstein, R. (2013). "Absciscic acid synthesis and response." The Arabidopsis Book **11**: e0166.
- Gamir, J., M. Cerezo, et al. (2014). "The plasticity of priming phenomenon activates not only common metabolomic fingerprint



but also specific responses against *P. cucumerina*." Plant signaling & behavior **9**(5): 227-240.

Genisel, M., S. Erdal, et al. (2015). "The mitigating effect of cysteine on growth inhibition in salt-stressed barley seeds is related to its own reducing capacity rather than its effects on antioxidant system." Plant Growth Regulation **75**(1): 187-197.

Genisel, M., H. Turk, et al. (2013). "Exogenous progesterone application protects chickpea seedlings against chilling-induced oxidative stress." Acta physiologiae plantarum **35**(1): 241-251.

Gong, H., K. Chen, et al. (2008). "Effects of silicon on defense of wheat against oxidative stress under drought at different developmental stages." Biologia plantarum **52**(3): 592-596.

Hamayun, M., S. A. Khan, et al. (2010). "Effect of salt stress on growth attributes and endogenous growth hormones of soybean cultivar Hwangkeumkong." Pak. J. Bot **42**(5): 3103-3112.

Hampson, C. R. and G. Simpson (1990). "Effects of temperature, salt, and osmotic potential on early growth of wheat (*Triticum aestivum* L.). I. Germination." Canadian Journal of Botany **68**(3): 524-528.

Hernández-Ruiz, J. and M. B. Arnao (2008). "Melatonin stimulates the expansion of etiolated lupin cotyledons." Plant growth regulation **55**(1): 29-34.

Hernandez-Ruiz, J., A. Cano, et al. (2004). "Melatonin: a growth-stimulating compound present in lupin tissues." Planta **220**(1): 140-144.

Huang, J. and R. Redmann (1995). "Salt tolerance of *Hordeum* and *Brassica* species during germination and early seedling growth." Canadian Journal of Plant Science **75**(4): 815-819.

- Hubbard, M., J. Germida, et al. (2012). "Fungal endophytes improve wheat seed germination under heat and drought stress." Botany **90**(2): 137-149.
- Ibrahim, H., M. Eldeen, et al. (2016). "Comparison of Germination and Seedling Characteristics of Wheat Varieties from China and Sudan under Salt Stress." Agronomy Journal **108**(1): 85-92.
- Jahromi, F., R. Aroca, et al. (2008). "Influence of salinity on the in vitro development of Glomus intraradices and on the in vivo physiological and molecular responses of mycorrhizal lettuce plants." Microbial Ecology **55**(1): 45-53.
- Jamil, M., C. C. Lee, et al. (2005). "Salinity (NaCl) tolerance of Brassica species at germination and early seedling growth." Electronic Journal of Environmental, Agricultural and Food Chemistry **4**(4): 970-976.
- Khan, M. A. and D. J. Weber (2006). Ecophysiology of high salinity tolerant plants, Springer Science & Business Media.
- Kishorekumar, A., C. A. Jaleel, et al. (2007). "Comparative effects of different triazole compounds on growth, photosynthetic pigments and carbohydrate metabolism of Solenostemon rotundifolius." Colloids and Surfaces B: Biointerfaces **60**(2): 207-212.
- Larkindale, J. and M. R. Knight (2002). "Protection against heat stress-induced oxidative damage in Arabidopsis involves calcium, abscisic acid, ethylene, and salicylic acid." Plant physiology **128**(2): 682-695.
- Lee, H. Y., Y. Byeon, et al. (2014). "Melatonin as a signal molecule triggering defense responses against pathogen attack in Arabidopsis and tobacco." Journal of pineal research **57**(3): 262-268.

- Lei, X. Y., R. Y. Zhu, et al. (2004). "Attenuation of cold-induced apoptosis by exogenous melatonin in carrot suspension cells: the possible involvement of polyamines." Journal of pineal research **36**(2): 126-131.
- Li, C., P. Wang, et al. (2012). "The mitigation effects of exogenous melatonin on salinity-induced stress in *Malus hupehensis*." Journal of pineal research **53**(3): 298-306.
- Li, Q.-Y., H.-B. Niu, et al. (2008). "Protective role of exogenous nitric oxide against oxidative-stress induced by salt stress in barley (*Hordeum vulgare*)." Colloids and Surfaces B: Biointerfaces **65**(2): 220-225.
- Matraszek, R., B. Hawrylak-Nowak, et al. (2015). "Protein hydrolysate as a component of salinized soil in the cultivation of *Ageratum houstonianum* Mill.(Asteraceae)." Acta Agrobotanica **68**(3): 247-253.
- Mittova, V., M. Tal, et al. (2003). "Up-regulation of the leaf mitochondrial and peroxisomal antioxidative systems in response to salt-induced oxidative stress in the wild salt-tolerant tomato species *Lycopersicon pennellii*." Plant, cell & environment **26**(6): 845-856.
- Møller, I. M. (2001). "Plant mitochondria and oxidative stress: electron transport, NADPH turnover, and metabolism of reactive oxygen species." Annual review of plant biology **52**(1): 561-591.
- Moloi, M. J. and A. J. van der Westhuizen (2006). "The reactive oxygen species are involved in resistance responses of wheat to the Russian wheat aphid." Journal of plant physiology **163**(11): 1118-1125.
- Munns, R. and M. Tester (2008). "Mechanisms of salinity tolerance." Annual Review of Plant Biology **59**: 651-681.

- Nakano, Y. and K. Asada (1981). "Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts." Plant and cell physiology **22**(5): 867-880.
- Ng, L. M., K. Melcher, et al. (2014). "Abscisic acid perception and signaling: structural mechanisms and applications." Acta Pharmacologica Sinica **35**(5): 567-584.
- Panda, S. and M. Khan (2009). "Growth, Oxidative Damage and Antioxidant Responses in Greengram (*Vigna radiata* L.) under Short-term Salinity Stress and its Recovery." Journal of Agronomy and Crop Science **195**(6): 442-454.
- Paredes, S. D., A. Korkmaz, et al. (2009). "Phytomelatonin: a review." Journal of Experimental Botany **60**(1): 57-69.
- Parida, A. K. and A. B. Das (2005). "Salt tolerance and salinity effects on plants: a review." Ecotoxicology and environmental safety **60**(3): 324-349.
- Petrussa, E., V. Casolo, et al. (2008). "Activity of a channel in Arum spadix mitochondria during thermogenesis." Journal of plant physiology **165**(13): 1360-1369.
- Posmyk, M. M., M. Bałabusta, et al. (2009). "Melatonin applied to cucumber (*Cucumis sativus* L.) seeds improves germination during chilling stress." Journal of pineal research **46**(2): 214-223.
- Rouhi, H. R., M. A. Aboutalebian, et al. (2011). "Effects of hydro and osmopriming on drought stress tolerance during germination in four grass species." International journal of Agriscience **1**(2): 107-114.
- Song, L., W. Ding, et al. (2006). "Nitric oxide protects against oxidative stress under heat stress in the calluses from two ecotypes of reed." Plant Science **171**(4): 449-458.

- Szafrńska, K., S. Glińska, et al. (2012). "Changes in the nature of phenolic deposits after re-warming as a result of melatonin pre-sowing treatment of *Vigna radiata* seeds." Journal of plant physiology **169**(1): 34-40.
- Tan, D.-X., R. Hardeland, et al. (2012). "Functional roles of melatonin in plants, and perspectives in nutritional and agricultural science." Journal of Experimental Botany **63**(2): 577-597.
- Tan, D.-X., L. C. Manchester, et al. (2007). "Phytoremediative capacity of plants enriched with melatonin." Plant signaling & behavior **2**(6): 514-516.
- Tan, W., J. Liu, et al. (2008). "Alterations in photosynthesis and antioxidant enzyme activity in winter wheat subjected to post-anthesis water-logging." Photosynthetica **46**(1): 21-27.
- Tseng, M. J., C.-W. Liu, et al. (2007). "Enhanced tolerance to sulfur dioxide and salt stress of transgenic Chinese cabbage plants expressing both superoxide dismutase and catalase in chloroplasts." Plant Physiology and Biochemistry **45**(10): 822-833.
- Tuna, A. L., C. Kaya, et al. (2008). "The combined effects of gibberellic acid and salinity on some antioxidant enzyme activities, plant growth parameters and nutritional status in maize plants." Environmental and Experimental Botany **62**(1): 1-9.
- UMEZAWA, T., K. SHIMIZU, et al. (2001). "Effects of non-stomatal components on photosynthesis in soybean under salt stress." Japanese Journal of Tropical Agriculture **45**(1): 57-63.
- Wang, Y., L. Yu, et al. (2004). "Vigor tests used to rank seed lot quality and predict field emergence in four forage species." Crop Science **44**(2): 535-541.
- Wani, S. and V. Kumar (2015). "Plant stress tolerance: engineering ABA: a Potent phytohormone." Transcriptomics **3**: 113.

- Wassmann, R., S. Jagadish, et al. (2009). "Climate change affecting rice production: the physiological and agronomic basis for possible adaptation strategies." Advances in Agronomy **101**: 59-122.
- Xie, Z., D. Jiang, et al. (2003). "Relationships of endogenous plant hormones to accumulation of grain protein and starch in winter wheat under different post-anthesis soil water statuses." Plant Growth Regulation **41**(2): 117-127.
- Xu, X.-y., R. Fan, et al. (2011). "Proteomic analysis of seed germination under salt stress in soybeans." Journal of Zhejiang University-Science B **12**(7): 507-517.
- Yadav, P. V., M. Kumari, et al. (2011). "Seed priming mediated germination improvement and tolerance to subsequent exposure to cold and salt stress in capsicum." Research Journal of Seed Science **4**: 125-136.
- Yang, S.-H., L.-J. Wang, et al. (2007). "Ultraviolet-B irradiation-induced freezing tolerance in relation to antioxidant system in winter wheat (*Triticum aestivum* L.) leaves." Environmental and Experimental Botany **60**(3): 300-307.
- You-Ming, Y., X. Chu-Nian, et al. (2001). "Effects of plant growth regulators on secondary wall thickening of cotton fibres." Plant Growth Regulation **35**(3): 233-237.
- Zhang, H., W.-B. Shen, et al. (2005). "A rapid response of  $\beta$ -amylase to nitric oxide but not gibberellin in wheat seeds during the early stage of germination." Planta **220**(5): 708-716.
- Zhang, N., Q. Sun, et al. (2015). "Roles of melatonin in abiotic stress resistance in plants." Journal of Experimental Botany **66**(3): 647-656.

- Zhang, N., B. Zhao, et al. (2013). "Melatonin promotes water-stress tolerance, lateral root formation, and seed germination in cucumber (*Cucumis sativus* L.)." Journal of pineal research **54**(1): 15-23.
- Zhao, Y., L. W. Qi, et al. (2011). "Melatonin improves the survival of cryopreserved callus of *Rhodiola crenulata*." Journal of pineal research **50**(1): 83-88.
- Zheng, C., D. Jiang, et al. (2009). "Effects of salt and waterlogging stresses and their combination on leaf photosynthesis, chloroplast ATP synthesis, and antioxidant capacity in wheat." Plant Science **176**(4): 575-582.
- Zheng, C., D. Jiang, et al. (2009). "Exogenous nitric oxide improves seed germination in wheat against mitochondrial oxidative damage induced by high salinity." Environmental and Experimental Botany **67**(1): 222-227.