

Original Research Article

Jaggery yield and nutritional quality as influenced by sugarcane varieties suitable for Andhra Pradesh

Abstract:

Jaggery is produced from sugarcane in addition to sugar. It plays a great role in rural economy of India. However, till date the jaggery producers are dependent on the sugarcane varieties which are released for those areas. The major factor that governing the consumer preference and marketing of jaggery is its external appearances i.e., colour, texture and storability which in turn depend on sugarcane varieties having high sucrose content, purity and low colloids. Hence this study was taken up to identify suitable sugarcane varieties for high yield and good quality jaggery. A study was conducted at Agricultural Research Station, Perumallapalle, Acharya N.G. Ranga Agricultural University, Andhra Pradesh, India to identify the sugarcane varieties most suitable for quality jaggery production. Ten promising sugarcane varieties were planted in randomized block design with three replications. All cultivation practices had followed equally for all varieties. The sugarcane varieties were evaluated qualitatively for pH, EC, reducing sugars, ash content, sucrose, moisture, harvested at proper maturity, crushed to extract juice and prepared jaggery. The jaggery was micronutrients viz., Fe, Mn, Zn and Cu and calculated net rendement value (NRV). The jaggery was found to have 70.6 to 86.9% sucrose, 5.12 to 6.80% of reducing

Remove this unnecessary line.

sugars, 3.52 to 4.32% of ash, 5.0 to 6.1% of moisture, 9.58 to 11.20 % of recovery, 49.85 to 67.70 of NRV values, 98.6 to 114.1 kg jaggery t^{-1} of sugarcane. The jaggery prepared from 2012 T 115 had recorded significantly the highest jaggery yield (114.1 kg t^{-1} of cane), high recovery percent (11.20%) and high NRV (67.70) than other varieties. The varieties viz., 2012 T 183, 2012 T 88 and 2012 T 53 were produced Grade 1 jaggery with high NRV as per Indian standards. The

variety 2012 T 115 had higher Fe and Mn (12.15 and 0.38 mg of Fe and Mn/ 100g of jaggery, respectively).

Keywords: Jaggery yield, sugarcane varieties, mineral and nutritional quality, NRV and South India.

Introduction

Sugarcane is the main source of sweetener's in India. Jaggery is one of the oldest and most important cottage industries in India. Jaggery is a natural, traditional sweetener made by the concentration of sugarcane juice and is known all over the world [1] in different local names [2]. In India, of the total sugarcane produced, 53% is processed into white sugar, 36% into jaggery and khandsari, 3% for chewing as cane juice, and 8% as seed cane [3]. India produces more than 70% of the total jaggery production of the world [4]. Nearly 40% of cane grown in Andhra Pradesh is utilized towards jaggery manufacture. Jaggery is not only used as sweetening agent but also used in several sweet food preparations. In Rayalaseema region of Andhra Pradesh, an appreciable percentage of cane is being utilized for jaggery manufacture. Being an eco friendly sweetener with additional nutritional value jaggery holds good export potential. Jaggery is far complex than sugar, as it is made up of longer chains of sucrose. Hence, it is digested slower than sugar and releases energy slowly and not spontaneously. This provides energy for a longer time and is not harmful for the body. The dietary intake of jaggery prevents the atmospheric pollution related toxicity and the incidence of lung cancer [3]. Quality of jaggery is very sensitive to various parameters such as sugarcane variety, cultivation practices, fertilizers used, stage of harvest, method of juice extraction etc [5, 6]. The composition of extracted juice in terms of its pH, purity, TSS affects the quality of jaggery. To sustain the market and export potential of jaggery it is imperative that the jaggery yield and quality need to be sustained [7]. In general sugarcane farmers face several problems such as low extraction percentage of juice, low recovery and poor quality of jaggery. Quality jaggery comprises of high sucrose and purity with less reducing sugars.

It has been found that there is wide variation in quality of jaggery depending on varieties used in its preparation [8, 9]. Jaggery quality depends mainly on juice and hence factors affecting juice quality also affect the jaggery quality. Studies showed that jaggery quality preferable depends on chemical composition of juice irrespective of method of boiling and clarification [10]. Keeping this in views, a study was taken up to identify a variety with high jaggery yield and good quality of jaggery which is suitable for Andhra Pradesh and South India

Materials and methods

Ten sugarcane varieties viz., 2012 T 53, 2012 T 73, 2012 T 88, 2012 T 98, 2012 T 106, 2012 T 115, 2012 T 180, 2012 T 183, 2003 V 46 and Co C 671 were planted in randomized block design with three replications at Agricultural Research Station, Perumallapalle, ANGRAU, Andhra Pradesh, India. Planting was taken up in the month of February, during both the years (2017 and 2018) and harvested at December of the same year for jaggery preparation. The cane was weighed after harvest and crushed for juice. The extracted juice of all the varieties by using traditional open pan. In jaggery preparation, juice clarification is most important process for acquiring good colour[11]. Clarification of juice was carried out by using bhendi plant extract as natural juice clarificant (250g /120 lit of juice). Lime (40g / 120 lts of juice) was added at the time of boiling of juice during jaggery preparation to bring juice pH to neutral from acidic state.

Give the proper space in between two words. Do this in remain all.

Try to write the full words of spelling.

For analysis of jaggery, 65 g of jaggery sample was weighed and dissolved in 500 ml of water to make a homogeneous solution (0.5M) and then the analysis was carried out with this solution. Brix reading was recorded with brix hydrometer; pol reading was noted by polarimeter using lead acetate as juice clarificant. Physico-chemical properties were assessed by the method developed by [12] . Reducing sugars measured by the method described by [13]. The colour of jaggery was determined by percent transmission of light by using UV-VIS spectrophotometer. The jaggery samples were digested and used for estimation of nutrients (Fe, Mn, Zn and Cu) by

using atomic absorption spectrophotometer (Varian AAS 42). Moisture percent was estimated by the method described by [14].

As per Indian standards (IS 1923; 1990), the jaggery shall be prepared in the form of solid lumps with firm consistency. The colour of jaggery should be golden yellow to light brown, free from dirt, other extraneous matter and also substance harmful to health. It should be sweet to taste and should not be sour, salty or any other objectionable taste. Besides grade I and II (Table 1), jaggery should confirm the characteristics, requirements and methods of test for grading Indian jaggery. Specific characteristics and grading designation of Agmark standard are given in Table 1.

Net rendement value was calculated as $NRV = (\text{Sucrose \%} - \text{Reducing sugars \%}) - 3.5 \times \text{ash \%}$. Based on NRV, the quality of jaggery was considered as follows:

List 1 : Range of Net Rendement Value

Range of Net Rendement Value	Grade	Quality
>65	A1	Excellent
60 - 65	A2	Good
45 - 60	B	Medium
<45	C	Poor

Results and Discussion

The data in Table 2 showed that jaggery yield had significantly affected by all varieties. The variety 2012 T 115 recorded significantly the highest jaggery yield (114.1 kg of jaggery t^{-1} of sugarcane) followed by 2012 T 183 (112.8 kg t^{-1} of cane). This might be due to high recovery percent from cane with 2012 T 115 (11.20%) followed by 2012 T 183 (10.98%) and also due to high cane yield [15, 16]. The sucrose content in jaggery has significantly differed with varieties. The sucrose content in varieties 2012 T 115 (86.9%) and 2012 T 53 (86.5%) was on par with each other. For superior jaggery (Grade 1), the variety should possess high sucrose content (>80%) and low reducing sugars (<10%). The jaggery prepared from all the varieties had low reducing sugars i.e. less than 10% which is one of the important parameter for grade I jaggery. The variety 2012 T

53 had low reducing sugars (5.12%) and it was at par with 2012 T 115 (5.26%). Low reducing sugar is preferable for better quality of jaggery because it is generally high hygroscopic [8]. The moisture content in jaggery of various varieties ranged between 5.2% (2012 T 115) and 8.1% (2012 T 98) (Table 3). According to Indian standard, the jaggery with 5 - 6% moisture content was categorised under Grade 1 (Table 1). Another important feature of the variety for quality jaggery production is low fibre content that should improve the extraction percent of juice and ultimately the recovery percent. Higher jaggery recovery from cane juice was obtained in 2012 T 115 (21.08%) followed by 202 T 183 (20.86%). The purity of jaggery of the varieties 2012 T 183, 2012 T 53, 2012 T 106 and 2012 T 115 was at par with each other. Regarding ash content, 2012 T 115 showed less ash content (3.46%) followed by 2012 T 106 (3.52%). Electrical conductivity of jaggery which shows salt content owing to its hygroscopicity is lowest in 2012 T 183 (0.182 dS m⁻¹) followed by 2012 T 115 (0.208 dS m⁻¹) (Table 3). Highest EC values were recorded with CoC 671, Co 94008, 2012 T 53 which are more hygroscopic compared to other varieties. Higher values indicated that susceptibility of jaggery for moisture absorption. As far as colour is concerned, the most suitable varieties were found to be 2012 T 115 (20.50), Co 94008 (21.56) and 2012 T 53 (21.60). These varieties could be used for producing light golden coloured jaggery. The production of attractive colour jaggery in turn depends on the extent of clarity of sugarcane juice. The fresh cane juice contains appreciable quantity of colloidal impurities and these are to be removed for manufacturing the quality jaggery. The pH of the juice determines the crystalline texture of the jaggery. The different varieties of sugarcane were showed non significant effect on pH of jaggery. The quality crystalline jaggery can be produced by adjusting the pH above 6.0. The natural sugarcane juice has a pH of 5.5. It can be adjusted by using the lime for this purpose. The low levels of pH cause inversion by hydrolyzing the sugar and thus affecting the jaggery quality [7, 16].

The higher NRV was obtained with varieties 2012 T 115 (67.70), 2012 T 106 (66.15), 2012 T 183 (66.78) and 2012 T 88 (65.69). It indicated that these varieties grouped under A1 grade with excellent quality jaggery [17]. The Fe and Mn content in jaggery were significantly affected by varieties (Table 4). The Fe content from jaggery produced with 2012 T 115 (12.15 mg/100g) was at par with 2012 T 183 (11.92 mg/100g). However high Mn content (0.39 mg/100g) was recorded with 2012 T 183 and it was at par with 2012 T 115 (0.38mg/100g). The Zn and Cu content in jaggery produced from different varieties was not significantly influenced by varieties [18].

Conclusion

At present, jaggery is graded at national level on Agmark system of solid jaggery grading (based on physical characteristics) and BIS standards IS:1923. Out of ten varieties 2012 T 115 proved to be suitable regarding maximum jaggery yield, high NRV, low electrical conductivity, low reducing sugars and light colour. The study revealed that quality grade 1 jaggery with high sucrose, low ash content, low reducing sugars were obtained from 2012 T 183, 2012 T 88 and 2012 T 53. The variety 2012 T 115 had higher Fe and Mn (12.15 and 0.38 mg of Fe and Mn/ 100g of jaggery, respectively).

Kindly give the space for paragraph making.

Adjust the making of spaces. In remain all articles.

Table 1: Standard specifications for cane jaggery

Parameters	Requirements for	
	Grade I	Grade II
Sucrose % (minimum)	80	70

Reducing sugars % (maximum)	10	20
Moisture % (maximum)	5	7
suphated ash % (maximum)	3.5	5.0

Kindly write correct spelling.

Table 2: Quality parameters of jaggery affected by elite sugarcane varieties

Variety	Jaggery yield (kg t ⁻¹)	Sucrose (%)	Purity (%)	Reducing sugars (%)	Moisture (%)	Recovery % from cane	Recovery % from juice
2012 T 53	109.8 ^c	86.5 ^a	92.5 ^a	5.26 ^a	7.6 ^c	10.25 ^c	20.54 ^a
2012 T 73	101.3 ^{de}	81.2 ^{cd}	86.5 ^d	5.94 ^{bc}	6.0 ^b	10.86 ^a	19.25 ^{ab}
2012 T 88	112.2 ^b	84.5 ^b	90.2	6.35 ^d	7.8 ^c	10.02 ^{cd}	20.80 ^a
2012 T 98	99.12 ^f	70.6 ^e	88.5 ^{bc}	6.0 ^c	8.1 ^{de}	9.98 ^{cd}	17.26 ^e
2012 T 106	107.6 ^c	84.4 ^b	91.6 ^a	5.92 ^{bc}	7.4 ^c	10.56 ^b	20.04 ^{ab}
2012 T 115	114.1 ^a	86.9 ^a	91.0 ^a	5.12 ^a	5.2 ^a	11.20 ^a	21.08 ^a
2012 T 180	102.7 ^d	79.1 ^{cd}	92.0 ^a	6.74 ^e	6.8 ^b	10.57 ^b	18.55 ^{cd}
2012 T 183	112.8 ^b	86.6 ^a	92.8 ^a	5.81 ^b	6.2 ^b	10.98 ^a	20.86 ^a
Co94008	101.5 ^{de}	72.5 ^d	89.2 ^{bc}	6.01 ^c	8.0 ^{de}	9.94 ^{cd}	19.36 ^{bc}
CoC 671	98.6	77.2 ^{cd}	84.1 ^e	5.96 ^{bc}	5.5 ^a	9.58 ^e	18.62 ^{cd}
Treatments	*	**	*	*	*	*	*
p-value	0.042	0.008	0.032	0.027	0.014	0.034	0.044

Table 3: Physico chemical properties and recovery percent of jaggery as influenced by sugarcane varieties

Variety	Ash (%)	pH	EC (dS m ⁻¹)	Colour intensity (OD value)	NRV
---------	---------	----	--------------------------	-----------------------------	-----

2012 T 53	3.46 ^a	5.82	0.264 ^{bc}	21.60 ^{ab}	62.83
2012 T 73	4.12 ^{de}	5.77	0.251 ^b	23.53 ^{cd}	60.84
2012 T 88	3.56 ^a	5.60	0.260 ^{bc}	22.47 ^{ab}	65.69
2012 T 98	4.21 ^{de}	5.74	0.258 ^{bc}	29.10 ^e	49.85
2012 T 106	3.52 ^a	5.94	0.250 ^b	30.40 ^e	66.18
2012 T 115	3.91 ^{cd}	5.81	0.208 ^a	20.50 ^a	67.70
2012 T 180	3.89 ^{cd}	5.77	0.262 ^d	22.10 ^{ab}	58.75
2012 T 183	3.72 ^{ab}	5.87	0.182 ^a	23.13 ^{cd}	66.78
Co94008	4.32 ^e	5.34	0.288 ^{de}	21.56 ^{ab}	51.36
CoC 671	4.02 ^d	5.91	0.352 ^e	25.42 ^d	57.17
Treatments	*	N.S.	*	*	*
p-value	0.041	0.102	0.034	0.048	0.042

Table 4: Nutrient content (mg /100g of jaggery) of jaggery prepared from elite sugarcane varieties

Variety	Fe	Zn	Mn	Cu
2012 T 53	10.85 ^b	0.48	0.32 ^{ab}	0.28
2012 T 73	11.62 ^{ab}	0.52	0.28 ^{bcd}	0.20
2012 T 88	8.62 ^c	0.46	0.35 ^a	0.26
2012 T 98	9.74 ^{bc}	0.59	0.26 ^{bcd}	0.30
2012 T 106	7.58 ^d	0.41	0.31 ^{ab}	0.27
2012 T 115	12.15 ^a	0.50	0.38 ^a	0.24
2012 T 180	10.56	0.41	0.30 ^{ab}	0.32
2012 T 183	11.92 ^a	0.48	0.39 ^a	0.28
Co94008	8.20 ^{cd}	0.50	0.22 ^{cd}	0.26
CoC 671	7.02 ^e	0.44	0.32 ^{ab}	0.24
Treatments	*	N.S.	*	N.S.
p-value	0.026	0.094	0.041	0.124

Arrange references uniform way as per instructions.

References:

1. Priyanka S, Amit B, Singh MM. Validation of elite sugarcane varieties for quality jaggery production in subtropical India. Sugar Tech. 2018 Jul 13, <https://doi.org/10.1007/s12355-018-0647-6>.
2. Thakur AK. Potentiality of jaggery (Gur) manufacturing in Punjab state. Proceedings of the National Seminar on Status, Problems and Prospects of Jaggery and Khandsari Industry in India, Indian Institute of Sugarcane Research. Lucknow, India. 1999. 95-96.

3. Rao PVKJ, Das M and Das SK. Jaggery - A traditional Indian sweetener. Indian Journal of Traditional Knowledge 2007. 6: 95-102.
4. Singh J, Singh RD, Anwar SI, Solomon S. Alternative sweeteners production from sugarcane in India: Lump Sugar (jaggery). Sugar Tech 2011. 13: 366-371.
5. Nath A, Dutta D, Kumar P and Singh JP. Review on Recent Advances in Value Addition of Jaggery based Products. J Food Process Technol 2015. 6: 440. doi:10.4172/2157-7110.1000440
6. Nevkar GS, Patil DS and Nale VN. Studies on standardization of jaggery powder production process. J Agric Res Technol 2011. 36(3): 490-492.
7. Dilip. A. Pawar, Maruti. S. Jadhav, Charudatta A. Nimbalkar. Techniques and Advances in Jaggery Processing: A review. Res. J Chem Env Sci 2017. 5 (3) June 2017. 14-20
8. Rakiyappan T and Janki P. Jaggery quality of some commercial and promising sugarcane varieties. Co operative Sugar 1996.. 27(12):909-913.
9. Singh P, Singh M and Sharma BL. Screening of sugarcane varieties for quality jaggery in North India. In proceedings of strategic governance and technological advancement for sustainable agriculture, 4th Uttar Pradesh agricultural science congress, Kanpur. 2016. 2-4 March, Pp:45.
10. Singh J, Solomon S and Kumar D. Manufacturing jaggery: a product of sugarcane as health food. Journal of Agro techno 2013. S11:1-3.
11. Mungare TS, Jadhav HD, Patil JP, Hasure RP, Jadhav BS and Singh J. Clarification technique for producing quality jaggery. Cooperative Sugar 2000. 32(4): 283-285.
12. Spencer GL and Meade GP. Standard procedure for juice analysis. Cane Hand Book, London: Willey. 1945.
13. Hulme AC and Narain R. The ferricyanide method for the determination of reducing sugars: A modification of the Hagedorn-Jensen-Hanes technique. Biochemical journal 1931. 25 (4): 1051.
14. Jain PC and Singh P. Moisture determination of jaggery in microwave oven. Sugar Tech 2000. 2(3):51-52
15. Mishra A. Evaluation of early maturing promising sugarcane varieties for jaggery yield and quality. 1991. Ind. Journal of Agronomy 36: 315-316.
16. Madan HK, Jaiswal UK, Kumar JS and Khanna SK. Improvement in gur (jaggery) making plant for rural areas. Journal of rural tech 2004. 194-196.

17. Sharma SC, Johary PC and Rao GSC. Some observations on the chemical composition of cane juice of different promising varieties grown in North Bihar condition and their classification characteristics. Indian Sugar 1979. 29(3):141-147
18. Shweta, Pawar K, Gehlot R, Kumar R and Sudhir R. Screening and physico chemical study of quality jaggery prepared from different early and mid season sugarcane varieties.. Asian journal of dairy and food Res 2022 (41): 56-63

UNDER PEER REVIEW