

## Original Research Article

### **NEW INSIGHT INTO A SUBDUCTION-RELATED OROGEN: A REAPPRAISAL OF THE GEOTECTONIC FRAMEWORK AND EVOLUTION OF THE MIDDLE AND WEST PARTS OF THE SOUTHEAST ANATOLIAN OROGENIC BELT (TÜRKİYE)**

#### **ABSTRACT**

The geotectonic framework and the evolutionary history of the Southeast Anatolian Orogenic Belt are closely related to the assemblage of eastern and western Gondwana and the subsequent events from the opening of the southern branch of the Neo-Tethys to the final collision. The first geotectonic event is the subduction of the prototethys under the northern Gondwana during the Ediacaran and accordingly the formation of igneous rocks within the lower units of Bitlis-Pütürge Massifs. The first orogeny affecting the region was the Cadomian orogeny. The southern branch of the Neo-Tethys began to open between the Arabian plate North of Gondwana) and today's southeastern Anatolian metamorphic massifs in the Late Triassic, and oceanic spreading continued until the late Cretaceous. The ophiolites and an intra-oceanic arc were formed during the Late Cretaceous (92 to 82Ma and 84–72 Ma respectively) in a SSZ tectonic environment formed by the northward subducting South branch of Neo-Tethys ocean crust. The Arabian platform entered the subduction zone and as a result ophiolites thrust on the Arabian plate margin, the metamorphic massifs were fragmented and migrated to the South onto the ophiolites and arc magmatics in the Maastrichtian. Despite the collision, the continental subduction continued and a break-off of subducted slab was formed. A widespread marine transgression is realized onto the Arabian Platform and ophiolites from latest Cretaceous to Early Miocene to the South of the Bitlis-Pütürge metamorphics. The remnant of the ocean continued until Late Miocene to the North of the Bitlis-Pütürge massifs as marine basins with different depths and morphological characteristics. The magma formed by the partial melting of the mantle wedge, the rising deep asthenosphere mantle and the continental crust forms Maden arc over the ophiolites and the Bitlis-Pütürge Massifs in the Middle Eocene. Behind the Maden arc, shallow-deep marine carbonates and clastics were deposited in a back-arc basin (Kırkgeçit basin). The closure which started in the Late Eocene and ended in the Late Miocene enabled Southeast Anatolian Orogenic Belt to take its actual position.

**Key words:** Southeast Anatolia, Neo-Tethys, tectonics, geological evolution, subduction

## **1.INTRODUCTION**

The Southeast Anatolian Orogenic Belt (SEAOB) forms a belt over 1000 km in length from Iskenderun Bay to triple joinction of Türkiye-Iran-Irak between the Arabian Platform and the Anatolian (Türkiye) microplate. There are 4 major units in the SEAOB: a) metamorphic massifs, b) ophiolites and arc related rocks, c) Maden Complex, d) Upper Cretaceous–Neogene cover units. The geological studies of these units prior to the 1990s were largely based on the studies in the field (relationships between different units) and limited number of chemical analyzes [9, 10, 14, 35-38, 52, 61, 65, 67, 70, 73- 75, 92, 104, 106, 125, 137, 140, 145, 157, 165- 169, 185, 187, 194]. Therefore, in these studies before the 1990s, the geotectonic models of the region were created according to these insufficient data. Since it was generally accepted that like other ophiolites in the world, the Turkish ophiolites also were formed at the ocean floor spreading center and thrust on the continental crust. Pearce al. [135] indicated that many of the world's best-known ophiolites have petrological and geochemical characteristics that suggest formation above a subduction zone (supra subduction-zone; SSZ). After this acceptance, the formations of ophiolites cropping out in the region have also been reinterpreted according to this theory [27-33, 43, 46, 47, 83-85, 96, 130, 133, 146, 149, 150, 155, 177, 178]. Various geochemical analyzes, isotope studies, and geochronological data on metamorphic massifs [33, 46, 109, 177, 178], ophiolites and arc-related magmatic rocks [28, 32, 43, 46, 47, 84, 85, 96, 130, 133, 146, 149, 150, 156] were also used in the development of these new models.

The purpose of this study is, by Our field studies for 40 years, our observations based on the relationships between different units, our geochemical-geochronological data and using the data of researchers working for different purposes in the region, to revise the geotectonic framework and the evolutionary history of middle and west parts of the Southeast Anatolian Orogenic Belt.

## **2. MAJOR GEOTECTONIC UNITS**

The Southeast Anatolian Orogenic Belt (SEAOB) is located between the Arabian Platform and the Anatolian microplate and is separated from it by the Southeast Anatolian Thrust Belt (Bitlis-Zagros Suture Zone). The Arabian Platform which represents the northwestern part of the Arabian Plate, a part of the SEAOB. The Arabian foreland located at the south of the Bitlis-Zagros suture zone has a basement composed mainly of Precambrian

rocks, overlain with a thick pile of shallow water sedimentary formations of Early Paleozoic to Miocene ages [164].

Except for the Arabian Platform, the SEAOB is mainly composed of the Neoproterozoic to Early Cenozoic orogenic elements, i.e. regional metamorphic rocks, ophiolites, arc-related magmatics, and volcanic and sedimentary rocks that align roughly parallel to the general trending of the SEAOB (Fig.1).




Figure 1- Major geological units of Southeast Anatolian Orogenic Belt (simplified from MTA, 2004).

## 2.1. Metamorphic Massifs

The metamorphic massifs named as Southeast Anatolian metamorphic complex by Ketin [92], include Bitlis-Pütürge- Engizek-Keban-Malatya and Binboga Massifs. Bitlis-Pütürge-Engizek massifs forming the southern belt form subparallel units to the Keban-Malatya-Binboğa massifs forming the northern belt

### 2.1.1. Bitlis Pütürge-Engizek Massifs:

The Bitlis-Pütürge-Engizek Massifs are an arcuate belt of allochthonous metamorphic massifs. This belt is approximately 30 km wide, dipping northwards at low to moderate angles, and they extend approximately parallel to the southeast Anatolian thrust belt

(Fig.1). They are separated from Arabian Platform by a narrow tectonic belt consisting of ophiolitic and flysch units [28, 39, 43, 73, 117, 137, 138, 195]. The massifs consist of a Neoproterozoic-Cambrian high grade metamorphic lower unit and a Devonian-Triassic lower-grade metamorphic cover. The lower unit consists of granitoid gneiss, amphibolite, and mica-schists [33, 51, 62, 63, 71, 116, 118, 177].  $^{207}\text{Pb}/^{206}\text{Pb}$  single-zircon age determinations on the metagranites from meta-granites in the Bitlis massif (Ustaömer et al. 2009; 2012) and on the augen gneisses in the Pütürge Massif [33] reveal that they crystallize at an age range of 572-5230 Ma (Ediacaran–Early Cambrian). Geochemical characteristics of the metagranites and augen gneisses suggest the existence of Andean type arc-related magmatism. The augen gneisses the  $\epsilon\text{Hf}(t)$  values of augen gneisses suggest the involvement of older continental crust in magma genesis [33].

Devonian volcanics outcropping in the Arabian platform south of the thrust zone and contemporaneous age with augen gneiss and meta granites were formed in a back arc basin [72].

The lower unit underwent high-grade metamorphism by the closure of the proto-tethys and final amalgamation of exotic terranes during the Cadomian orogeny the northeastern part of Gondwana [49, this study). Paleozoic-Lower Mesozoic metamorphic platform sediments unconformably overlie the lower unit. There is no sedimentary or igneous rock yielding Cambrian-Ordovician aged units. The first marine clastic and carbonate rocks overlying the lower unit are of middle Devonian age [71]. This cover unit consists of muscovite schist containing mid-Devonian fossils with kyanite-bearing quartzite lenses, garnet staurolite mica schists, and Permian recrystallized limestones [63, 71]. Late Triassic characterized by radiolarite meta-mudstone, meta-basalt meta-tuff and meta-shale indicates that the sea deepened suddenly and the region rifted. This rifting marks the opening of the southern branch of the Neo-Tethys. Rifting occurred between the Arabian platform and the present metamorphic massifs. Present metamorphic belt thereafter remained as submerged continental margin up to the Late Campanian-Early Maastrichtian. The lower-unit and the upper unit both together rocks were metamorphosed under greenschist facies conditions during the Upper Cretaceous [63, 188].

#### 2.1.2. Keban-Malatya Metamorphics

The Malatya metamorphics cropping out in the west of Malatya and the Keban metamorphics cropping out around Keban-Baskil (Elazığ) and Pertek (Tunceli) areas display similar successions.

The Malatya Metamorphics crop out between Malatya and Kahramanmaraş, and consist of meta-carbonates, mica schist, phyllite, meta-clastic rocks, and meta-cherts [140]. Özgül et al. [124], based on the very limited number of fossils they have found, accept that the Malatya metamorphics were formed in the Late Permian to Early Triassic. The Malatya metamorphics cropping out around Gölbaşı, the Karanlıkdere ophiolite which is an extension of the Koçali ophiolite, outcrops in the south and the Berit ophiolite in the north. The contacts between ]metamorphics and ophiolite are tectonic. The tectonic contact between metamorphics and Karanlıkdere ophiolite dips to the north and the metamorphics thrust over the ophiolites [42, this study] and the tectonic contact between the metamorphics and Berit ophiolite is a southward dipping [154]. This situation shows that the Malatya metamorphics are found as an allochthonous unit over the ophiolites in the north and south and that the two ophiolites are the continuation of each other under the metamorphics (Fig. 2).




Figure 2- Cross-section between Doruk M. and Koçali.

A tectonic relationship is observed between the Malatya metamorphics and the Pütürge metamorphics in the vicinity of Çelikhhan (Adıyaman) [190]. This shows that the Malatya metamorphics and the Pütürge metamorphics are parts of the same large massif. They were metamorphosed in greenschist facies during the Late Cretaceous [36, 155, 196].

The Keban metamorphic rocks consist of meta-carbonates, meta-conglomerates meta-sandstone, and phyllite-chlorite-sericite schist with intrusions of meta-diabase dykes [36, 90]. Keban metamorphics thrust over the Late Cretaceous Elazığ magmatics in the south. The northward dipping of the thrust is cut by the granitoid of the Elazığ magmatics. The granitoid also intrudes the Keban platform Permo-Carboniferous carbonate deposits NW of Birivan (Ulupınar) village. The contact between granitoid and metamorphosed Keban Platform carbonates displays well-exposed hornfels and skarn rocks. Such a skarn contact is also found

SE of Aşvan village, near the Keban Dam, between a diorite intrusion and the Keban marbles with a large magnetite mineralization. Keban metamorphics also crop out tectonically over the late Cretaceous aged Elazığ magmatics around Pertek. The tectonic contact between the two units is covered by the Eocene Kırkgeçit Formation to the east and west of Pertek [36]. Keban metamorphics are found as small allochthonous blocks over the Elazığ magmatics in the vicinity of Elazığ.

Kipman [93] suggested that the age of the Keban metamorphics is Permo-Carboniferous according to fossils *Glomospira* and *Ammodiscus* families identified in the crystallized limestones.

All metamorphics of SEA OB were metamorphosed in greenschist facies due to northward subduction of the southern branch of the Neo-Tethys in the Late Cretaceous.

Field data indicate that all metamorphic massifs in the SEA OB are parts of an once-united giant tectonostratigraphic unit [43, 197].

## **2.2.Ophiolites**

SEA OB ophiolites are an important part of the 3000 km long Neotethys ophiolite belt extending from Italy to Oman. Neotethys is divided into two branches in the region where today's Turkey and Iran are located. These are 1) main branch, 2) south branch. While the Neotethys ophiolites are mostly MORB type in the west, they are around 170-140Ma aged (e.g., Ligurian in Italy, Mirdita in Albania, Pindos in Greece, Refahiye in Turkey, and Makran in Iran), those in the central and eastern parts show typical SSZ geochemical signatures and 125-86Ma aged (Troodos in Cyprus, Kızıldağ, Koçali, Ispendere, Kömürhan–Guleman in southeastern Turkey, Neyriz in Iran, and Oman) [1, 43, 69, 108, 179]. Many researchers have conducted studies on the SEA OB ophiolites locally or regionally [21, 22, 26-31, 37, 39, 43, 47, 56, 58, 59, 83-86, 98, 100, 125, 130-134, 148-150, 180]. The SEA OB ophiolites extend for approximately 1000 km from the Iskenderun Bay in the West to the Turkey-Iran-Iraq triple intersection in the East and include the Kızıldağ, Koçali, Ispendere, Kömürhan, Guleman, Gevaş, Cilo ophiolites, and numerous small unnamed ophiolite fragments (Fig.3).

The Kızıldağ ophiolite, located in the westernmost part of SEA OB, was thrust over the thick Cambrian-Cretaceous autochthonous Arabian platform and is unconformably overlain by the Late Maastrichtian–Late Miocene autochthonous sediments [157]. The Kızıldağ ophiolite contains all of the lithological units seen in an ideal ophiolite succession: harzburgitic mantle peridotite, the dunitic mantle-crust transition zone (DTZ), ultramafic-mafic cumulates, sheeted dykes and volcanic rocks [43, 47, 56, 58, 59, 83- 86, 131- 134].




Figure 3- Distribution of the ophiolites in the study area (From Bingöl et al. 2018)

The Koçali ophiolite is a part of the Koçali complex which consists of the Triassic Tarasa volcanic rocks, the Konak formation, and the Late Cretaceous Kale formation [136, 137]. The Kale Formation extends towards the east to the vicinity of Çermik. Further east, the unit cropping out around Çermik (Diyarbakır), containing an ophiolite sequence and showing the same characteristics as the Kale Formation, was named Koçali ophiolite by Bingöl [39]. The unit corresponding to the Kale formation, which crops out to the west of Gerger and consists of mantle peridotites, cumulates, diabase dyke complex and basalts, was named as Koçali ophiolite [28]. The Tarasa volcanics and the Konak formation were thrust onto the Koçali ophiolite. The Koçali ophiolite has been thrust onto the Upper Campanian Karadut Complex. The Tarasa volcanic rocks, the Konak formation and the Koçali ophiolite stratigraphically overlain by Upper Maastrichtian-Eocene sedimentary units of the Arabian Platform [28] and the Çüngüş Formation and Pütürge metamorphics overlie them tectonically in the West of Sincik (Adıyaman) [190] (Fig. 4).

Some ophiolite fragments were tectonically overlain by the Çüngüş Formation belonging to Arabian Platform and the Çüngüş Formation was thrust by the Pütürge metamorphics. The Karanlık Dere ophiolite cropping out to the East of Gölbaşı (Adıyaman) is



an extension of the Koçali ophiolite under the neo-autochthonous cover of the Arabian Platform [42, 43]. The Malatya Metamorphics cropping out in the north of the Karanlık Dere tectonically overlain the Karanlık Dere ophiolite in the south and the Berit ophiolite in the north. Well, the Malatya Metamorphics overlie the ophiolites as allochthonous unit (Fig.2). Therefore, the Berit ophiolite and the Koçali ophiolite are parts of the same ophiolite (Fig.2). Between the Kızıldağ ophiolite and the Koçali ophiolite, a large number of unnamed ophiolite fragments crop out under young sediments and volcanics. The Koçali ophiolite consists of harzburgitic mantle peridotites, gabbros, plagiogranite, sheeted dikes, and basalts. In the Çermik anticline, arc-related volcano-clastics are observed on the ophiolite, while in the Karanlık Dere, the ophiolite is cut by granitoid dykes.




Figure 4 Cross-section between Aşağıçimen (Sincik-Adıyaman) Çat Dam (Çelikhan-Adıyaman)

The Guleman ophiolite cropping out in the Southeast of Elazığ shows a very different tectonic situation. The Guleman ophiolite thrust over the Late Miocene Lice formation, which is the youngest unit of the Arabian Platform. It is thrust by the Pütürge metamorphics in the South of the Hazar Lake, and by the Bitlis metamorphics in the Northeast. It is depositionally overlain by the Late Maasstrichtian–Early Eocene Hazar Group and middle Eocene Maden Complex [43, 64, 127, 137, 139, 145]. Guleman ophiolite presents an ideal ophiolitic sequence consisting of mantle peridotites, dunitic mantle-crust transition zone (DTZ), ultramafic-mafic crustal rocks [43, 149]. The basalts called Caferi volcanics by Özkan and Öztunalı [125] are controversial. However, the Guleman ophiolite and arc magmatics developed on the ophiolite are examined in detail, it is seen that these basalts form the uppermost part of the ophiolite and also form the base of the arc magmatics (Fig.5 and 6).






Figure 5- Cross-section Between Alacakaya (Elazığ)-Pertek (Tunceli)




Figure 6- Cross-section Between Çermik (Diyarbakır)-Keban (Elazığ)

The Kömürhan and Ispendere ophiolites, which are the western extension of the Guleman ophiolite, thrust over the middle Eocene Maden Complex developed over the Pütürge metamorphics, and are overlain by the arc-related magmatics (Fig.7). Mantle peridotites are missing in both ophiolites. They consist of ultramafic cumulates, cumulate gabbros intruded by ultramafic-mafic dykes and stocks, sheeted dykes, basaltic pillow lavas, and lava flows. While the Kömürhan ophiolite is cut by granitic dikes, a tectonic relationship is observed between the Ispendere ophiolite and granitic rocks [27, 29, 43]. The lower part of the Kömürhan ophiolite containing amphibolite, pyroxenite, and garnet-peridotites is metamorphosed in greenschist and amphibolite facies [188]. The granitic dykes of the arc magmatites cut the Kömürhan ophiolite and form contact metamorphism around it. [27, 29].

The U-Pb zircon datings provide the ages 92 to 82 Ma for the Southeastern Taurus ophiolites [43, 83, 84, 98]. These ages indicate that ophiolites were formed in a maximum

time period of  $\sim 10$  Ma [43]. The whole-rock geochemical, geochronological, and isotropic data described by different researchers [28, 43, 47, 56, 58, 59, 83-86, 100, 101, 131- 134, 149, 150] strongly suggest that the Southeast Anatolian ophiolites were generated in SSZ tectonic settings during the Late Cretaceous.




Figure 7- Cross-section Between Çermik (Diyarbakır)- Baskil (Elazığ)

### 2.3.Arc-related magmatic rocks

Perinçek [139] suggests that a very complex unit consisting of Late Cretaceous basic-andesitic igneous rocks and sedimentary rocks has been named the Yüksekova complex at the easternmost end of SEAOb, and a group of rocks similar to this unit also crops out around Elazığ-Malatya. Hempton and Savcı [77] showed that this unit predominantly consists of igneous rocks and named the Elazığ complex in the Elazığ area. Hempton [75] also named this unit the Elazığ volcanic complex. We carried out detailed studies and geological mapping for the first time on this unit in the north-northeast of Elazığ, and we published the results of these studies in two articles [35, 36]. Although we used the name Yüksekova Complex in these articles, in later studies it was revealed that the unit was not a complex and therefore named it as Elazığ magmatics [172]. Some researchers, on the other hand, named the plutonic rocks in different regions of the unit with the name of that place; e.g: Baskil batholite [188]), Baskil granitoid [146, 148], Baskil magmatic[17], Pertek granitoid [96], Şifrin group [190]. Ural et al. [176] use the name Yüksekova complex.

Elazığ magmatics crop out most widely between the Elazığ and Malatya provinces. The unit crops out in a very wide area around the city center of Elazığ, between Kovancılar, Keban, Baskil districts, Pertek (Tunceli), Kale (Malatya) districts [4, 15, 16, 26, 27, 29, 31,

32, 35, 36, 38, 40, 41, 43, 96, 98, 146, 156, 176] (Fig. 8). The most comprehensive study on the unit was given by Beyarslan and Bingöl [2018].




Figure 8- Geological map of the Late Cretaceous magmatics in the Elazığ-Malatya regions. (From Beyarslan and Bingöl, 2018).

The Elazığ magmatics mainly overlie ophiolites. The relationships between Elazığ magmatics and ophiolites are sometimes tectonic and sometimes transitive. The Elazığ magmatics thrust over the middle Eocene Maden Complex to the South and North of Hazar Lake and to the South of the Elazığ-Bingöl highway (Fig. 9).

The Keban metamorphics thrust onto the Elazığ magmatics in the Keban and Pertek regions and the granitic rocks of the Elazığ magmatics cut the thrusting contact. It can be observed some small skarn zones at this contact [36]. While the Late Cretaceous Harami formation transgressively overlies the Elazığ magmatics, the Early Paleocene Kuşcular, Late Paleocene-Early Eocene Seske [94], and Upper Bartonian-Priabonian Kırkgeçit Formations [91] unconformably overlies the Elazığ magmatics. The Kırkgeçit Formation also overlies the tectonic contact between the Keban metamorphics and the Elazığ magmatics in the vicinity of Pertek town [36].




Figure 9- Cross-section between Ergani (Diyarbakır)- Perisuyu (Tunceli).

The Elazığ magmatics consist of volcanic and volcano-sedimentary rocks and intrusive rocks. Volcanic rocks outcropping in a wide area between Elazığ city center-Kovancılar and Pertek composed of basalt lavas and lava flow, basaltic andesite, andesite, andesite-pyroclastic alternation dacite and occasionally rhyolite [32, 35, 36, 38, 41]. Fine-grained andesite and dacite dykes transect the basalts. The andesitic lava flows overlie the basaltic rocks. Andesites starting with lava flows pass upwards andesite/pyroclastic alternation and then continue with pyroclastics. The pyroclastic rocks consist of agglomerate, pyroclastic breccia, lapillistone and tuff. The dacitic dykes cut vertically all the alternation of andesite/pyroclastic rocks. These dykes, 0,5 to 2 meters thick and 100 to 200 meters long, feed small dacite domes. The dacitic and rhyolitic rocks are relatively infrequent. The andesitic to rhyolitic rocks are characterized by the calc-alkaline series. Lin et al. [98]), who made a detailed U-Pb age determination on Elazığ magmatics, reported that the volcanic rocks were formed in the range of 84-81 Ma. According to Beyarslan and Bingöl [32], the age of andesite determined by the  $^{206}\text{Pb}/^{238}\text{U}$  method is 82 Ma. Karaoğlu et al. [84] give an age of 74 Ma for a rhyolite sample on the Kömürhan ophiolite and 83 Ma for a sample of a rhyolite on the Göksun ophiolite.

Intrusive rocks crop out most commonly in the North-Northeast of Elazığ city, in the vicinity of Pertek (Tunceli), in the Baskil district (Elazığ) and Ispendere (Malatya). In addition, these rocks crop out in the region between Çelikhan and Sincik (Adıyaman) [190]. Intrusive rocks consist of a wide lithological composition from gabbro to granite and at lesser



rates monzonite-syenite. Different lithological units intersect each other and also other units such as basaltic volcanics and metamorphics, ophiolites. According to the lithological features in the regions they examined, Akgül [4] divided them into diorite and tonalite groups, Sar [156], on the other hand, divides them into granitic and dioritic groups. Lin et al. [98] divide the Elazig magmatics into three groups according to their crystallization ages and magmatic series. They are: (1) 84-81 Ma: tholeiitic suite that consists of extrusive (basalt and andesite) and intrusive (gabbro and diorite) rocks: (2) 80-79 Ma: calc-alkaline suite of monzonite, granodiorite, and granite; (3) 74-72 Ma: calc-alkaline suite of intrusions (gabbro, monzodiorite and monzonite). Beyarslan and Bingöl [32] group Elazig magmatics as (1) first-stage intrusions—mostly gabbro-diorite-tonalite and a lesser granodiorite- granite, 2) second-stage intrusions—mostly granodiorite-granite and a lesser tonalite, 3) late-stage intrusions—mostly monzodiorite- monzonite-syenite subgroup) subgroups. The granodiorite and granite of the second stage have intruded into ophiolites, volcanic rocks, and first group rocks. They also cut the tectonic contact between the Keban metamorphics and Elazığ magmatics [4, 29, 32,36, 38 40, 172]. The late-stage subgroup crops out at the North of Elazığ city and between Çelikhan and Sincik (Adıyaman). These outcrops of late-stage intrusive rocks named Şifrin group by Yıldırım [190] in the Çelikhan region are mostly composed of monzonite. However, in different proportions, gabbro, diorite, quartz diorite, tonalite, quartz monzonite, syenite, quartz syenite, granite, granodiorite, monzodiorite, quartz monzodiorite are also found. In the studies carried out by Pişkin [142], these rocks were named leucocratic quartz monzonites and according to K/Ar measurements, it is indicated that these rocks are 62 Ma or older. Esence granitoid cropping out between Göksun and Afşin (Kahramanmaraş) [119-121] was intruded into the Malatya Metamorphics and Late Cretaceous Göksun Ophiolite. The 85 to 77 Ma ages obtained by the K-Ar method [83, 130] from Esence granitoid, display that this granitoid is the western extension of the intrusive rocks of the Elazığ Magmatics.

Zircon U-Pb age determination made from Elazig magmatics with low K-tholeiitic, calc-alkaline, and shoshonitic series features that the arc-related magmatism took place between 84–72 Ma in an intra-oceanic arc-system that developed on a northwardly dipping oceanic crust in the Upper Cretaceous.

## 2.4. Maden Complex

The unit that is widely exposed in SEAOb has been different named by different researchers: i.e “Maden Unit” [145] “Maden Complex” [10, 75, 141, 186, 187], “Baykan

Complex” [159], “Karadere Formation” [2], “Maden Formation” [51, 104, 127], and “Maden Group” [64].

The unit is located in different positions compared to other units. It is located under the Bitlis-Pütürge metamorphic massifs to the South, while it unconformably overlies the Pütürge massif on the Malatya-Pütürge road. In the district of Maden, from which the unit is named, the Maden Complex unconformably overlies the Guleman ophiolite and the Hazar group. In the same region, it is observed that the Guleman ophiolite and the Hazar group were thrust over the Maden Complex (Fig 4 and 10).




Figure 10. Cross-section in the south and southeast of Lake Hazar.

The Maden Complex has a complex internal structure. The complex begins with transgressive sediments continues upwards with sandstone, silicified red chert, and red-colored mudstones. Lateral and vertical lithological discontinuities are very common in the complex. The red-colored marly and clayed limestone is interbedded by volcanic rocks. The Ypresian-Lutetian clastic sediments unconformably overlie the Pütürge metamorphic massif on the Malatya-Pütürge road [62, 63, 188]. There are many olistrostromal parts consisting of Upper Lutetian limestone, andesitic epiclastics, sandstone, volcanic, and diabase blocks. The uppermost of the unit is composed of basalt, andesite, pyroclastic, and hypabyssal rocks. The hypabyssal rocks composed of diabase and tonalite intruded through Bitlis-Pütürge metamorphic basement [51, 142, 190, 193]. Large tectonic lenses of tourmaline-bearing

micro-leucogranite occurs above the autochthonous sediments (Çakçak Tepe-Malatya), and in the Pütürge metamorphic rocks [188]. Moreover, tonalitic and andesitic vein rocks belonging to the Maden Complex are intruded into pütürge metamorphics at Sakız, Çakçak Tepe-Gazitahara Tepe and Baizge regions [190, 193]. The enrichment of large-ion lithophile elements (LILE), depletion of high field strength elements (HFSE), and positive Pb and negative Nb-Ta anomalies [68] indicate that magma yielding the volcanic rocks and dykes of the Maden Complex derives from the a lithospheric mantle source affected by continental contamination.

## **2.5.Upper Cretaceous–Paleogene Cover Units**

As we briefly summarized above, sedimentary units of the Arabian platform crop out in the south of Pütürge and Engizek Massifs, while marine sediments from Late Cretaceous to Miocene crop out in the North. There are 6 group or formation sedimentary units tcrop out at the North of the Pütürge and Engizek Massifs. They are: 1-Hazar group (Maastrichtian-Early Eocene), 2) Harami Formation (Late Maastrichtian), 3) Kuşçular Formation (Early Paleocene), 4) Seske Formation (Late Paleocene to Early Eocene), 5) Kırkgeçit Formation (middle Eocene-Oligocene) and 6) Plio-Quaternary sedimentary rocks [36, 50, 91, 94, 169-172].

### **2.5.1.Hazar Group**

The typical locality of the Hazar Group, which is named differently such as Hazar unit [145, 159], Hazar complex [137], Hazar formation [126], Hazar Group [9], is in the East of Hazar Lake (Elazığ). The group unconformably overlays the Late Cretaceous Guleman ophiolite. Coarse-grained ophiolite-derived conglomerates form the base of the Group. Shale and limestone alternations dominate towards the upper levels. The unit continues with shallow-marine mudstones and sandstones and ends up with neritic limestones at the top

According to the fossils determined within the group, the age of the group is Maastrichtian–Early Eocene [9, 137, 159]. The possible equivalents of the Hazar Group are reported to be Maastrichtian to middle Eocene in the Malatya and Palu [137, 186].

### **2.5.2.Harami Formation**

The unit, first described by Erdoğan [66] to the North of Gölbaşı (Adıyaman), crops out in limited areas around Gölbaşı and Elazığ. The Harami Formation crops out in the North (around Harput) and in the South of Elazig city center. The Harami Formation depositionally overlays the Elazığ magmatics. While the contact is angular unconform in some places, it is



vertical transitive in some places at least locally [24,7]. The ophiolites are depositionally overlain by the Harami Formation in the Gölbaşı area. The formation is overlain by the Paleocene Kuşçular Formation, middle-upper Eocene Kırkgeçit formation and the Quaternary Harput volcanics (7, 34, 36, 80, 91]. The Harami formation begins with Elazig magmatics-derived conglomerate level in some place, the base of the formation consists of sandstones and sandy limestones in some other places [5, 7, 24, 36]. The unit continues by shallow-marine limestone. The age of the unit is commonly accepted as (Late) Maastrichtian [36, 169, 173, 187]. The upper age range of the unit has been lately extended into the Late Paleocene by Herece and Acar [78]. Other areas where the Harami formation outcrops are the West of Elazig (Baskil) and the surroundings of Gölbaşı district of Adıyaman province. The Harami Formation which is represented by alternating pelagic limestone, shale, marl, radiolarite, manganiferous shale, and mudstone tectonically overlies the Esence Granitoid and Göksun ophiolite and is thrust by Malatya Metamorphics, in the Gölbaşı (Adıyaman)-Kahramanmaraş areas [191]. Elsewhere, in another area, the ophiolites are depositionally overlain by the Harami Formation (Yıldırım, 2015).

#### 2.5.3.Kuşçular Formation

The Kuşçular Formation crops out in the West of Elazığ, Baskil area. It rests unconformably on the Keban metamorphics, the Elazığ magmatics and the Harami limestones, and is unconformably overlain by a prominent carbonate unit, the Seske Formation. The Formation consists of conglomerates, sandstones, red mudstones, and gypsum levels. Its Early Palaeocene age is inferred from the bio-sstratigraphic ages of the underlying and overlying formations [94, 122, 171]. The Kusçular Formation was deposited in the Early Paleocene in a tectonically-controlled foreland basin [94].

#### 2.5.4.Seske Formation

The Seske formation crops out in narrow areas around Adıyaman and Elazığ [66, 78,128,170, 173]. The Seske Formation unconformably overlies the Elazığ magmatics and the Kuşçular Formation and is unconformably overlain by Kırkgeçit Formasyon. Although the Seske Formation is mostly represented by limestones around Elazig, it shows local lithological differences. The formation, which starts with massive limestones at the base in some areas, passes into bedded limestones and mudstones with red pelagic foraminifera towards the upper levels. In some places, it consists of massive limestones. The unit consists of shallow-marine limestone containing Late Paleocene-Early Eocene foraminiferal assemblages [173].

#### 2.5.5.Kırkgeçit Formation

Compared to other cover sedimentary units outcropping to the North of the Bitlis-Pütürge Massifs and ophiolites, the Kırkgeçit formation crops out in a wide area from Malatya to Van. The Kırkgeçit Formation rests unconformably on the Keban metamorphics, the Elazığ magmatics and the other (Late) Maastrichtian-Early Paleocene sedimentary units such as Harami, Kuşçular, Seske Formations. It is unconformably overlain by the Late Miocene–Early Pliocene Karabakır Formation and Quaternary volcanic (Harput volcanics) and sedimentary rocks. The Kırkgeçit Formation consists of a wide range of lithofacies and fossil assemblages [6, 19, 20, 36, 50, 78, 91, 94, 122, 128, 129, 173] . The facies characteristics indicate a deposition environment highly irregular basin floor topography and various depositional environments, from very shallow-marine to pelagic. The fossils it contains indicate that the age of the formation ranges from the middle Eocene to the Oligocene.

### 3. DISCUSSION

SEAOB was under the influence of intense tectonic events from Neoproterozoic to the Late Miocene. Therefore, primary relationships between units are not seen everywhere. There are many allochthonous units in the belt. The oldest unit in the belt is the lower-unit of the Bitlis-Pütürge metamorphics, and the crystallization ages of the augen gneisses and metagranites in the lower unit vary between 570-510 Ma. The oldest orogeny affecting the region is the Cadomian orogeny. U–Pb magmatic zircon ages indicate that Cadomian magmatism took place between 600 and 500 Ma and was especially intense during a 45-Myr timespan ca 570–525 Ma in Iran and Anatolia [33, 109, 177]. Researchers working in Turkey and Iran agree that the Late Proterozoic units cropping out in Iran and Turkey, including the Lower unit of the Bitlis-Pütürge massifs, are the remnants of an orogenic (Cadomian) belt along the northern margin of Gondwana. The geochemical, isotopic and geochronological features of augen gneiss and metagranites in the lower unit reveal the existence of Andean-type magmatic arcs and back-arc basins forming the northern margin of Gondwana, with southward subduction of Proto-Tethys oceanic lithosphere [33, 55, 72, 109, 111, 112, 114, 115, 143, 158, 178, 184] (Fig.11a). Avigad et al [18], who studies on the origin of the Mediterranean, suggest that the base of the Taurus Mountains chiefly consists of the graywacke succession formed in the mid- to Late Ediacaran back-arc basin over the southward subduction proto-Tethys Ocean. The arc and back-arc units were metamorphosed to various degrees and intruded by Ediacaran granites during Cadomian orogeny. The Ediacaran magmatism in the north of Gondwana is quite intense and there is a very prolonged

flare-up [108, 109]. Most of the Cambrian-Ordovician-Silurian lower Devonian rocks are absent. The first sedimentary unit overlying the metamorphic lower unit is the mid-Devonian aged transgressive sedimentary rocks. The deposition starting from the mid- Devonian continues under shelf environment conditions until the Late Triassic and abruptly change into deep-sea environment in the late Triassic. These sudden change conditions indicate a Late Triassic rifting between metamorphic Massifs and northern Gondwana. There is general acceptance that this rifting is onset of the opening of southern branch of Neo-Tethys. The Tarasa volcanics and Konak Formation of the Koçali complex which are the remains of an upper part of the oceanic crust, contain Late Triassic radiolarian fauna [180]. These radiolarian fauna indicating that the opening of the southern Neotethys Ocean began in the Triassic time. After rifting, spreading continued for about 140 million years (from Carnian? to Cenomanian?) between Gondwanaland and future Southeast Anatolian metamorphic massifs resulting in the creation of the southern branch of the Neotethys (Fig.11c). Karaoğlu et al. [87] and Robertson et al. [155] suggest that there were two different active oceanic realms within the southern branch of the Neotethys Ocean during the Late Cretaceous. One realm, called the Berit Ocean, was located between the Tauride platform to the North and the Bitlis–Pütürge microcontinent to the South; the other oceanic realm was situated between the Bitlis–Pütürge microcontinent to the North and the Arabian Platform to the South. Therefore, as we explained in the section ophiolites (section 2.2), the Malatya metamorphics are allochthonous on the Berit ophiolite and the Koçali ophiolite. In that case, these two ophiolites are products of the same oceanic crust.

The whole-rock geochemical, geochronological, and isotropic data described by different researchers strongly suggest that the Southeast Anatolian ophiolites were formed during the Late Cretaceous (92 to 82My) in a SSZ tectonic environment formed by the northward subducting of south branch of Neo-Tethys ocean crust [32, 43, 47, 56, 58, 59, 83, 84, 100, 101, 132, 133, 148, 149]. The recent geochronological data of ophiolites obtained from crustal rocks (92-82Ma) [43, 85,100] have revealed that the northwards subduction of southern Neo-Tethyan oceanic lithosphere started prior to 92 Ma. The continued northward subduction characterized by a moderate and constant dip of the subducted rock resulted in the formation of an intra-oceanic arc (Elazığ magmatics—volcanics, volcanoclastics, and granitoid) during the Late Cretaceous (84–72 Ma) (Fig. 11d and e).

The units on the subduction zone which are ophiolites, arc magmatics and southeastern Anatolian metamorphics migrated towards the South depending on the South-North compression at the end of the Cretaceous. The ophiolites thrust over the Arabian

platform together with the Karadut Complex. The massif forming the metamorphics was fragmented and migrates southward over ophiolites and arc magmatics and metamorphosed under greenschist facies conditions Fig. 11f). During all these tectonic events, the intrusive rocks of Elazig magmatics continue to form and cut the thrust zone between metamorphics and arc magmatics, metamorphics and ophiolites. After the southward thrust of the ophiolite and other units, the subsiding ocean is closed and the continental Arabian Plate enters the subduction zone. Despite the emplacement of the first ophiolitic nappes onto the Arabian continent during the Campanian– Early Maastrichtian period, the oceanic environment survived in the North of the Arabian platform [196, 198].

There is still much debate on when the northward subduction of Arabian Plate beneath Anatolia ceased and when the closure of the southern Neotethys and subsequent continental collision actually took place [9, 29, 32, 43, 45, 46, 73, 195].

There are three main alternative theories related to the time of the collision: 1) in the Late Cretaceous [29, 88, 95], 2) in the Late Eocene [11,182], or 3) during the Oligocene to Early Miocene [9, 151, 153, 154, 195]. In order to fully explain this issue, the units formed in the region after the Late Cretaceous thrust need to be examined in detail. The metamorphic massifs were fragmented and were thrust over the ophiolites by the Late Cretaceous thrust. The marine environment continues to the north and south of these massifs. Rising eustatic sea-level [105] possibly combined with isostatic regional subsidence following ophiolite emplacement resulted in a widespread marine transgression onto the Arabian Platform and the Koçali and the Kızıldağ ophiolites from Latest Cretaceous to Early Miocene times. The South of the Bitlis-Pütürge-Engizek massifs and Hatay areas, was dominated by shallow marine conditions in these periods. Maastrichtian–Early Eocene Hazar Group and Maastrichtian–Late Paleocene Harami Formation indicate the presence of the remnant of southern Neotethys Ocean realm over subduction zone to the North of the Bitlis-Pütürge metamorphics. The Hazar Group was deposited on the Guleman ophiolites in a shallow marine environment. The Harami Formation, on the other hand, was deposited in an east-west oriented basin over the Elazig magmatics. Both basins are not very large. Perinçek and Kozlu [140] suggest that the Harami Formation was deposited during periods when the island arc volcanism forming the Elazığ magmatics (Yüksekova Complex) was inactive. İnceöz [80] accepts that the formation was deposited after the Elazig magmatites completed their formation. Aksoy et al. [7] hypothesize that the Harami formation started in the shallow parts of the inner Tauride ocean during the late arc phases forming the Elazig magmatics. According to Herece et Acar [78], the Late Maastrichtian–Late Paleocene (Selandian) Harami Formation was deposited in

shallow shelf environment. All field data show that the formation of the Harami Formation started in the last phase of the intra-oceanic arc, and the basin floor is of very different depth and shape.

Until today, the most discussed issue in the region is the formation of the middle Eocene aged Maden Complex. The seven different models have been proposed for the formation of the this Complex. According to these models, the geotectonic environments in which the Maden complex is formed are:

- (1) a synorogenic “back-deep” type basin [145],
- (2) an immature island arc [64, 65],
- (3) an Eocene rifting zone [23],
- (4 ) a marginal basin formed behind the arc above a south-dipping subduction zone [141],
- (5) a back-arc basin [162],
- (6) a collisional belt [186, 187],
- (7) a lithospheric removal and asthenospheric upwelling associated with the extensional collapse of Southeastern Anatolian [68]

However, none of these models can fully explain the formation of the Maden Complex.


After the thrusting of the ophiolites on the the Arabian Platform, even though the subsiding ocean is closed, the subduction continued and due to the continental lithosphere and oceanic lithosphere exhibit different buoyancy, their contrasting buoyancy will finally lead in the breakoff of subducted slab. The subsiding ocean closure in Latest Maastrichtian and breakoff probably weakening the lithospheric mantle. This would have provided suitable conditions for subduction of the Arabian Plate because the breakoff of the subducted oceanic crust would have provided the pulling force for the subduction of the Arabian Plate Fig. 11g). The geochemical data suggest that the volcanic rocks of the Maden Complex are derived from a lithospheric mantle source. However, the positive and negative  $\epsilon\text{Nd}(t)$  values indicate the involvement of continental material [68]. The presence of the tourmaline-bearing leucogranites above the autochthonous Maden sediments has been interpreted by a large-scale intracrustal subduction [188]. Slab breakoff would open a slab window that allows the hot asthenosphere beneath the slab to rise into the mantle wedge, resulting in the intensification of magmatism. Magmatic rocks generated by such a process display compositional diversity with varying sources, such as the mantle wedge, the deep asthenosphere mantle, and the continental crust [183, 199]. The magma formed by the partial melting of the mantle wedge,

the rising deep asthenosphere mantle and the continental crust forms Maden arc over the ophiolites, Bitlis-Pütürge Massifs and the Hazar Group in the middle Eocene (Fig.11h). Above subduction zones, while Maden arc develops, behind it, shallow-deep marine carbonates and clastics (Kırkgeçit Formation) were deposited in a back-arc basin [8]. During the middle –Early Eocene and Early Oligocene, marine sediments were deposited in a large basin (Kırkgeçit Basin) to the North of the Bitlis– Pütürge Massifs [8, 79] (Fig.11i).


The last marine sedimentary rocks cropping out in the region belong to the early Miocene Alibonca Formation and are not very common. Closure of the sea in the Early Miocene was related to regional uplift following the closure of Neotethys and regional continent–continent collision in the middle Miocene, marking the beginning of the Neotectonic period [54, 76, 81, 160, 162].

The continuous northward migration of the Arabian Plate led to the disruption of the Tethys seaway and the final closure related to continental collision of Arabia and Eurasia. Figures (11 a-j) show the actual situation of the SEAOb. According to some researchers [11, 103], in its central segment of the SEAOb, the collision between Arabia and Eurasia started possibly from the Latest Eocene. After the Middle Eocene the large Kırkgeçit basin was closed. In early Miocene only a very shallow and narrowly distributed marine basin continues. Alibonca formation was deposited in this narrow basin. All these data indicate that the closure of the Southern Branch of the Neo-Tethys began in the late Eocene and was completed in the late Miocene. The continental collision of the Arabian Plate with the Eurasian Plate gave rise to the East Anatolian Accretionary Complex and the Caucasus-Iran-Anatolia (CIA) volcanic province [3, 99, 151, 162, 164, 175, 195]. This collision zone is associated with widespread “post-orogenic” [174] or “post-collisional” [48, 135, 161] volcanic eruptions. From this moment onward, the ongoing northward movement of the Arabian plate (still continuing today) [12, 102, 144], and the retreat of the Hellenic subduction zone to the west [25, 82, 97] led to westward tectonic escape of Anatolia along the North and East Anatolian Faults [53, 160, 163].


A) Neoproterozoic (Ediacaran)



B) Late Devonian-Early Triassic




C) Late Triassic-Early Cretaceous







#### D) Late Cretaceous (Turonian-Campanian)




#### E) Late Cretaceous (Campanian)




## F) Late Cretaceous (Maastrichtian)




## G) Late Cretaceous (Maastrichtian)




## H) Early Paleocene-Middle Eocene



# I) Middle Eocene-Early Miocene



## J) Actual (central part between Ergani (Diyarbakır)-Keban (Elazığ))



## K) Actual (Central part, between Çermik (Diyarbakır)-Keban (Elazığ))




Figure 11A-K- Tectonic evolution of the Souteast Anatolian orogenic Belt.

#### 4.CONCLUSIONS

Major advances are taking place in the study of geotectonics of the SEA OB from various aspects that include: 1) Detailed field studies of the internal structures of different units and their relations with each other, 2) the various geochemical, geochronological and isotopic evidences gathered by new methodology. A synthesis of these results lead to the following major conclusions:

-The oldest rocks of SEA OB is the Ediacaran Early Cambrian lower unit rocks of the Bitlis-Pütürge metamorphics. The geological, geochemical and geochronological data of augen gneiss, metagranites demonstrate the presence of a Cadomian active margin setting along the Northern Gondwana margin. Terranes within the Cadomian active margin were involved in the Cadomian orogeny from ~570 to ~520 Ma.

- The absence of Cambrian-Ordovician rocks onto the metamorphic basement indicates that the basement was exhumed and transgressed by sedimentary rocks during mid-Devonian.

- In the Late Triassic, the southern branch of the Neo-Tethys began to open with the rifting that started between the Arabian platform and the proto-Bitlis Pütürge massifs, and the oceanic expansion continued until the beginning of the Late Cretaceous.

- A northwards subduction of southern Neo-Tethyan oceanic lithosphere started prior to 92 Ma and as a result SE Anatolian ophiolites and arc magmatics were formed. At the end of the Late Cretaceous, the ophiolites were thrust over the Arabian platform and the metamorphic massifs were fragmented and drifted southward over the ophiolites and arc magmatics.

- After the thrusting of the ophiolites on the the Arabian Platform the subsiding ocean is closed and the continental Arabian plate enters the subduction zone. Even though the subsiding ocean is closed, the subduction continued and the breakoff of subducted slab has occurred.

- Following ophiolite emplacement resulted in a widespread marine transgression at the South of Bitlis-Pütürge Massifs onto the Arabian platform and the Koçali and the Kızıldağ ophiolites from Latest Cretaceous to Early Miocene times. In the same period, remnant basins of neotethys develop in the North of Bitlis-Pütürge Massifs.

-The magma formed by the partial melting of the mantle wedge, the rising deep asthenosphere mantle and the continental crust forms Maden arc over the ophiolites, Bitlis-

Pütürge Massifs and the Hazar Group in the middle Eocene. In the same periode, back-arc basin deposits in the north of the Maden arc are formed.

-The final closure, which started in the Late Eocene, ended in the Late Miocene. After the final closure, all units forming the SEA OB were thrust towards the south.

## 5. REFERENCES

1. Abbate E., Bortolotti V., Principi G., 1980. Apennine ophiolites: a peculiar oceanic crust. in: Rocci G., ed., Tethyan Ophiolites, 1, western area: Ofioliti special issue 59–96.
2. Açıkbash D., Baştuğ C., 1975. Geological report and oil possibilities of northern fields of V. Region Caca-Hani region. TPAO, No. 917.
3. Agard P., Omrani J., Jolivet L. Mouthereau F., 2005. Convergence history across Zagros (Iran): constraints from collisional and earlier deformation. *International Journal of Earth Sciences* 94, 401–419.
4. Akgül B., 1993. Petrographic and petrological properties of magmatic rocks around Piran Village. Unpubl. PhD Thesis, Firat University, Graduate School of Natural and Applied Sciences, Elazig, 128pp.
5. Aksoy E., 1993. General geological features of the west and south of Elazig. *Turkish Journal of Earth Sciences* (in Turkish) 1/1, 113-123.
6. Aksoy E., Turan M., Turkmen I., Özkul, M. 1995. Evolution of Elazig Basin in Tertiary. Black Sea Technical University 30th year Symposium abstracts, Trabzon, Turkey, p.45.
7. Aksoy E., Turkmen I., Turan M., Meriç E., 1999. New findings on stratigraphic position and depositional environment of Harami Formation (Upper Campanian-Maastrichtian), south of Elazig. *The Bulletin of Turkish Association of Petroleum Geologists* 11 (1), 1–15.
8. Aksoy E., Turkmen I., Turan M., 2005. Tectonics and sedimentation in convergent margin basins: an example from the Tertiary Elazig Basin, Eastern Turkey. *Journal of Asian Earth Sciences* 25, 459-472.
9. Aktaş G., Robertson A.H.F., 1984. The Maden Group, S.E. Turkey: Evolution of a Neotethyan active margin, in: the geological evolution of the eastern Mediterranean. Edited by J.E. Dixon and A.H.F. Robertson. Geological Society of London, Special Publication No. 17, 375–402.

10. Aktaş G., Robertson A.H.F., 1990. Tectonic evolution of the Tethys suture zone in SE Turkey: evidence from the petrology and geochemistry of Late Cretaceous and Middle Eocene Extrusives E.M. Moores, A. Panayiotou, C. Xenophontos, eds., *Ophiolites-Oceanic Crustal Analogues*. Proceedings of the International Symposium 'Troodos 1987', Geological Survey Department, Cyprus, 311-329.
11. Allen M.B., Armstrong H.A., 2008. Arabia–Eurasian collision and the forcing of mid Cenozoic global cooling. *Palaeogeography, Palaeoclimatology, Palaeoecology* 265 (1–2), 52–58. Doi:10.1016/j.palaeo.2008.04.021.
12. Allmendinger R. W., Reilinger R. Loveless J., 2007. Strain and rotation rate from GPS in Tibet, Anatolia and the Altiplano. *Tectonics*, 26, TC3013. doi: 10.1029/2006TC002030.
13. Al-Riyami K., Robertson A.H.F., Dixon J.E., Xenophontos C., 2002. Origin and emplacement of the Late Cretaceous Baer–Bassit ophiolite and its metamorphic sole in NW Syria. *Lithos* 65, 225–260.
14. Altınlı E., 1963. 1:500 000 scaled geological map of Turkey. (Cizre explanatory text of the geological map of Turkey): Mineral Research and Exploration Institute, Ankara, Turkey.
15. Arslan G., 2014. Mineralogical, petrographic and geochemical properties of plutonic rocks around Çolaklı (Elazığ). Unpubl. Master Thesis, Firat University, Graduate School of Natural and Applied Sciences, Elazığ, 56pp.
16. Arslan G., Beyarslan M., 2016. Mineralogical, petrographic and geochemical properties of plutonic rocks around Çolaklı (Elazığ). *C. U. Journal of the Faculty of Engineering and Architecture* 31 (1), 345–361.
17. Asutay H.J., 1988. Geology of Baskil (Elazığ) surroundings and petrology of Baskil magmatites, *Bulletin of Mineral Research and Exploration*, Ankara, Turkey 107, 46-72.
18. Avigad D., Abbo A., Gerdes A., 2016. Origin of the Eastern Mediterranean: Neotethys rifting along a cryptic Cadomian suture with Afro-Arabia. *Geo. Soc. am. bull.* <http://dx.doi.org/10.1130/B31370.1>.
19. Avşar N., 1983. Stratigraphic and micropaleontological studies in the northwest of Elazığ. Unpublish. PhD Thesis, Firat Univ. Graduate School of Natural and Applied Sciences, Elazığ, Turkey.

20. Avşar N., 1991, Presence of *Nummulites fabianii* (Prever) group (*Nummulites ex. gr. fabianii*) and associated foraminifers in the Elazığ region. Bulletin of the Mineral Research and Exploration, Ankara Turkey 112, 71-76.

21. Bağcı U., 2013. The geochemistry and petrology of the ophiolitic rocks from the Kahramanmaraş region, southern Turkey. Turkish. J. Earth Sci. 22, 1–27.

<https://doi.org/10.3906/yer-1203-1>.

22. Bağcı U., Parlak O., Höck V., 2005. Whole rock and mineral chemistry of cumulates from the Kızıldağ (Hatay) ophiolite (Turkey): clues for multiple magma generation during crustal accretion in the southern Neotethyan Ocean. Mineral Mag 69, 53–76.

23. Bastuğ M. C., 1980. Sedimentation, deformation and melange emplacement in the Lice basin, Dicle-Karabegan area, Southeast Turkey: Unpubl. PhD. thesis, Middle East Technical University, Ankara, Turkey.

24. Baykendi O. 1998. The geology of Tadım, Dedeyolu, Bademşınarı (Elazığ) villages around and petrographic properties of igneous rocks. Unpubl. Master Thesis, Firat University, Graduate School of Natural and Applied Sciences, Elazığ.

25. Berckhemer H., 1977. Some aspects of the evolution of marginal seas deduced from observations in the Aegean region. In: Montadert, L., ed., Structural History of the Mediterranean Basins. Technip. Paris, Split, Yugoslavia, 303–313.

26. Beyarslan M., 1991. Ispendere (Kale-malatya) Ofiyolitleri'nin Petrografik Özellikleri. Unpubl. Master Thesis, Firat University, Graduate School of Natural and Applied Sciences, Elazığ.

27. Beyarslan M., 1996. Kömürhan ophiolite biriminin petrografik ve petrolojik özellikleri. Unpubl. PhD, Thesis, Firat University, Graduate School of Natural and Applied Sciences, Elazığ.

28. Beyarslan M., 2017, Supra-subduction zone magmatism of the Koçali ophiolite, SE Turkey. J Afr. Earth Sci. 129, 390–402.

29. Beyarslan M., Bingöl A.F., 2000. Petrology of a supra-subduction zone ophiolite (Elazığ, Turkey). Can. J. Earth Sci. 37, 1411–1424.

30. Beyarslan M., Bingöl A.F., 2010. Ultramafics and mafic bodies in cumulates of Ispendere and Kömürhan Ophiolites (SE Anatolian Belt, Turkey) Turkish J. Sci. Technol. 5(1), 19–36.

31. Beyarslan M., Bingöl A.F., 2014. Petrology of the Ispendere, Kömürhan and Guleman Ophiolites (Southeast Turkey): Subduction Initiation Rule (SIR) Ophiolites and Arc Related Magmatic. 3rd Annual International Conference on Geological and Earth Sciences,



22-23 September 2014, Singapore, proceedings. pp. 50–59. <https://doi.org/10.5176/2251-3353-GEOS14.31>

32.Beyarslan M., Bingöl A.F., 2018. Zircon U-Pb age and geochemical constraints on the origin and tectonic implications of late cretaceous intra-oceanic arc magmatic in the Southeast Anatolian Orogenic Belt (SE-Turkey). *J. Afr. Earth Sci.* 147, 477–497.

33.Beyarslan M., Lin Y-C., Bingöl A.F., Chung S-L., 2016. Zircon U-Pb age and geochemical constraints on the origin and tectonic implication of Cadomian (Ediacaran-Early Cambrian) magmatism in SE Turkey. *J. Asian Earth Sci.* 130, 223–238.

34.Beyarslan M., Ertürk M.A., Rizeli M.E., Sar A., 2022. Doğu Anadolu Fay Sistemi boyunca gelişen Kuvaterner yaşlı mafik alkali Harput Volkanik Kayaçları'nın petrojenezi ve tektonik konumu, Güneydoğu Anadolu Orojenik Kuşağı (Elazığ). *El-Cezerî Journal of Science and Engineering* Vol: 9, No:1, 171-188. DOI:10.31202/ecjse.955277

35.Bingöl A.F. 1982. Elazığ – Pertek – Kovancılar arası volkanik kayaçların petrografik ve petrolojik incelenmesi. *F.Ü. Fen Fakültesi Dergisi*, 1, 9–21.

36.Bingöl A.F. 1984. Geology of Elazığ area in the Eastern Taurus region, in the geology of the Taurus Belt. Edited by O. Tekeli and M.C. Göncüoğlu. *International Symposium Proceedings, Maden Tetkik ve Arama Enstitüsü, Ankara, Turkey* 209–216.

37.Bingöl A.F., 1986. Petrographic and petrological characteristics of intrusive rocks of Guleman ophiolite (Eastern Taurus – Turkey). *Geosound* 13/14, 41–57.

38.Bingöl A.F., 1988. Petrographical and petrological features of the intrusive rocks of Yüksekova Complex in the Elazığ region (Eastern Taurus – Turkey). *Journal of Fırat University* 3/2, 1– 17.

39.Bingöl A.F., 1994. Çermik yöresinde (Güneydoğu Türkiye) Koçali Karmaşığına ait magma kayalarının jeokimyası ve petrolojisi. *TÜBİTAK Yerbilimleri Dergisi* 3, 55–61.

40.Bingöl A.F., Aydoğdu S., 1994. Dutluköy (Elazığ) yöresi magmatik kayaçlarının petrografik ve jeokimyasal özellikleri. In: *Çukurova Üniversitesi Müh. Mimarlık Fak. 15. Yıl Sempozyumu, Adana (Turkey)*, 199–214.

41.Bingöl A.F., Beyarslan M., 1996. Elazığ Mağmatitleri'nin jeokimyası ve petrolojisi. In: *Proceedings of Karadeniz Teknik Üniversitesi 30. yıl Sempozyumu*. Edited by S. Korkmaz and M. Akçay. Trabzon, Turkey, pp. 208–224.

42.Bingöl A.F., Beyarslan M., Akgül B., Erdem E., 1997. Karanlıkdere (Gölbaşı-Adıyaman) magmatitlerinin petrolojisi. In: *Proceedings of Selçuk Üniversitesi, Mühendislik-Mimarlık Fakültesi 20. Yıl Jeoloji Sempozyumu*. Edited by S. Temur. Konya, Turkey, 135–148.

43. Bingöl A.F., Beyarslan M., Lin Y-C., Lee H-Y., 2018. Geochronological and geochemical constraints on the origin of the Southeast Anatolian ophiolites, Turkey. *Arabian Journal of Geosciences* 11, 569. <https://doi.org/10.1007/s12517-018-3880-0>
44. Boulton S.J., 2009. Record of Cenozoic sedimentation from the Amanos Mountains, Southern Turkey: Implications for the inception and evolution of the Arabia–Eurasia continental collision. *Sedimentary Geology* 216, 29–47
45. Boulton S.J., Robertson A.H.F., 2007. The Miocene of the Hatay area, S Turkey: transition from the Arabian passive margin to an underfilled foreland basin related to closure of the Southern Neotethys Ocean. *Sedimentary Geology* 198, 93–124.
46. Cavazza W., Cattò S., Zattin M., Okay A.I., Reiners P., 2018. Thermochronology of the Miocene Arabia-Eurasia collision zone of southeastern Turkey. *Geosphere*; v. 14, no. 5. <https://doi.org/10.1130/GES01637.1>
47. Chen C., Su B-X., Uysal I., Avcı E., Zhang P-F., Xiao Y., He Y-S., 2015. Iron isotopic constraints on the origin of peridotite and chromitite in the Kızıldağ ophiolite, southern Turkey. *Chem. Geol.* 417, 115–124.
48. Chung S.L., Chu M.F., Zhang Y., Xie Y., Lo C.H., Lee T.Y., Lan C.Y., Li X., Zhang Q., Wang Y., 2005. Tibetan tectonic evolution inferred from spatial and temporal variations in post-collisional magmatism. *Earth Sci. Rev.* 68, 173–196.
49. Collins A.S., Pisarevsky S.A., 2005. Amalgamating eastern Gondwana: the evolution of the Circum-Indian Orogens. *Earth-Sci. Rev.* 71, 229–270.
50. Cronin B.T., Hartley A.J., Çelik H., Hurst A., Türkmen I., Kerey I.E., 2000. Equilibrium profile development in graded deep-water slopes: Eocene, Eastern Turkey. *Journal of the Geological Society of London*, 157, 943–955.
51. Çağlayan M. A., İnal R., Şengün M., Yurtsever A., 1984. Structural setting of Bitlis Massif. In: Tekeli, O., and Goncuoglu, M. C., eds., *Geology of the Taurus belt; International Symposium Proceedings*, Ankara, Turkey, Maden Tetkik ve Arama Enstitüsü, p. 245-254.
52. Coğulu E., Delaloye M., Vuagnat M., Wagner J.J., 1975. Some geochemical, geochronological and petrophysical data on the ophiolite massif from the Kızıldağ, Hatay, Turkey. *Comptes rendus des Séances de la Société de Physique et d'Histoire Naturelle de Genève* 10, 141–150.
53. Dewey J.F., Şengör A.M.C., 1979. Aegean and surrounding regions: complex multiplate and continuum tectonics in a convergent zone. *Geological Society of America Bulletin, Part I*, 84–92

- 54.Dewey J.F., Hempton M.R., Kidd W.S.F., Şaroğlu F., Şengör A.M.C., 1986. Shortening of continental lithosphere: The neotectonics of Eastern Anatolia—a young collision zone. In: Coward, M.P., and Ries, A.C., eds., *Collision Tectonics: Geological Society of London Special Publication* 19, 1–36. <https://doi.org/10.1144/GSL.SP.1986.019.01.01>
- 55.Dharma Rao C.V., Santosh M., Dong Y., 2012. U–Pb zircon chronology of the Pangidi–Kondapalle layered intrusion, Eastern Ghats Belt, India: constraints on Mesoproterozoic arc magmatism in a convergent margin setting. *J. Asian Earth Sci.*, 49, 362–375.
- 56.Dilek Y., Delaloye M., 1992. Structure of the Kızıldağ ophiolite, a slow-spread Cretaceous ridge segment North of the Arabian promontory. *Geology* 20, 19–22.
- 57.Dilek Y., Eddy C. A., 1992. The Troodos (Cyprus) and Kızıldağ (S. Turkey) ophiolites as structural models for slow-spreading ridge segments. *Journal of Geology* 100, 305–322.
- 58.Dilek Y., Thy P., 1998. Structure, petrology, and seafloor spreading tectonics of the Kızıldağ ophiolite, Turkey. In: Mills, R.A., Harrison, K., eds., *Modern ocean floor processes and the geological record. Geological Society of London Special Publication*, vol. 148, 43–69.
- 59.Dilek Y., Thy P., 2009. Island arc tholeiite to boninitic melt evolution of the cretaceous Kızıldağ (Turkey) ophiolite: model for multi-stage early arc-forearc magmatism in Tethyan subduction factories. *Lithos* 113, 68–87.
- 60.Dilek Y., Furnes H., Shallo M., 2007. Suprasubduction zone ophiolite formation along the periphery of Mesozoic Gondwana. *Gondwana Research* 11, 453–475.
- 61.Dubertret L., 1953. Géologie des roches vertes du nord-ouest de la Syrie et du Hatay (Turquie): Notes et Mémoires sur le Moyen-Orient 6, 227 pp.
- 62.Erdem E., 1994. Pütürge (Malatya) Metamorfikleri'nin petrografik ve petrolojik özellikleri. Unpubl. PhD Thesis, Fırat University, Graduate School of Natural and Applied Sciences Elazığ.
- 63.Erdem E., Bingöl A.F., 1995. Pütürge (Malatya) Metamorfiklerinin Petrografik Özellikleri. *F.Ü.Fen Müh. Bilim. Derg.* 7/1, 73–85.
- 64.Erdoğan B., 1977. Geology geochemistry and genesis of the sulfide deposits of the Ergani-Maden region SE Turkey. Ph.D. Thesis, University of New Brunswick, Canada.
- 65.Erdoğan B., 1982. Geology and volcanic rocks of the Southeast Anatolian ophiolite belt of the Ergani–Maden region. *Bulletin of the Geological Society of Turkey* 25, 49–59.
- 66.Erdoğan T., 1975. Gölbaşı yöresinin jeolojisi. TPAO Arama Grup Raporu no. 229.

- 67.Erendil M., 1984. Petrology and structure of the upper crustal units of the Kizildag ophiolite. In: Tekeli O., Göncüoğlu A.M., eds, Proceedings of the International Symposium on the Geology of the Taurus Belt. Turkey, Ankara, Turkey, Mineral Res Explor Institute, 269–284
- 68.Ertürk M.A., Beyarslan M., Chung S-L., Lin T-H., 2018. Eocene magmatism (Maden Complex) in the Southeast Anatolian Orogenic Belt: Magma genesis and tectonic implications. *Geoscience Frontiers* 9, 1829-1847.
- 69.Furnes H., De Wit M., Dilek Y., 2014. Four billion years of ophiolites reveal secular trends in oceanic crust formation. *Geosci Frontiers* 5, 571–603.
- 70.Garfunkel Z., Derin B., 1984. Permian Early Mesozoic tectonism and continental margin formation and its implications for the history of the Eastern Mediterranean. In: Dixon, J.E., and Robertson A.H.F., eds., *The Geological Evolution of the Eastern Mediterranean*. Geological Society, London, Special Publication, 17, 177-186.
- 71.Göncüoğlu M.C., Turhan N., 1984. Geology of the Bitlis Metamorphic Belt. In Tekeli, O., and Göncüoğlu, M.C., eds., *International Symposium on Geology of the Taurus Belt Proceedings*. Mineral Research and Exploration Institute of Turkey, Ankara, 237–244.
- 72.Gürsu S, Möller A., Göncüoğlu M.C., Köksal S., Demircan H., Köksal F.T., Kozlu H., Sunal G., 2015. Neoproterozoic continental arc volcanism at the northern edge of the Arabian Plate, SE Turkey. *Precambrian Res.* 258, 208–233.
- 73.Hall R., 1976. Ophiolite emplacement and evolution of the Taurus suture zone, southeast Turkey: *Geological Society of America Bulletin* 87, 1078–1088.
- 74.Hempton M.R., 1984. Results of detailed mapping near Lake Hazar (Eastern Taurus Mountains). In: the geology of the Taurus Belt. Edited by O. Tekeli and M.C. Göncüoğlu. *International Symposium Proceedings*. Maden Tetkik ve Arama Enstitüsü, Ankara, Turkey, 223–228.
- 75.Hempton R.M., 1987. Constraints on Arabian plate motion and extensional history of the Red Sea. *Tectonics* 6 (6), 687–705.
- 76.Hempton M.R., Savcı G., 1982. Elazığ Volkanik Karmaşığı'nın petrolojik ve yapısal özellikleri. *T.J.K. Bült.* 25, 143–150.
- 77.Helvacı C., Griffin W.L., 1983. Metamorphic feldspathization of metavolcanics and granitoids, Avnik Area, Turkey. *Contributions to Mineralogy and Petrology* 83, 309–319.
- 78.Herece E.I., Acar Ş., 2016. Upper Cretaceous-Tertiary geology/stratigraphy of Pertek and its vicinity (Tunceli, Turkey). *Bull. Min. Res. Exp.* 153, 1-44.

79.Hüsing S.K., Zachariasse W-J., van Hinsbergen, D.J.J., Krijgsman, W., Inceöz, M., Harzhauser, M., Mandic, O., Kroh, A., 2009. Oligocene-Miocene basin evolution in SE Anatolia, Turkey: constraints on the closure of the eastern Tethys gateway. Geological Society, London, Special Publications 311, 107-132 doi:10.1144/SP311.4

80.Inceöz M., 1994. Geological features of the northern and eastern vicinities of Harput (Elazığ). Unpubl. PhD Thesis, Firat University, Graduate School of Natural and Applied Sciences Elazığ, 112pp (in Turkish with English abstract).

81.Jackson J., McKenzie, D., 1984. Active tectonics of the Alpine-Himalayan Belt between W. Turkey and Pakistan. Geophysical Journal of the Royal Astronomical Society 77, 185–264.

82.Jolivet L., 2001. A comparison of geodetic and finite strain pattern in the Aegean, geodynamic implications. Earth and Planetary Science Letters, 187, 95–104.

83.Karaoğlu F., Parlak O., Klötzli U., Thöni M., Koller F., 2012. U–Pb and Sm–Nd geochronology of the ophiolites from the SE Turkey: implications for the Neotethyan evolution. Geodinamica Acta 25(3–4), 146–161.

84.Karaoğlu F., Parlak O., Klötzli U., Thöni M., Koller F., 2013a. U-Pb and SmNd geochronology of the Kızıldağ (Hatay, Turkey) ophiolite: implications for the timing and duration of suprasubduction zone type oceanic crust formation in southern Neotethys. Geol. Mag. 150, 283–299.

85.Karaoğlu F., Parlak O., Klötzli U., Koller F., Rızaoğlu T., 2013b. Age and duration of intra-oceanic arc volcanism built on a suprasubduction zone type oceanic crust in southern Neotethys, SE Anatolia. Geoscience Frontiers 4, 399–408.

86.Karaoğlu F., Parlak O., Robertson A.H.F., Thöni M., Klötzli U., Koller F., Okay A.I., 2013c. Evidence of Eocene high-temperature/high-pressure metamorphism of ophiolitic rocks and granitoid intrusion related to Neotethyan subduction processes (Doğanşehir area, SE Anatolia. In: Robertson, A.H.F., Parlak, O., and Ünlügenç, C., eds., Geological Development of Anatolia and the Easternmost Mediterranean Region. Geol. Soc. London Spec. Publ. 372, 249–272.

87.Karaoğlu F., Parlak O., Hejl E., Neubauer F., Urs K.S., 2016. The temporal evolution of the Active Margin along the Southeast Anatolian Orogenic Belt (SE Turkey): evidence from U-Pb, Ar-Ar and Fission Track Chronology. Gondwana Res. 33, 190–208.

<https://doi.org/10.1016/j.jgr.2015.12.011>.

- 88.Karig D. E., Kozlu H., 1990. Late Paleogene– Neogene evolution of the triple junction region near Maraş, south-central Turkey. *Journal of the Geological Society, London*, 147, 1023–1034.
- 89.Kaya A., 2004. Gezin (Maden-Elazığ) çevresinin jeolojisi. *Pamukkale Univ. Müh. Fak. Dergisi* 10/1, 41-50.
- 90.Kaya A., 2016. Tectono-stratigraphic reconstruction of the Keban metamorphites based on new fossil findings, eastern Turkey. *J Afr Earth Sci* 124, 245–257.
- 91.Kaygılı S., 2021. Reassessment of the age and depositional environment of the Kırkgeçit Formation based on larger benthic foraminifera, NW Elazığ, Eastern Turkey. *Turkish Journal of Earth Sciences*. 45, 580-600.
- 92.Ketin I., 1966. Tectonic units of Anatolia. General Directorate of Mineral Research and Exploration (MTA) Bulletin 66, 20–34 (in Turkish with English abstract).
- 93.Kipman E., 1981. Kebanın Jeolojisi ve Keban sariyacı. *İstanbul Üniversitesi Yerbilimleri*, 1-2, 75-81.
- 94.Koç Taşgın C., 2017. Soft-sediment deformation related to syntectonic intraformational unconformity in the early Palaeocene alluvial-fan deposits of Kuşçular Formation in the Elazığ sector of Tauride foreland, eastern Turkey. *Journal of African Earth Sciences* 134, 665-677.
- 95.Kozlu H., 1997. Tectono-stratigraphic units of the Neogene basins (İskenderun, Misis-Andirin) and their tectonic evolution in the eastern Mediterranean region. PhD Thesis, Natural Science Institute, Cukurova University, Turkey.
- 96.Kürüm S., Akgül B., Öztüfekçi Önal A., Boztuğ D., Harlavan Y., Ural M., 2011. An example for arc type granitoids along collisional zones the Pertek granitoid taurus orogenic belt Turkey. *Int. J. Geosci.* 2 (3), 214–226.
- 97.Le Pichon X., Angelier J., Sibuet J.-C., 1982. Plate boundaries and extensional tectonics. *Tectonophysics* 81, 239–256.
- 98.Lin Y.-C., Chung S.-L., Bingöl A.F., Beyarslan M., Lee H.-Y., Yang J.-H., 2015. Petrogenesis of late Cretaceous Elazig magmatic rocks from SE Turkey: New age and geochemical and Sr-Nd- Hf isotopic constraints. *Goldschmidt 16–21 August Prag, Abstracts*, 1869.
- 99.Lin Y-C., Chung S-L., Bingöl A.F., Yang L., Okrostsvaridze A., Pang K-W., Lee H-Y., Lin T-H., 2020. Diachronous initiation of post-collisional magmatism in the Arabia-Eurasia collision zone. *Lithos* 376–377 105394. <https://doi.org/10.1016/j.lithos.2020.105394>

100. Lin K-Y., Wang K-L., Chung S-L., Lizuka Y., Bingöl A.F., 2016. Suprasubduction zone characteristics of the Guleman Ophiolite, SE Turkey: evidence from peridotite geochemistry. Abstract. AGU fall meeting, San Fransisco, 12–16 December.
101. Lin K-Y., Wang K-Y., Chung S-L., Bingöl A.F., Iizuka Y., Lee H-Y., 2020. Tracking the magmatic response to subduction initiation in the forearc mantle wedge: Insights from peridotite geochemistry of the Guleman and Kızıldağ ophiolites, Southeastern Turkey. *Lithos* 105737. <https://doi.org/10.1016/j.lithos.2020.105737>
102. McClusky S., Balassanian S., Barka A., Demir A., Ergintav S., Georgiev I., 2000. Global Positioning System constraints on plate kinematics and dynamics in the eastern Mediterranean and Caucasus. *Journal Geophysical Research* 105, 5695–5719.
103. McQuarrie N., van Hinsbergen D.J., 2013. Retrodeforming the Arabia-Eurasia collision zone: age of collision versus magnitude of continental subduction. *Geology* 41, 315–318.
104. Michard A., Whitechurch H., Ricou L.E., Montigny R., Yazgan, E. 1984. Tauric subduction (Malatya–Elazığ provinces) and its bearing on the tectonics of the Tethyan realm in Turkey. In: the geological evolution of the eastern Mediterranean. Edited by J.E. Dixon and A.H.F. Robertson. Geological Society of London, Special Publication No. 17, 361–373.
105. Miller B.V., Samson S.D., d'Lemos R.S., 1999. Time span of plutonism, fabric development, and cooling in a Neoproterozoic magmatic arc segment: U–Pb age constraints from syn-tectonic plutons, Channel Islands, UK. *Tectonophysics* 312, 79–95.
106. MTA 2004. Geological Map of Turkey, scale 1/500.000. Mineral Research and Exploration, Ankara.
107. Moghadam H.S., Li X-H., Santos J.F., Stern R.J., Griffin W.L., Ghorbani G., Sarebani N., 2017. Neoproterozoic magmatic flare-up along the N. margin of Gondwana: the Taknar complex, NE Iran. *Earth Planet. Sci. Lett.* 474, 83–96.
108. Moghadam H.S., Stern R.J., 2011, Geodynamic evolution of Upper Cretaceous Zagros ophiolites: formation of oceanic lithosphere above a nascent subduction zone. *Geol. Mag.* 148 (5–6), 762–801
109. Moghadam H.S., Li Q., Griffin W., Ster, R., Santos J., Lucci F., Beyarslan M., Ghorbani G., Ravankhah A., Tilhac R., 2021. Prolonged magmatism and growth of the Iran-Anatolia Cadomian continental arc segment in Northern Gondwana. *Lithos* 105940. <https://doi.org/10.1016/j.lithos.2020.105940>
110. Murphy J.B., Eguiluz L., Zulauf G., 2002. Cadomian Orogens, peri-Gondwanan correlatives and Laurentia-Baltica connections. *Tectonophysics* 352, 1–9.



111. Murphy J.B., Fernández-Suárez J., Keppie J.D., Jeffries T.E., 2004. Contiguous rather than discrete Paleozoic histories for the Avalon and Meguma terranes based on detrital zircon data. *Bulletin of Geological Society of America* 32 (7), 585–588.
112. Nadimi A., 2007. Evolution of the Central Iranian Basement. *Gondwana Research*, 12, 324–333. <https://doi.org/10.1016/j.gr.2006.10.012>.
113. Nance R.D., Murphy J.B., Keppie J.D., 2002. A Cordilleran model for the evolution of Avalonia. *Tectonophysics* 352, 11–31.
114. Nance R.D., Gutierrez-Alonso G., Keppie J.D., Linnemann U., Murphy J.B., Quesada C., Strachan R.A., Woodcock N.H., 2010. Evolution of the Rheic ocean. *Gondwana Res.* 17, 194–222.
115. Neubauer F., 2002. Evolution of late Neoproterozoic to early Palaeozoic tectonic elements in Central and Southeast European Alpine mountain belts: review and synthesis. *Tectonophysics* 352, 87–103.
116. Nutman A.P., Mohajjel M., Bennett V.C., Fergusson C.L., 2014. Gondwanan Eoarchean-Neoproterozoic ancient crustal material in Iran and Turkey: zircon U-Pb-Hf isotopic evidence. *Can. J. Earth Sci.* 51, 272–285.
117. Okay A., 2008. Geology of Turkey: A synopsis. *Anschnitt* 21, 19–42.
118. Okay A.I., Arman M.B., Göncüoğlu M.C., 1985. Petrology and phase relations of the kyanite-eclogites from eastern Turkey. *Contributions to Mineralogy and Petrology* 91, 196–204.
119. Önal A., 1995, Polat-Begre (Doğanşehir, Malatya) yöresinde yüzeyleyen magmatik kayaçların petrografik ve petrolojik özellikleri. Unpubl. PhD Thesis, Firat University, Graduate School of Natural and Applied Sciences Elazığ, 159pp.
120. Önal A., Beyarslan M. 2001. Doğanşehir-Sürgü Kadili Civarındaki MaPetrographic Properties of Gematic Rocks. *THIS. Journal of the Faculty of Engineering and Architecture* c.16, 2: 67–79, Konya.
121. Önal A., Bingöl A.F., 2000, Geochemical Characterization and Petrogenesis of the Polat Granitoid in Eastern Taurus Belt, Turkey. *Journal of the Geological Society of India*, 56, 235–251
122. Ozcan E., Les, G.Y., Jovane L., Catanzariti R., Frontalini F., Coccioni R., Giorgioni M., Rodelli D., Rego E.S., Kaygılı S., Asgharian Rostami, M., 2019. Integrated biostratigraphy of the middle to upper Eocene Kırkeçit Formation (Baskil section, Elazığ, eastern Turkey): larger benthic foraminiferal perspective. *Mediterranean Geoscience Reviews* <https://doi.org/10.1007/s42990-019-00004-6>

123.Özçelik M., 1982. The petrology and geochemistry of volcanic rocks and associated sulphide deposits of the S.E. Anatolian ophiolite belt, near Malatya, Turkey. PhD thesis, University of Durham, 454 pp.

124.Özgül N., Turş A., Özyardımcı N., Şenol M., Bingöl I., Uysal S., 1982. Geology of Munzur Mountains. MTA report no: 6995. Ankara, Turkey (in Turkish).

125. Özkan Y. Z., Öztunalı Ö., 1984. Petrology of the magmatic rocks of Guleman ophiolite. In: Tekeli O., Göncüoğlu M.C., eds., Geology of the Taurus Belt. Proceeding of International Symposium. MTA, Ankara, 285-294.

126.Özkaya İ., 1975. Structural geology of the Sason region. Proceedings of the 50th Anniversary of the Republic, MTA Ankara 21-40.

127.Özkaya İ., 1978. Ergani Maden region stratigraphy, Türkiye Geo. Exchange rate. Bulletin 21/2, 120-139.

128.Özkul M., 1988 Sedimentological investigations on the Kırkgeçit Formation in the western Elazığ. Unpubl. PhD Thesis, Fırat University, Graduate School of Natural and Applied Sciences Elazığ, (in Turkish with English abstract).

129.Özkul M., Kerey I.E., 1996. Facies analysis on deep sea, shelf complex: Kırkgeçit Formation (Middle Eocene–Oligocene), Baskil-Elazığ. Turkish Journal of Earth Sciences 5, 57–70 (in Turkish with English abstract).

130.Parlak O., 2006. Geodynamic significance of granitoid magmatism in the southeast Anatolian orogen: geochemical and geochronological evidence from Göksun-Afşin (Kahramanmaraş-Turkey) region. International Journal of Earth Sciences 95, 609-627.

131.Parlak O., Höck,V., Kozlu H., Delaloye M., 2004. Oceanic crust generation in an island arc tectonic setting, SE Anatolian Orogenic Belt (Turkey). Geological Magazine 141, 583-603.

132.Parlak O., Rızaoğlu T., Bağcı U., Karaoğlu F., Höck V., 2009. Tectonic significance of the geochemistry and petrology of ophiolites in Southeast Anatolia, Turkey. Tectonophysics 473, 173–187.

133.Parlak O., Karaoğlu F., Rızaoğlu T., Nurlu N., Bağcı U., Höck V., Önal A., Kürüm S., Topak Y., 2012. Petrology of the Ispendere (Malatya) ophiolite from the Southeast Anatolia: implications for the late Mesozoic evolution of the southern Neotethyan Ocean. In: Robertson, A.H.F., Parlak, O., and Ünlügenç, U., eds., Geological development of the Anatolian continent and the easternmost Mediterranean region. Geol Soc. London Spec. Publ. 372, 219–247

134. Parlak O., Karaoğlu F., Rızaoğlu T., Klötzli U., Koller F., Billor Z., 2013. UPb and  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  geochronology of the ophiolites and granitoids from the Tauride belt: implications for the evolution of the Inner Tauride suture. *J. Geodyn.* 65, 22–37.
135. Pearce J.A., Lippard S.J., Roberts S., 1984. Characteristics and tectonic significance of supra-subduction zone ophiolites. In: Kokelaar, B.P., Howells, M.F., eds., *Marginal pressure geology*. Geological Society of London special publication, vol 16, 77–89.
136. Perinçek D., 1978. Geological investigation of the Çelikhan-Sincik-Koçali (Adıyaman province) area and investigation of oil possibilities. Doctoral Thesis (unpublished). I.U. Faculty of Science, 212pp.
137. Perinçek D., 1979. Geology and petroleum possibilities of Palu – Karabegan – Elazığ – Sivrice – Malatya area. TPAO Archive, Ankara, Report No. 1361, (unpublished).
138. Perinçek D., 1990. Stratigraphy of Hakkari Province and its surroundings. *TPJD Bulletin* 2/1, 21. (in Turkish).
139. Perinçek D., Çelikdemir M.E., 1979, Geology and petroleum possibilities of Palu-Karabegan- Elazığ- Sivrice-Malatya region (unpublished). TPAO Archive Report No: 1361.
140. Perinçek D., Kozlu H., 1984. Stratigraphy and structural relations of the units in the Afşin-Elbistan-Doğanşehir region (Eastern Taurus). In: Tekeli O., Göncüoğlu M.C., eds, *Geology of the Taurus Belt. Proceedings of International Symposium Proceedings on the Geology of the Taurus Belt Miner Res Explor Inst Turkey, Spec. Publ.* 181—198.
141. Perinçek D., Özkaya İ., 1981. Tectonic evolution of the northern margin of the Arabian Plate: H.Ü. *Geosciences B*, 91-100
142. Pişkin Ö., 1972. Etude mineralogique de la region situee a L'Est de Çelikhan ( Taurus Oriental, Adıyaman –Turkey). *Geo. Soc. Turkey Bull.* V.21/2, 107-111.
143. Ramezani J., Tucker R.D., 2003. The Saghand Region, Central Iran: U-Pb Geochronology, Petrogenesis and Implications for Gondwana Tectonics. *American Journal of Science* 303, 622-665. <http://dx.doi.org/10.2475/ajs.303.7.622>
144. Reilinger R., McClusky S., Vernant P., Lawrence S., Ergintav S., Çakmak, R., et al., 2006. GPS constraints on continental deformation in the Africa–Arabia–Eurasia continental collision zone and implications for the dynamics of plate interactions. *Journal of Geophysical Research*, 111, V05411, doi:10.1029/2005JB004051.
145. Rigo de Righi M., Cortesini A., 1964. Gravity tectonics in foothills structure belt of Southeast Turkey. *Amer. Petrol. Geol. Bull.* 48 (12), 1911–1937.

- 146.Rızaoğlu T., 200., Andean-type Active Margin Formation in the Eastern Taurides: Geochemical and Geochronological Evidence from the Baskil Granitoid (Elazığ, SE Turkey), *Tectonophysics*, 473, 188–207.
- 147.Rızaoğlu T., Parlak O., İşler F., 2005. Geochemistry and tectonic significance of Esence Granitoid (Göksun-Kahramanmaraş) SE Turkey. *Yerbilimleri* 26, 1–13.
- 148.Rızaoğlu T., Parlak O., Höck V., İşler F., 2006. Nature and significance of late cretaceous ophiolitic rocks and its relation to the Baskil granitoid in Elazığ region, SE Turkey. *Geol Soc. London Spec. Publ.* 260, 327– 350
- 149.Rizeli M.E., 2021. GD Anadolu Orojenik Kuşağı Ofiyolitlerinin Manto Peridotitleri Bileşimleri ve Petrolojisi: Ana Element, İz Element Jeokimyası, Mineral Kimyası ve Fe, Mg ve Os İzotopları. Unpubl. PhD Thesis, Firat University, Graduate School of Natural and Applied Sciences Elazığ, 205pp (in Turkish with English abstract).
- 150.Rizeli M.E., Beyarslan M., Wang K-L., Bingöl A.F., 2016. Mineral chemistry and petrology of mantle peridotites from the Guleman ophiolite (SE Anatolia, Turkey): evidence of a forearc setting. *J.Afr. Earth Sci.* 123, 392–402.
- 151.Robertson A. H. F., 2000. Mesozoic-Tertiary tectonic-sedimentary evolution of a south Tethyan oceanic basin and its margins in southern Turkey. In Bozkurt, E., Winchester, J. A. and Piper, J. D. A., eds., *Tectonics and Magmatism in Turkey and the Surrounding Area*. Geological Society London, Special Publication, 173, 97–138.
- 152.Robertson A. H. F., Dixon J. E., 1984. Introduction: aspects of the geological evolution of the Eastern Mediterranean. In: Dixon, J. E., and Robertson, A. H. F., eds., *The Geological Evolution of the Eastern Mediterranean*. Geological Society, London, Special Publications 17, 1–74.
- 153.Robertson A.H.F., Ünlügenç U.C., Inan N., Taşlı K., 2004. The Misis– Andırın complex: a mid-tertiary melange related to late-stage subduction of the southern Neotethys in S Turkey. *J. Asian Earth Sci.* 22, 413–453.
- 154.Robertson A.H.F., Ustaömer T., Parlak O., Ünlügenç U.C., Taşlı K., Inan N., 2006. The Berit transect of the Tauride thrust belt, S Turkey: Late Cretaceous–Early Cenozoic accretionary/collisional processes related to closure of the Southern Neotethys. *J. Asian Earth Sci.* 27, 108–145.
- 155.Robertson A.H.F., Parlak O., Rızaoğlu T., Ünlügenç U.C., Ustaömer T., Inan N., Taşlı K., 2007. Late cretaceous-middle tertiary tectonic evolution of the Tauride thrust belt and the evolution of S Neotethys: evidence from SE Anatolia (Elazığ-Maden region). In:

Deformation of the continental crust: the legacy of Mike Howard. Ries A.C., Butler R.W.H., Graham, R.H., eds., Geol. Soc. London Spec Publication 272, 231–270.

156.Sar A., Ertürk M. A., Rizeli M., 2019. Genesis of Late Cretaceous intra-oceanic arc intrusions in the Pertek area of Tunceli Province, eastern Turkey, and implications for the geodynamic evolution of the southern Neo-Tethys: Results of zircon U-Pb geochronology and geochemical and Sr-Nd isotopic analyses, *Lithos*, 105263. <https://doi.org/10.1016/j.lithos.2019.105263>.

157.Selçuk H., 1981. Étude géologique de la partie méridionale du Hatay (Turquie). Thèse de Doctorat. Université de Genève, Suisse (unpublished).

158.Stampfli G.M., Von Raumer J.F., Borel G.D., 2002. Paleozoic evolution of pre-Variscan terranes: from Gondwana to the Variscan collision. In: Martínez Catalán, J.R., Hatcher, R.D.J.R., Arenas, R., Díaz García, F. (Eds.), *Variscan–Appalachian dynamics: the building of the late Paleozoic basement: Geological Society of America Special Paper*, 364, pp. 263–280.

159.Sungurlu O., 1974. Geology and petroleum possibilities of the northern part of petroleum district VI: second petroleum Congress of Turkey Proceedings, Ankara.

160.Şengör A.M.C., 1980. Principles of the neotectonics of Turkey. Turkish Geological Society Publication, 40pp (in Turkish)

161.Şengör A.M.C., Kidd W.S.F., 1979. The post-collisional tectonics of the Turkish-Iranian Plateau and a comparison with Tibet. *Tectonophysics* 55, 361–376.

162.Şengör A.M.C., Yılmaz Y., 1981. Tethyan evolution of Turkey, A plate tectonic approach. *Tectonophysics* 75, 181- 241.

163.Şengör A.M.C., Görür N., Şaroğlu F., 1985. Strike-slip faulting and related basin formation in zones of tectonic escape: Turkey as a case study. In: Biddle, K.T., Christie-Blick, N., eds., *Strike-slip Deformation, Basin Formation, and Sedimentation. Society of Economic Paleontologists and Mineralogists Special Publication*, vol. 37, 227–264 (in honor of J.C. Crowell)

164.Şengör A.M.C., Özeren M.S., Keskin M., Sakıncı M., Özbakır A.D., Kayan I., 2008. Eastern Turkish high plateau as a small Turkic-type orogen: Implications for post-collisional crust-forming processes in Turkic-type orogens. *Earth-Science Reviews* 90 (1-2), 1-48.

165.Tarhan N., 1986a. Geology of Göksun-Afşin-Elbistan region. *Geological Engineering*, 19, 3-9.

- 166.Tarhan N., 1986b. Evolution and origin of granitoid magmas related to the closure of Neo-Tethys in eastern Taurus (Turkey). *Bull Min Res Expl, MTA Ankara*, 107, 84–100.
- 167.Tekeli O., Erendil M., 1984. Geology and petrology of the Kızıldağ Ophiolite (Hatay). In: Tekeli, O., Göncüoğlu, M.C., eds., *Int. Symposium on the Geology of the Taurus Belt*. MTA Publ, Ankara, pp. 127–147.
- 168.Tolun N., 1960. Stratigraphy and tectonics of southeastern Anatolia: *Revue de la Faculté des Sciences de l'Université d'Istanbul. Series B*, v. 25, p. 3-4.
- 169.Turan M., 1984. Stratigraphy and tectonics of the Baskil-Aydinlar (Elazig) region. Unpubl. PhD Thesis, Firat University, Graduate School of Natural and Applied Sciences Elazig, (in Turkish with English abstract).
- 170.Turan M., Bingöl A.F., 1991. Tectono-stratigraphic features of the region between Kovancılar and Baskil (Elazığ). *Ahmet Acar Geology Symposium Proceedings* 211–227.
- 171.Turan M., Turkmen I., 1996. Stratigraphy and sedimentological features of Kuşçular Formation (Lower Paleocene). *Turkish Journal of Earth Sciences* 5, 109–121 (in Turkish with English abstract).
- 172.Turan M., Aksoy E., Bingöl A.F., 1995. Features of the geodynamic evolution of the Eastern Taurus in the vicinity of Elazig. *Firat University. Journal of Science and Engineering Sciences*, 7(2), 177–199.
- 173.Türkmen I., Inceöz M., Aksoy E., Kaya M., 2001. New findings on Eocene stratigraphy and paleogeography of Elazığ area. *Bulletin of Earth Sciences Application and Research Center of Hacettepe University* 24, 81–95 (in Turkish with English abstract).
174. Turner S., Sandiford M., Foden J., 1992. Some geodynamic and compositional constraints on "postorogenic" magmatism. *Geology* 20, 931–934.
- 175.Tüysüz N., Erler A., 1995. Geology and geotectonic implications of Kazikkaya area, Kagizman-Kars (Turkey). In: Örcen, S., ed., *Geology of the Black Sea Region. Proceedings of the International symposium on the Geology of the Black sea Region*, 7–11 September 1995, Ankara, Turkey. General directorate of mineral research and exploration and chamber of geological engineering, Ankara, 76–81.
- 176.Ural M., Arslan M., Göncüoğlu U.K., Kürüm S., 2015. Late Cretaceous arc and back-arc formation within the southern Neotethys: whole-rock, trace element and Sr-Nd-Pb isotopic data from basaltic rocks of the Yüksekova Complex (Malatya-Elazığ, SE Turkey). *Ophiolite* 40 (1), 57–72.

- 177.Ustaömer P.A., Ustaömer T., Collins A.S., Robertson A.H.F., 2009. Cadomian (Ediacaran–Cambrian) arc magmatism in the Bitlis Massif, SE Turkey: magmatism along the developing northern margin of Gondwana. *Tectonophysics* 473, 99–112.
- 178.Ustaömer P.A., Ustaömer T., Gerdes A., Robertson A.H.F., Collins, A.S., 2012. Evidence of Precambrian sedimentation/magmatism and Cambrian metamorphism in the Bitlis Massif, SE Turkey utilizing whole-rock geochemistry and U-Pb LA-ICP -MS zircon dating. *Gondwana Res.* 21, 1001-1018.
- 179.Uysal I., Ersoy E.Y., Karsli O., Dilek Y., Sadiklar M.B., Ottely C.J., Tiepolo M., Meisel T., 2012. Coexistence of abyssal and ultra-depleted SSZ type mantle peridotites in a neo-Tethyan Ophiolite in SW Turkey: constraints from mineral composition, whole-rock geochemistry (major–trace–REE–PGE), and re–Os isotope systematics. *Lithos* 132–133, 50–69.
- 180.Uzunçimen S., Tekin U.K., Bedi Y., Perinçek D., Varol E., Soycan H., 2011. Discovery of the late Triassic (middle Carnian–Rhae- tian) radiolarians in the volcano-sedimentary sequences of the Koçali complex, SE Turkey: correlation with the other Tauride units. *J. Asian Earth Sci.* 40:180–200.
- 181.Varol E., Bedi Y., Tekin U.K., Uzunçimen S., 2011. Geochemical and petrological characteristics of late Triassic basic volcanic rocks from the Koçali complex, SE Turkey: implications for the Triassic evolution of southern Tethys. *Ophiolite* 36(1), 101–115.
182. Vincent S.J., Morton A.C., Carter A., Gibbs S., Barabadze T.G., 2007. Oligocene uplift of the Western Greater Caucasus: an effect of initial Arabia–Eurasia collision. *Terra Nova* 19, 160–166.
- 183.von Blanckenburg F., Davies J. H., 1995. Slab breakoff: A model for syncollisional magmatism and tectonics in the Alps. *Tectonics*, Volume14, Issue1 120-131.
- 184.von Raumer J. F., Stampfli G. M., Ricardo Arenas R., Sánchez Martíne S., 2002, Ediacaran to Cambrian oceanic rocks of the Gondwana margin and their tectonic interpretation. *int. J. Earth Sci.(Geol Rundsch)* 104:1107– 1121. DOI 10.1007/s00531-015-1142-x
- 185.Vuagnat M., Plural E., 1967. Quelques reflexions sur le massif basique–ultrabasique du Kızıldağ, Hatay, Turquie. *Comptes rendus de la Société de Physique et d'Histoire Naturelle de Genève* 2(3):210–216.
- 186.Yazgan E., 1983. A geotraverse between theArabian Platform and the Munzur Mountains. *Field Guidebook, Excursion 5, International Symposium on Geology of the Taurus Belt*, Ankara, 17 p.

187. Yazgan E., 1984. Geodynamic evolution of the eastern Taurus region. In: *Geology of the Taurus Belt*. Edited by O. Tekeli and M.C. Goncuoglu. Mineral Research and Exploration Institute, Ankara, International Symposium Proceedings, 199-208.
188. Yazgan E., Chessex R., 1991. Geology and tectonic evolution of the southeastern Taurides in the region of Malatya. *Bulletin of the Turkish Petroleum Geologists Association*, 3(1), 1–42.
189. Yazgan E., Michard A., Whitechurch H., Montigny R., 1983. Le Taurus de Malatya (Turquie orientale), element de la suture sud-tethysien. *Bulletin de la Société Géologique de France*, 25, 59-69.
190. Yıldırım E., 2010. Petrology of igneous rocks between Çelikhan and Sincik (Adıyaman). Unpubl. PhD Thesis, Firat University, Graduate School of Natural and Applied Sciences Elazığ. 244pp.
191. Yıldırım E., 2015. Geochemistry, petrography and tectonic significance of the ophiolitic rocks, felsic intrusions and Eocene volcanic rocks of an imbrication zone (Helete area, Southeast Turkey). *Journal of African Earth Sciences* 107, 89–107.
192. Yıldırım E., Bingöl A.F., 2012. Petrography and petrology of the Karadere Formation: Çelikhan-Sincik (Adıyaman). *Firat Univ. Journal of Engineering Sciences*, 24(2), 157-167.
193. Yıldırım E., Bingöl A.F., Yıldırım N., 2011. Are the Vein Rocks, Which Cuts the Pütürge Metamorphics at the Çelikhan-Sincik Region, Belong to the Maden Complex? Do the Vein Rocks Intersecting the Pütürge Metamorphites in the Çelikhan-Sincik Region belong to the Mineral Complex? 63th Geological Congress of Turkey (Abstract).
194. Yılmaz Y., 1985. Geology of the Cilo Ophiolite: An ancient ensimatic island arc fragment on the Arabian Platform, SE Turkey: *Ophiolite*, 10(2/3), 457–484.
195. Yılmaz Y., 1993. New evidence and model on the evolution of the southeast Anatolian Orogen. *Geological Society of America Bulletin*, 105: 251–271.
196. Yılmaz Y., 2018. Southeast Anatolian Orogenic Belt revisited (geology and evolution), *Can. J. Earth Sci.* 00:1–18 (0000) [dx.doi.org/10.1139/cjes-2018-0170](https://doi.org/10.1139/cjes-2018-0170)
197. Yılmaz Y., Yiğitbaş E., 1991. The different ophiolitic metamorphic assemblages of the SE Anatolia and their significance in the geological evolution of the region. *Proceedings of the 8th Petroleum Congress*, 128–140.
198. Yılmaz Y., Yiğitbaş E., Çemen I., 2022. Tectonics of the Southeast Anatolian Orogenic Belt. *The Earth and Space Science Open Archive*. <https://doi.org/10.1002/essoar.10510308.1>



199. Zhu D-C., Wang Q., Chung S-L., Cawood P.A., Zhao Z-D., 2019. Gangdese magmatism in southern Tibet and India–Asia convergence since 120 Ma. *Geo. Soc. London, Spec. Publ.* 483, 583–604.

UNDER PEER REVIEW