

Evaluation of herbicides for the management of complex weed flora in French bean (*Phaseolus vulgaris* L.) under mid hill conditions of Himachal Pradesh, India

ABSTRACT

A field experiment was conducted at Experimental Farm of the Department of Vegetable Science and Floriculture, CSK HPKV, Palampur during *Kharif*, 2022 to study the effect of herbicide combinations for the management of weeds in French bean (*Phaseolus vulgaris* L.). The experiment was laid out in Randomized Block Design with three replications and comprised of eleven weed control treatments viz., oxyfluorfen 150 gha⁻¹ (pre-emergence), pretilachlor 1000 gha⁻¹ (pre-emergence), imazethapyr 100 gha⁻¹ (pre-emergence), quizalofop-ethyl 100 gha⁻¹ (pre-emergence), oxyfluorfen 100 gha⁻¹ (pre-emergence) fb quizalofop-ethyl 70 gha⁻¹ (post-emergence), pretilachlor 700 gha⁻¹ (pre-emergence) fb quizalofop-ethyl 70 gha⁻¹ (post-emergence), imazethapyr 70 gha⁻¹ (pre-emergence) fb quizalofop-ethyl 70 gha⁻¹ (post-emergence), pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) fb quizalofop-ethyl 50 gha⁻¹ + chlorimuron 3 gha⁻¹ (post-emergence), pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) fb HW, 2 HWs (30 and 45 DAS) and weedy check. Results of the study revealed that hand weeding (twice) and pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) fb HW effectively reduced weed count and dry matter of the weed species. Highest weed control efficiency of 75.6 per cent was obtained from hand weeding (twice) treatment followed by 63.7 per cent with pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) fb HW. The treatment, hand weeding (twice) also produced significantly higher pod yield (16.14 t/ha), gross (₹ 2, 83,795) and net (₹ 2, 66,732) returns due to weed control. Amongst herbicidal treatments, pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) fb HW proved best as it resulted in significantly higher pod yield (14.24 t/ha), gross (₹ 2, 38,006) and net (₹ 2, 31,129) returns due to weed control. The treatment, Imazethapyr 100 gha⁻¹ gave the highest marginal benefit cost ratio (MBCR) of 178.9.

Keywords: French bean, Imazethapyr, Quizalofop-ethyl, Oxyfluorfen, Pretilachlor, Weed flora

INTRODUCTION:

French bean (*Phaseolus vulgaris* L.), being the monopoly of hill farmers (Tripathi et al., 1986 and Sood et al., 2003), is commercially cultivated in an area of about 228.0 thousand hectares with a production of 2257 thousand metric tons in the country. “In Himachal Pradesh, it is mainly cultivated as a market crop in mid and high hill areas, covering 3.82 thousand hectares with a production of about 50.87 thousand metric tons. Its cultivation has become more popular amongst growers on account of its off-seasonality, relative ease in cultivation and highest profit margins. Of the various reasons for its low productivity, weeds pose the serious threat on account of frequent irrigation and high fertility which provide conducive environment for their growth and development and consequently reduce yield by 20-60 per cent” (Anonymous, 2009). “Though, the weeds can be effectively managed with the application of pre-emergence herbicides at critical period of crop weed competition but the continuous application of pre-emergence herbicides in crops alters annual-perennial balance in favour of perennial weeds”. (Kumar et al. 2010) “Use of pre-emergence herbicides at low doses in conjunction with manual weeding 30-40 days after seeding is environmentally safe, socially acceptable and economically viable” (Kumar et al. 2011). “However, unavailability of labour at critical period of crop-weed competition and sometimes unfavourable field conditions do not permit manual weeding. In literature, sufficient information on pre-emergence herbicides to control weeds has been reported from various quarters but the information on post-emergence herbicides or their combinations is lacking. Many a times,

extension workers and farmers demand information on post-emergence herbicides or their combinations particularly when they fail to spray pre-emergence herbicides due to one or other reasons and paucity of labour for manual weeding. Hence, it becomes imperative to identify appropriate herbicide(s) and their combinations to manage the complex weed flora in French bean". [48]

"Weeds have been recognized as a serious problem ever since the man started domesticating/cultivating plants and since then the battle against weeds is ever ending and often the costliest agronomic input for successful crop production. Weeds are the most under estimated crop pests in agriculture and cause maximum reduction in crop yields than other pest and diseases. Of the total annual loss of agricultural produce from various pests in India, weeds roughly account for 37%, insects for 29%, diseases for 22% and other pests for 12%" (Yaduraju, 2006). "Weeds are the silent robbers of plant nutrients, moisture, sunlight and compete for space that would otherwise be available to main crop. Weeds also harbour pests and disease causing organisms, cause adverse allelopathic effects on crop and reduce the yield and quality of the produce". [48]

"Weeds cause approximately 9.5% yield loss of wheat globally"(Khan et al., 2022a). "Phalaris minor, Avenafatua, Chenopodium album, Lathyrusaphaca, Angalisarvensis, and Melilotusindica are the most common and troublesome ones" (Khan et al., 2023a, Khan et al., 2022a). "Since weeds possess competitive and deleterious effects on each growth phase of wheat, it is of prime importance to follow new systems for their management"(Khan et al., 2023b).

"The wide usage of herbicides increases the chances of weed resistance and farmer's dependence. In addition to the resistance, the hazard caused by herbicides and their persistent toxic effect on the quality of all life aspects after reaching the action site are other major issues related to the chemical control"(Javaid et al., 2022, Nadeem et al., 2022). "Despite those several herbicide side effects, its use is extremely important in augmenting crop productivity to face all the necessities of food security and sustainability of human populations" (Khan et al., 2023c).

"Manual method of weed control is labour intensive, cumbersome and time consuming whereas, the mechanical methods of weed control are reported to cause injury to the root system. Weeds can be effectively controlled by manual, mechanical and chemical methods. Unavailability of labour at critical period of crop-weed competition and sometimes unfavorable field conditions do not permit manual weeding. Under such situations, use of herbicides becomes imperative. However, use of herbicides at low doses in conjunction with manual weeding is environmentally safe, socially acceptable and economically viable" (Kumar et al. 2011). Herbicides like fluchloralin as pre-plant-incorporation (PPI) and pendimethalin as pre-emergence (PE) have been recommended for weed control but these are effective only during initial period of crop growth (up to 30 DAS). Thus, for the effective control of weeds throughout the crop season, use of post-emergence herbicides or their combinations is necessary. There is every possibility that use of single post-emergence herbicide coupled with pre-emergence herbicide may prove effective and raise the income of farmers.

MATERIALS AND METHODS

The field investigation was carried out at Experimental Farm of the Department of Vegetable Science and Floriculture, CSK HPKV, Palampur[32° 6' North latitude and 76° 3' East longitude and 1290 m above mean sea level] during *kharif*, 2022. The site is falling

under mid-hill zone of Himachal Pradesh. The soil of this zone is of podzolic type with pH range of 5.0-6.0. Soil of experimental field was silty clay loam in texture, acidic in reaction, medium in organic carbon (0.71%), medium in available nitrogen (407 kg/ha), phosphorus (17.2 kg/ha) and potassium (162 kg/ha). Eleven treatment combinations namely, oxyfluorfen 150 g/ha (pre-emergence), pretilachlor 1000 g/ha (pre-emergence), imazethapyr 100 g/ha (pre-emergence), quizalofop-ethyl 100 gha⁻¹ (pre-emergence), oxyfluorfen 100 gha⁻¹ (pre-emergence) fb quizalofop-ethyl 70 gha⁻¹ (post-emergence), pretilachlor 700 gha⁻¹ (pre-emergence) fb quizalofop-ethyl 70 gha⁻¹ (post-emergence), imazethapyr 70 gha⁻¹ (pre-emergence) fb quizalofop-ethyl 70 gha⁻¹ (post-emergence), pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) fb quizalofop-ethyl 50 gha⁻¹ + chlorimuron 3 gha⁻¹ (post-emergence), pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) fb HW, 2 HWs (30 and 45 DAS) and weedy check were evaluated in a randomized block design with three replications. "Herbicidal sprays as per treatments were applied immediately after sowing (pre-emergence) and 30 days after sowing (post-emergence) with the help of knapsack sprayer using flat fan nozzle in 750 liters of water per hectare. Weed count and weed dry weight were recorded at 40, 60 days after sowing (DAS) and at harvest. Growth, yield attributes and yield were recorded at different growth and harvest times. The data were subjected to statistical analysis as per Panse and Sukhatme (1984) and the treatments were compared at 5 per cent level of significance to interpret the differences. The weed count data were analyzed after subjecting the original data to square root transformation *i.e.* $\sqrt{(x + 0.5)}$ and the treatment effects were compared using transformed means". [48] Weed control efficiency of different treatments was calculated as per the following formula given by Mishra and Tosh (1979).

$$\text{Weed control efficiency (\%)} = \frac{\text{DWC} - \text{DWT}}{\text{DWC}} \times 100$$

Where,

DWC - weed dry weight (g/m²) in control plot, and

DWT - weed dry weight (g/m²) in treated plot

RESULTS AND DISCUSSION

The dominant weed flora of the experiment site was comprised of *Digitaria sanguinalis*, *Trifolium repens*, *Artemisia vulgaris*, *Cyperus rotundus* and *Alternanthera philoxeroides*. A similar type of weed flora in French bean has also been reported by Rana *et al.* (2008) under the mid-hill conditions of Himachal Pradesh.

Total weed count

Weed control treatments had significantly influenced the population of total weeds at all the stages of observation (Table 1). Significantly highest count of total weeds was observed in weedy check. All the weed control treatments significantly reduced the population of total weeds over weedy check at all the stages of the observation. Hand weeding (twice) had the lowest population of total weeds at all the stages of observation.

Application of pretilachlor 500 gha⁻¹+ imazethapyr 50 gha⁻¹(pre-emergence) *fb* HW resulted in significantly lower population of total weeds at all the stages of observation but it was at par with pretilachlor 1000 gha⁻¹(pre-emergence) at 80 DAS. This was due to effective control of the weeds with the spray of pretilachlor 500 gha⁻¹+ imazethapyr 50 gha⁻¹(pre-emergence) *fb* HW which reduced the species-wise weed population and ultimately resulted in lowest weed count. The treatment, oxyfluorfen 100 gha⁻¹(pre-emergence) *fb* quizalofop-ethyl 70 gha⁻¹(post-emergence) was found to be least effective against total weed population at all the stages of observation.

Table 1: Effect of weed control treatments on total weed count (No./m²) at different stages of observation in French bean

Treatment	Dose (g/ha)	40 DAS	60 DAS	At harvest
Oxyfluorfen	150	13.1 (170.7)	13.4 (180.0)	15.7 (245.3)
Pretilachlor	1000	11.4 (130.7)	11.8 (138.7)	14.8 (220.0)
Imazethapyr	100	14.1 (197.3)	14.1 (198.7)	17.0 (289.3)
Quizalofop-ethyl	100	14.7 (216.0)	15.5 (238.7)	17.8 (316.0)
Oxyfluorfen/ <i>fb</i> quizalofop-ethyl	100 <i>fb</i> 70	16.3 (265.3)	15.8 (249.3)	18.6 (344.0)
Pretilachlor/ <i>fb</i> quizalofop-ethyl	700 <i>fb</i> 70	15.3 (234.7)	14.1 (197.3)	16.9 (284.0)
Imazethapyr/ <i>fb</i> quizalofop-ethyl	70 <i>fb</i> 70	15.4 (236.0)	15.1 (226.7)	18.1 (326.7)
Pretilachlor+ imazethapyr/ <i>fb</i> quizalofop-ethyl+ chlorimuron ethyl	500 + 50 <i>fb</i> 50 + 3	15.1 (226.7)	14.1 (200.0)	17.2 (294.7)
Pretilachlor + imazethapyr <i>fb</i> HW	500 + 50	9.1 (82.7)	10.0 (100.0)	14.3 (204.0)
Hand weeding (twice)	-	1.2 (1.3)	7.2 (52.0)	12.0 (144.0)
Weedy check	-	19.3 (373.3)	21.1 (445.3)	24.0 (576.0)
SE (m)±		0.3	0.4	0.3
CD (P = 0.05)		1.0	1.3	1.0

Values given in the parentheses are the mean of original values, Data subjected to ($\sqrt{x + 0.5}$) square root transformation; DAS: days after sowing, PE: pre-emergence, PoE: post-emergence, HW: hand weeding and *fb*: followed by

Total weed dry weight

The effects of weed control treatments on total weed dry matter accumulation have been presented in Table 2. There was gradual increase in total weed dry matter accumulation from 40 DAS up to harvest (80 DAS) in all weed control treatments. Weed control treatments had significantly influenced the dry matter accumulation of total weeds at all the stages of observation. Significantly higher total weed dry matter accumulation was recorded in weedy check at all the stages of observation. All the weed control treatments showed significant reduction in total weed dry matter accumulation over weedy check at all the stages of observation.

Table 2: Effect of weed control treatments on dry weight of weeds (g/m²) and weed control efficiency at different stages of observation

Treatment	Dose	40	60	At harvest
-----------	------	----	----	------------

	(g/ha)	DAS	DAS	DAS
Oxyfluorfen	150	2.2 (4.7)	7.1 (49.3)	8.7 (74.7)
Pretilachlor	1000	1.8 (2.7)	5.8 (33.3)	7.5 (56.0)
Imazethapyr	100	2.1 (4.0)	5.8 (33.3)	7.8 (61.3)
Quizalofop-ethyl	100	1.9 (3.1)	5.7 (32.0)	7.4 (54.7)
Oxyfluorfen/ <i>fb</i> quizalofop-ethyl	100 <i>fb</i> 70	2.6 (6.1)	6.0 (36.0)	9.8 (96.0)
Pretilachlor/ <i>fb</i> quizalofop-ethyl	700 <i>fb</i> 70	2.4 (5.9)	7.9 (61.3)	8.3 (68.0)
Imazethapyr/ <i>fb</i> quizalofop-ethyl	70 <i>fb</i> 70	2.3 (5.1)	6.9 (46.7)	8.3 (69.3)
Pretilachlor+ imazethapyr/ <i>fb</i> quizalofop-ethyl+ chlorimuron ethyl	500 + 50 <i>fb</i> 50 + 3	2.1 (3.8)	5.7 (32.0)	8.3 (68.0)
Pretilachlor + imazethapyr/ <i>fb</i> HW	500 + 50	1.5 (1.8)	5.3 (28.0)	7.3 (53.3)
Hand weeding (twice)	-	0.7 (0.0)	4.0 (16.0)	5.6 (30.7)
Weedy check	-	3.1 (9.2)	9.3 (85.3)	10.3 (106.7)
SE (m)±		0.3	0.3	0.4
CD (P = 0.05)		0.8	0.7	1.1

Values given in the parentheses are the mean of original values, Data subjected to ($\sqrt{(x + 0.5)}$) square root transformation; DAS: days after sowing, PE: pre-emergence, PoE: post-emergence, HW: hand weeding and *fb*: followed by

Among the treatments, hand weeding (twice) had significantly lower total weed dry matter accumulation as compared to other treatments. Among various herbicidal treatments, pretilachlor 500 gha⁻¹+ imazethapyr 50 gha⁻¹ (pre-emergence) *fb* HW behaving statistically alike with quizalofop-ethyl 100 gha⁻¹ (post-emergence), pretilachlor 1000 gha⁻¹ (pre-emergence), imazethapyr 100 gha⁻¹(pre-emergence) and pretilachlor 500 gha⁻¹+ imazethapyr 50 gha⁻¹(pre-emergence) *fb*quizalofop-ethyl 50 gha⁻¹+ chlorimuron ethyl 3 gha⁻¹(post-emergence) resulted in lowest dry matter accumulation of total weeds at all the stages on account of effective control of the weeds with the application of pretilachlor 500 gha⁻¹+ imazethapyr 50 gha⁻¹ (pre-emergence) *fb* HW which significantly reduced the species-wise weed count and ultimately resulted in lowering the weed dry matter accumulation.

Weed control efficiency

The data on effect of different treatments on weed control efficiency have been presented in Table 3. Hand weeding (twice) resulted in highest weed control efficiency of 75.6 per cent.

Table 3: Effect of weed control treatments on dry weight of weeds (g/m²) and weed control efficiency at different stages of observation

Treatment	Dose (g/ha)	Weed control efficiency (%)
Oxyfluorfen	150	46.2

Pretilachlor	1000	59.3
Imazethapyr	100	54.7
Quizalofop-ethyl	100	52.8
Oxyfluorfen/ <i>fb</i> quizalofop-ethyl	100 <i>fb</i> 70	49.7
Pretilachlor/ <i>fb</i> quizalofop-ethyl	700 <i>fb</i> 70	52.3
Imazethapyr/ <i>fb</i> quizalofop-ethyl	70 <i>fb</i> 70	49.9
Pretilachlor+ imazethapyr/ <i>fb</i> quizalofop-ethyl+ chlorimuron ethyl	500 + 50 <i>fb</i> 50 + 3	56.8
Pretilachlor + imazethapyr/ <i>fb</i> HW	500 + 50	63.7
Hand weeding (twice)	-	75.6
Weedy check	-	-

Treatment	Dose (g/ha)	Dry matter accumulation (g/m ²)	Haulm yield (kg/m ²)
-----------	-------------	---	----------------------------------

However, amongst different herbicidal treatments, pretilachlor 500 g/ha + imazethapyr 50 gha⁻¹ (pre-emergence) *fb* HW had the highest weed control efficiency of 63.7 per cent which was followed by pretilachlor 1000 gha⁻¹ (pre-emergence) and pretilachlor 500 ggha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) *fb* quizalofop-ethyl 50 gha⁻¹ + chlorimuron ethyl 3 g/ha (post-emergence) which displayed 59.3 and 56.8 per cent weed control efficiencies, respectively. However, the lowest weed control efficiency of 46.2% was recorded in oxyfluorfen 150 gha⁻¹ (pre-emergence). The results are in confirmation with the findings of Kavadi et al. (2016), Singh et al. (2015), Gupta et al. (2015), Patel et al. (2016), Ram et al. (2012), Prachand et al. (2015), Bali et al. (2016), Rana et al. (2013), Devi et al. (2016) and Devaraju and Senthivel (2017) who observed the highest weed control efficiencies and lowest weed dry weight in their experimental studies.

Effect on crop

Dry matter accumulation (g/m²)

Dry matter accumulation of plants as influenced by different weed control treatments has been presented in Table 4. A critical analysis of the values in the table indicated that dry matter accumulation (g/m²) of the plants was drastically influenced by different weed control treatments. Hand weeding (twice) had significantly highest dry matter accumulation (g/m²) of plants but behaved statistically similar with the herbicidal treatment, pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) *fb* HW. Amongst weed control treatments, pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) *fb* HW had the highest dry matter accumulation (g/m²) when evaluated against other herbicidal treatments. The next best treatment was pretilachlor 1000 gha⁻¹ (pre-emergence), which was statistically different

Oxyfluorfen	150	30.7	3.4
Pretilachlor	1000	41.0	6.3
Imazethapyr	100	35.7	4.8
Quizalofop-ethyl	100	35.3	4.5
Oxyfluorfenfbquizalofop-ethyl	100 fb 70	33.5	3.8
Pretilachlorfbquizalofop-ethyl	700 fb 70	33.0	5.7
Imazethapyrfbquizalofop-ethyl	70 fb 70	32.3	4.8
Pretilachlor+ imazethapyrfbquizalofop-ethyl+ chlorimuron ethyl	500 + 50 fb 50 + 3	38.0	5.6
Pretilachlor + imazethapyrfb HW	500 + 50	46.0	6.8
Hand weeding (Twice)	-	48.0	7.5
Weedy check	-	22.0	2.2
SE (m)±		1.4	0.4
CD (P = 0.05)		4.2	1.1

Table 4: Effect of weed control treatments on dry mater accumulation (g/m^2) and haulm yield (kg/m^2) of plants

from pretilachlor 500 g/ha^{-1} + imazethapyr 50 g/ha (pre-emergence) fb HW and hand weeding (twice). The treatment, oxyfluorfen 150 g/ha (pre-emergence) had lowest dry matter accumulation (g/m^2) which was comparable with oxyfluorfen 100 g/ha^{-1} (pre-emergence) fb quizalofop-ethyl 70 g/ha^{-1} (post-emergence), pretilachlor 700 g/ha^{-1} (pre-emergence) fb quizalofop-ethyl 70 g/ha^{-1} (post-emergence) and imazethapyr 70 g/ha^{-1} (pre-emergence) fb quizalofop-ethyl 70 g/ha^{-1} (post-emergence). The results are in close conformity with the results obtained by Rana et al. (2004) in rajmash.

Haulm yield (kg/m^2)

The influence of different weed control treatments on haulm yield has been depicted in Table 4. It was reflected from the analytical values that haulm yield was significantly influenced by different weed control treatments. Though hand weeding (twice) had the highest haulm yield but it was statistically comparable with pretilachlor 500 g/ha^{-1} + imazethapyr 50 g/ha^{-1} (pre-emergence) fb HW, pretilachlor 700 g/ha^{-1} (pre-emergence) fb quizalofop-ethyl 70 g/ha (post-emergence) and pretilachlor 1000 g/ha^{-1} (pre-emergence). However, the lowest haulm yield of 3.4 kg/m^2 area was recorded in the treatment supplemented with the pre-emergence spray of oxyfluorfen 150 g/ha^{-1} . The results are in conformity with the findings of Rana et al. (2004) who observed the highest haulm yield in rajmash supplemented with the two hand weedings.

The effects of treatments on pod yield in tonnes per hectare of French bean have been presented in Table 5. A perusal of data revealed that pod yield (tonnes/ha) was significantly influenced by weed control treatments. Hand weeding (twice) had significantly highest pod yield when compared with other treatments. Among herbicidal treatments, pretilachlor 500 g/ha^{-1} + imazethapyr 50 g/ha^{-1} (pre-emergence) fb HW (post-emergence), though at par with hand weeding (twice) had the highest pod yield when compared with other

herbicidal treatments. The next best treatment was pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹(pre-emergence) fb quizalofop-ethyl 50 gha⁻¹+ chlorimuron ethyl 3 gha⁻¹(post-emergence). The treatment, oxyfluorfen 150 gha⁻¹(pre-emergence) produced the lowest pod yield as compared to other weed control treatments. This might be due to the effective weed control accomplished with the spray of pre and post-emergence herbicides coupled with hand weeding which produced more number of pods/plant with increased pod weight that ultimately resulted in enhanced pod yield. The results are in confirmation with the findings of earlier researchers (Singh et al. (2001), Rana et al. (2002b), Ram et al. (2012), Rana et al. (2013) in garden pea; Chaudhari et al. (2016) in green gram and Gupta et al. (2017) in black gram. Uninterrupted growth of weeds in the weedy check reduced French bean green pod yield by 66.5% as compared to the best treatment, hand weeding (twice). Green pod yield under the herbicidal treatments was 1.53 to 2.66 times higher than the weedy check treatment.

Economics

Gross returns

A perusal of data in Table 5 revealed that different weed control treatments increased the gross returns over weedy check. The treatments, handweeding (twice), pretilachlor 500 g/ha + imazethapyr 50 gha⁻¹(pre-emergence) fb HW, pretilachlor 1000 gha⁻¹(pre-emergence), pretilachlor 500 gha⁻¹+ imazethapyr 50 gha⁻¹(pre-emergence) fb quizalofop-ethyl 50 gha⁻¹+ chlorimuron ethyl 3 gha⁻¹ (post-emergence), pretilachlor 700 gha⁻¹(pre-emergence) fb quizalofop-ethyl 70 gha⁻¹(post-emergence) and imazethapyr 70 gha⁻¹(pre-emergence) fb quizalofop-ethyl 70 gha⁻¹(post-emergence) resulted in higher gross returns as compared to rest of the treatments. In general, the spray of herbicides coupled with hand weeding was better than the sole application of herbicides for effective weed management and obtaining higher gross returns (Kumar et al. 2017).

Gross returns due to weed control

Data on gross returns due to weed control of different weed control treatments have been presented in Table 5. The treatment, hand weeding (twice) had highest gross returns of Rs. 283795 per hectare due to weed control followed by pretilachlor 500 gha⁻¹+ imazethapyr 50 gha⁻¹(pre-emergence) fb HW (Rs. 238006 per hectare), pretilachlor 1000 gha⁻¹, pre-emergence (Rs. 175497 per hectare), pretilachlor 500 gha⁻¹+ imazethapyr 50 gha⁻¹(pre-emergence) fb quizalofop-ethyl 50 gha⁻¹+ chlorimuron ethyl 3 gha⁻¹, post-emergence (Rs. 174812 per hectare) and pretilachlor 700 gha⁻¹(pre-emergence) fb quizalofop-ethyl 70 gha⁻¹, post-emergence (Rs. 172257 per hectare). However, treatment oxyfluorfen 150 g/ha (pre-emergence) resulted in lowest gross returns of Rs. 76416 per hectare on account of the presence of complex weed flora which comprised of highest weed count and weed dry matter during the entire crop growth stages. This might have resulted in reduced cost of cultivation due to less cost incurred on spray of sole herbicide. The results are in accordance with findings of Shruti and Salankinkop (2015), Patel et al. (2017) and Kavadi et al. (2016).

Table 5: Effect of different weed control treatments on economics of French bean

Treatment	Dose (g/ha)	Pod yield (t/ha)	Cost of cultivation (INR/ha)	Gross returns (INR/ha)	Gross return due to weed control (INR/ha)	Cost of weed control (INR/ha)	Net return due to weed control (INR/ha)	MBCR
Oxyfluorfen	150	7.15	60699	215858	76416	1192	75224	63.1
Pretilachlor	1000	11.39	66913	314939	175497	1732	173765	100.3
Imazethapyr	100	10.31	65099	293374	153932	856	153076	178.9
Quizalofop-ethyl	100	10.68	66317	299634	160192	1528	158664	103.8
Oxyfluorfen <i>fb</i> quizalofop-ethyl	100 <i>fb</i> 70	9.59	65339	271836	132395	2187	130208	59.5
Pretilachlor <i>fb</i> quizalofop-ethyl	700 <i>fb</i> 70	11.37	67764	311699	172257	2587	169670	65.6
Imazethapyr <i>fb</i> quizalofop-ethyl	70 <i>fb</i> 70	11.24	67341	309916	170474	1974	168500	85.3
Pretilachlor + imazethapyr <i>fb</i> quizalofop-ethyl + chlorimuron ethyl	500 + 50 <i>fb</i> 50 + 3	11.51	67796	314253	174812	2410	172402	71.5
Pretilachlor + imazethapyr <i>fb</i> HW	500 + 50	14.24	76134	377447	238006	6876	231129	33.6
Hand weeding (twice)	-	16.14	89168	423237	283795	17063	266732	15.6
Weedy check	-	4.32	54748	139442	0	0	0	0
		9.5						
SE(m) ±		1.3						
CD (P = 0.05)		3.9						

Total cost of cultivation

Data on cost of cultivation of different weed control treatments have been presented in Table 5. The total cost of cultivation (Rs. 89168 per hectare) was recorded higher for hand weeding (twice) on account of the enhancement of wages of the labour deployed for carrying out weeding whereas, minimum cost of Rs. 54748 per hectare was observed under weedy check treatment.

Cost of weed control

Data on cost of weed control of different weed control treatments have been depicted in Table 5. Maximum cost of weed control (Rs. 17063 per hectare) was recorded under hand weeding (twice) treatment due to the deployment of more man power in performing manual weeding. The minimum cost of Rs. 856 per hectare was realized under the treatment, imazethapyr 100 gha⁻¹ (pre-emergence).

Net returns due to weed control

Net returns accrued under different weed control treatments followed almost the same trend as the gross returns. Net returns from hand weeding (twice) treatment were highest (Rs. 266732 per hectare) as compared to other weed control treatments on account of highest gross returns due to weed control (Table 5). This was followed by the combo mixture of pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) *fb* HW, pretilachlor 1000 gha⁻¹ (pre-emergence) and pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) *fb* quizalofop-ethyl 50 gha⁻¹ + chlorimuron ethyl 3 gha⁻¹ (post-emergence). Amongst herbicidal treatment, oxyfluorfen 150 gha⁻¹ (pre-emergence) produced the lowest net returns of Rs. 75224 per hectare due to higher weed count and weed dry matter. The results of the present investigation are in line with the findings of Tewari et al. (2003), Ram et al. (2012), Chaudhary et al. (2014) and Prachand et al. (2015) who obtained the highest net returns due to weed control from the herbicidal treatments coupled with hand weeding.

Marginal Benefit Cost Ratio (MBCR)

The data on marginal benefit cost ratio of different weed control treatments have been presented in Table 5. On account of lower cost of weed control and higher net returns due to weed control the treatment, imazethapyr 100 gha⁻¹ (pre-emergence), quizalofop-ethyl 100 g/ha (post-emergence) and pretilachlor 1000 gha⁻¹ (pre-emergence) had the highest marginal benefit cost ratio (MBCR) of 178.9, 103.8 and 100.3 respectively. Contrary to the higher cost of weed control the treatments, hand weeding (twice), pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) *fb* HW, pretilachlor 700 gha⁻¹ (pre-emergence) *fb* quizalofop-ethyl 70 gha⁻¹ (post-emergence), oxyfluorfen 150 gha⁻¹ (pre-emergence) and imazethapyr 70 gha⁻¹ (pre-emergence) *fb* quizalofop-ethyl 70 gha⁻¹ (post-emergence) had the lowest marginal benefit cost ratio (MBCR) of 15.6, 33.6, 59.5, 63.1 and 65.6 respectively, in comparison to other weed control treatments. Similar observations were also recorded by Goud and Dikey, (2016), Rana et al. (2004), Shekhar et al. (2021) in rajmash; Godara and Singh (2015) in cluster

bean; Prachand et al. (2015), Devi et al. (2016) in soybean; Tamang et al. (2015) in green gram; Ramesh and Radhika (2016), Devaraju and Santhivel (2017), Gupta et al. (2017) and Patel et al. (2017) in black gram.

Conclusion

The study indicated that weeds in French bean can be controlled effectively with combined application of pretilachlor 500 gha⁻¹ + imazethapyr 50 gha⁻¹ (pre-emergence) fb HW and hand weeding (twice) which further improved the dry matter accumulation, haulm yield and pod yield.

REFERENCES

1. Anonymous 2009. Pulses in India-An insight into the world's leading consumer of pulses. Annual Report, IIPR, Kanpur
2. Chaudhry S, Verma VK, Singh V, Pyare R and Singh AK. 2014. Studies on efficiency of herbicides against weeds of blackgram (*Vignamungo* L.). Advanced Research Journal of Crop Improvement **5**(1): 40-43
3. Kavadi NB, Patel CK, Patel AR and Thumber BR. 2016. Integrated weed management in blackgram. Indian Journal of Weed Science **48** (2): 222-224
4. Kumar A, Saxena A and Singh PK. 2014. Chemical and mechanical weed management for increased yield of French bean. Indian Journal of Weed Science **46** (4): 350-352
5. Kumar S, Angiras NN, Rana SS and Sharma P. 2010. Crop- weed competition studies in okra (*Abelmoschus esculentus* L. Moench) under mid-hill condition of Himachal Pradesh. Himachal Journal of Agricultural Research **36**: 13-19.
6. Kumar S, Angiras NN, Sharma P and Rana SS. 2011. Integrated weed management in okra (*Abelmoschus esculentus* L. Moench) under mid-hill conditions of Himachal Pradesh. Himachal Journal of Agricultural Research **37**: 10-16.
7. Panotra N and Kumar A. 2016. Weed management practices on winter French bean (*Phaseolus vulgaris* L.) under western Uttar Pradesh conditions. International Journal of Applied Science **4**(2): 275-283
8. Panse VG and Sukhatme PV. 1984. Statistical Methods for Agricultural Workers, ICAR, New Delhi. p. 359
9. Patel RI, Piyush KS, Patel PH and Patel VN. 2017. Effect of different crop management practices on growth and yield of fieldpea. The Bioscan **12**(1): 667-669
10. Prachand S, Kalhapure A and Kubde KJ. 2015. Weed management in soybean with pre and post-emergence herbicides. Indian Journal of Weed Science **47** (2): 163-165
11. Ram B, Punia SS, Meena DS and Tatarwal JP. 2012. Efficacy of post emergence herbicides on weed control and seed yield of rajmash (*Phaseolus vulgaris* L.). Journal of Food Legumes **25** (4): 306-309
12. Rana MC, Nag M, Rana SS and Sharma GD. 2013. Influence of post-emergence herbicides on weeds and productivity of garden pea (*Pisum sativum* L.) under mid hill conditions of Himachal Pradesh. Indian Journal of Agronomy **58** (2): 226-230

13. Rana SS, Sood P, Pathania P, Kumar S, Chadha S and Chaudhary DR. 2008. Crop-weed competition in french bean (*Phaseolus vulgaris* L.) under dry temperature conditions of North-western Himalaya. *Himachal Journal of Agricultural Research* **34** (2): 10-14
14. Rana SS. 2002. Evaluation of promising herbicide combinations for weed management in Rajmash (*Phaseolus vulgaris* L.) under dry temperate conditions of Himachal Pradesh. *Indian Journal of Weed Science* **34** (3 & 4): 227-230
15. Singh G, Kaur H, Aggarwal N and Sharma P. 2015. Effect of herbicides on weed growth and yield of green gram. *Indian Journal of Weed Science* **47** (1): 38-42
16. Sood S, Awasthi CP and Singh N. 2003. Biochemical evaluation of promising Rajmash (*Phaseolus vulgaris* L.) genotypes in Himachal Pradesh. *Himachal Journal of Agricultural Research* **29**(1&2): 65-69
17. Tripathi DP, Chandra S and Asthana AN. 1986. Technology for growing rajmash in plains. *Indian Farming* **36** (9): 12-15
18. Khan, B. A, Nadeem, M. A, NajeedAlawadi, H., Javaid, M. M, Mahmood, A., Qamar, R., Iqbal, M., Mumtaz, A., Maqbool, R., Oraby, H. & Elnaggar, N. (2023c). Synthesis, characterization, and evaluation of nanoparticles of clodinfopropargyl and fenoxaprop-*P*-ethyl on weed control, growth, and yield of wheat (*Triticumaestivum* L.). *Green Processing and Synthesis*, **12**(1), 20230105.
19. Khan, B. A., Nadeem, M. A., Iqbal, M., Yaqoob, N., Javaid, M. M., Maqbool, R., ...&Oraby, H. (2023b). Chitosan nanoparticles loaded with mesosulfuron methyl and mesosulfuron methyl+ florasulam+ MCPA isooctyl to manage weeds of wheat (*Triticumaestivum* L.). *Green Processing and Synthesis*, **12**(1), 20228152.
20. Khan, B. A., Nadeem, M. A., Javaid, M. M., Maqbool, R., Ikram, M., &Oraby, H. (2022b). Chemical synthesis, characterization, and dose optimization of chitosan-based nanoparticles of clodinfopropargyl and fenoxaprop-p-ethyl for management of *Phalaris minor* (little seed canary grass): First report. *Green Processing and Synthesis*, **11**(1), 1118-1127.
21. Khan, B.A. A, Nijabat., M. I. Khan., I. Khan., S. Hashim., M. A. Nadeem. & M. Ikram (2022a). Implications of Mulching on Weed Management in Crops and Vegetables. In: Akhtar, K., Arif, M., Riaz, M., Wang, H. (eds) *Mulching in Agroecosystems*. Springer, Singapore. P (199-213).
22. Khan, B. A., Nadeem, M. A., Nawaz, H., Amin, M. M., Abbasi, G. H., Nadeem, M., ...&Ayub, M. A. (2023a). Pesticides: impacts on agriculture productivity, environment, and management strategies. In *Emerging Contaminants and Plants: Interactions, Adaptations and Remediation Technologies* (pp. 109-134). Cham: Springer International Publishing.
23. Anonymous 2009. Pulses in India-An insight into the world's leading consumer of pulses. Annual Report, IIPR, Kanpur

24. Chaudhry S, Verma VK, Singh V, Pyare R and Singh AK. 2014. Studies on efficiency of herbicides against weeds of blackgram (*Vignamungo* L.). *Advanced Research Journal of Crop Improvement* **5**(1): 40-43
25. Javaid, M.M., A. Mahmood, D.S. Alshaya, M.D. AlKahtani, H. Waheed, A. Wasaya, S.A. Khan, M. Naqvi, M. Haider, M.A. Shahid, M.A. Nadeem, S. Azmat, B. A. Khan, R. M., Balal, k, A. Attia and S. Fiaz. 2022. Influence of environmental factors on seed germination and seedling characteristics of perennial ryegrass (*Lolium perenne* L.). *Scientific Reports*, **12**(1): 1-11.
26. Kavadi NB, Patel CK, Patel AR and Thumber BR. 2016. Integrated weed management in blackgram. *Indian Journal of Weed Science* **48** (2): 222-224
27. Khan, B. A, Nadeem, M. A, Najeeb Alawadi, H., Javaid, M. M, Mahmood, A., Qamar, R., Iqbal, M., Mumtaz, A., Maqbool, R., Oraby, H. and Elnaggar, N. (2023c). Synthesis, characterization, and evaluation of nanoparticles of clodinafopropargyl and fenoxaprop-P-ethyl on weed control, growth, and yield of wheat (*Triticum aestivum* L.). *Green Processing and Synthesis*, **12**(1): 20230105.
28. Khan, B. A, Nadeem, M. A., Iqbal, M., Yaqoob, N., Javaid, M. M., Maqbool, R., and Oraby, H. (2023b). Chitosan nanoparticles loaded with mesosulfuron methyl and mesosulfuron methyl+ florasulam+ MCPA isooctyl to manage weeds of wheat (*Triticum aestivum* L.). *Green Processing and Synthesis*, **12**(1): 20228152.
29. Khan, B. A, Nadeem, M. A., Javaid, M. M., Maqbool, R., Ikram, M., & Oraby, H. (2022b). Chemical synthesis, characterization, and dose optimization of chitosan-based nanoparticles of clodinafopropargyl and fenoxaprop-p-ethyl for management of *Phalaris minor* (little seed canary grass): First report. *Green Processing and Synthesis*, **11**(1): 1118-1127.
30. Khan, B. A, Nadeem, M. A., Nawaz, H., Amin, M. M., Abbasi, G. H., Nadeem, M., ... and Ayub, M. A. (2023a). Pesticides: Impacts on agriculture productivity, environment, and management strategies. In *Emerging Contaminants and Plants: Interactions, Adaptations and Remediation Technologies* (pp. 109-134). Cham: Springer International Publishing.
31. Khan, B.A. A, Nijabat., M. I. Khan., I. Khan., S. Hashim., M. A. Nadeem. & M. Ikram (2022a). Implications of mulching on weed management in crops and vegetables. In: Akhtar, K, Arif, M, Riaz, M., Wang, H. (eds) *Mulching in agroecosystems*. Springer, Singapore. P (199-213).
32. Kumar A, Saxena A and Singh PK. 2014. Chemical and mechanical weed management for increased yield of French bean. *Indian Journal of Weed Science* **46** (4): 350-352
33. Kumar S, Angiras NN, Rana SS and Sharma P. 2010. Crop- weed competition studies in okra (*Abelmoschus esculentus* L. Moench) under mid-hill condition of Himachal Pradesh. *Himachal Journal of Agricultural Research* **36**: 13-19.

34. Kumar S, Angiras NN, Sharma P and Rana SS. 2011. Integrated weed management in okra (*Abelmoschus esculentus* L. Moench) under mid-hill conditions of Himachal Pradesh. *Himachal Journal of Agricultural Research* **37**: 10-16.
35. Nadeem, M. A., Khan, B. A., Chadar, A. R., Maqbool, R., Raza, A., Javaid, M. M., ...& Irfan, M. (2022). Weed control and sustainable rice production through rice intensification system and conventional practices of weed competition periods and age of transplanted seedlings. *Semina: Ciências Agrárias*, 43(5), 2271-2292.
36. Panotra N and Kumar A. 2016. Weed management practices on winter French bean (*Phaseolus vulgaris* L.) under western Uttar Pradesh conditions. *International Journal of Applied Science* **4**(2): 275-283
37. Panse VG and Sukhatme PV. 1984. *Statistical Methods for Agricultural Workers*, ICAR, New Delhi. p. 359
38. Patel RI, Piyush KS, Patel PH and Patel VN. 2017. Effect of different crop management practices on growth and yield of field pea. *The Bioscan* **12**(1): 667-669
39. Prachand S, Kalhapure A and Kubde KJ. 2015. Weed management in soybean with preand post-emergence herbicides. *Indian Journal of Weed Science* **47** (2): 163-165
40. Ram B, Punia SS, Meena DS and Tatarwal JP. 2012. Efficacy of post emergence herbicides on weed control and seed yield of rajmash (*Phaseolus vulgaris* L.). *Journal of Food Legumes* **25** (4): 306-309
41. Rana MC, Nag M, Rana SS and Sharma GD. 2013. Influence of post-emergence herbicides on weeds and productivity of garden pea (*Pisum sativum* L.) under mid hill conditions of Himachal Pradesh. *Indian Journal of Agronomy* **58** (2): 226-230
42. Rana SS, Sood P, Pathania P, Kumar S, Chadha S and Chaudhary DR. 2008. Crop-weed competition in french bean (*Phaseolus vulgaris* L.) under dry temperature conditions of North-western Himalaya. *Himachal Journal of Agricultural Research* **34** (2): 10-14
43. Rana SS. 2002. Evaluation of promising herbicide combinations for weed management in Rajmash (*Phaseolus vulgaris* L.) under dry temperate conditions of Himachal Pradesh. *Indian Journal of Weed Science* **34** (3 & 4): 227-230
44. Singh G, Kaur H, Aggarwal N and Sharma P. 2015. Effect of herbicides on weed growth and yield of green gram. *Indian Journal of Weed Science* **47** (1): 38-42
45. Sood S, Awasthi CP and Singh N. 2003. Biochemical evaluation of promising Rajmash (*Phaseolus vulgaris* L.) genotypes in Himachal Pradesh. *Himachal Journal of Agricultural Research* **29**(1&2): 65-69
46. Tripathi DP, Chandra S and Asthana AN. 1986. Technology for growing rajmash in plains. *Indian Farming* **36** (9): 12-15
47. Yaduraju NT. 2006. Herbicide resistant crops in weed management. **In**: The Extended Summaries, Golden Jubilee National Symposium on Conservation Agriculture and Environment. October, 26-28, BHU, Banaras, p. 297-98.

48. SINGH R, CHAUDHARY D, RANA S. Evaluation of imazethapyr and its ready mix combinations for weed control in okra (*Abelmoschus esculentus* (L.) Moench). *Journal of Crop and Weed*. 2021;17(1):211-6.

UNDER PEER REVIEW