

# Effect of Prandtl Number on Deissler's Decay Law of MHD Turbulence at Four-point Correlations

## Abstract

Deissler's decay law plays a **great** significance in Homogeneous and MHD turbulence flow. Fluid Dynamics are an interesting part of research work which **focuses** on many branches of science, engineering science and also in meteorology. In turbulent flow, the fluid particles show **irregular** movement and unpredictable behavior. The effect of **the Prandtl** number on Deissler's energy decay law of MHD turbulence at 4-point correlations has been described.

Key word: Deissler's decay law, MHD turbulence, Fluid Dynamics, turbulent flow, kinematics viscosity, viscous shear stress, Spectral Equations

## Introduction

In fluid dynamics turbulent flow is a flow system characterized by **disorganization** and whose performance is actually irregular. In space and time, it shows small momentum circulation, high momentum convection and quick disparity of pressure and velocity. In this case, flow parameters are abruptly changed e.g., **kinematic** viscosity causes instability of the viscosity. The problem of turbulence is very difficult to solve for the case of nonlinearity. Turbulent flow problems are always treated statistically for its irregular conditions. Turbulent flow is always disorganized but not all disorganized flows are turbulent. In fact, turbulence is an inter-active movement of eddies of different sizes. As a consequence the velocity at any point varies both in magnitude and direction with respect to time. Such a diffused flow is characterized as turbulent flow. At Reynolds number 4,000, the nature of flow in **circular pipes** is always assumed to be turbulent.

For turbulent flow, a constant source of energy supply is required because turbulence dissipates rapidly as the kinetic energy is converted into internal energy by viscous shear stress. Turbulent fluctuations **indicate** the energy losses for the velocity and pressure distributions in turbulent flows. Reynolds, O. [11] had the first methodical investigation on turbulent **flow**. Reynolds [11], is one of the **renowned researchers** who studied turbulent flow.

In particular, a turbulent flow exhibits all of the features, e.g. disorganized, chaotic, irregular behavior. In brief, turbulent flow exhibits irregular temporal behavior at any selected spatial location. Throughout this work, decay of energy of Magneto-hydrodynamic Turbulent Flow for four-point correlations has been considered. Finally, the result has established how energy decays due to **the** effect of Prandtl Number. Kraichnan's [19] established logically different ideas from previous efforts for direct interaction approximation. Using Deissler's energy decay law Bkar, *pk et al.* [22] studied "the decay of energy of MHD turbulence for four-point correlation" and Bkar, *pk et al.* [23] **generalized** "it for dust particle system". He also obtained [24] "energy decay law for rotating dust **particles**". Bkar, *pk et al.* [25] also studied "Effects of first-order reactant on MHD turbulence at four-point correlation". Bkar, *pk et al.* [26] obtained "4-Point Correlations of Dusty Fluid MHD Turbulent Flow in a 1st order Chemical-Reaction". Further **Bkar, pk et al.** studied [27] "**energy decay of MHD turbulence in a rotating system for first order chemical reaction**". Taylor, 1921 [15] developed "the impression of the Lagrangian correlation coefficient". Taylor, G. I. [13, 14] and Von Karman, T. [17, 18] described "turbulence in terms of

collisions between discrete entities and then set up the thought of velocity correlation at two or more points". Taylor, G. I. derived the "energy spectrum" method to explain the probability density function for energy in the turbulent flow field. The study of turbulence had been generalized by Boussinesq [1] and Reynolds [11]. Reynolds, O. [11] first found the remarkable difference between laminar and turbulent flows. Based on the problems of practical importance Prandtl [10] established "mixing length" theory such as pipe flows over borders of exact shapes. In 1938 Taylor, G. I. [16] discussed the non-linearity of the dynamical equations and found the probability distribution of the difference between the velocity components at two points. Taylor, G. I. [13] defined correlation coefficients between the fluctuating quantities and established the design that the velocity of the fluid of turbulent motion is a random continuous function of position and time. Kolmogoroff's [6] contributed to understanding the physics of turbulence and gives interpolation and extrapolation of stationary sequences and in [7] he explained a refinement of previous hypotheses concerning the local structure of turbulence in a viscous incompressible fluid at high Reynolds number. Hopf, E. [4, 5] also constructed theory of the characteristic functional to turbulence and first order reactant in MHD turbulence before the final period. Some characteristics of turbulent motion are completed by Kampe de Fériet, J. [3]. Applying Fourier transformations they [4, 5, and 3] established the three dimensional energy spectrum functions. Monuvar Hossain, *et al.*, [28] obtained homogeneous fluid turbulence before the final period of decay for four-point correlation in a rotating system for first-order reactants. Azad *et al.* [29] obtained the effect of chemical reaction on statistical theory of dusty fluid MHD turbulent flow for certain variables at three-point distribution functions. Bkar, *pk et al.* [30] also obtained the effect of first order chemical reaction for Coriolis force and dust particles for small Reynolds number in the atmosphere over territory. Azad *et al.*, [31] established effect of chemical reaction on statistical theory of dusty fluid MHD turbulent flow for certain variables at three-point distribution functions. Shimin Yu *et al.*, [32] studied the effect of Prandtl number on mixed convective heat transfer from a porous cylinder in the steady flow regime. Using Deissler's decay law [20, 21] and Abdul Malek Ph.D Thesis [33] . Now we are going to study the effect of the Prandtl number on Deissler's decay law at four-point correlations. In this context, a few concepts and mathematical tools for the foundation of MHD turbulence have been discussed. This report shows some aspects of fluid dynamics that are relevant to the Deissler's energy decay law

#### Four-point Correlation and Spectral Equations

We take the momentum equation of MHD turbulence at the point  $p$  and the induction equation of magnetic field fluctuation four point correlation and equations at  $p'$ ,  $p''$  and  $p'''$  as

$$\frac{\partial u_l}{\partial t} + u_k \frac{\partial u_l}{\partial x_k} - h_k \frac{\partial h_l}{\partial x_k} = - \frac{\partial \omega}{\partial x_l} + \nu \frac{\partial^2 u_l}{\partial x_k \partial x_k} \quad (1)$$

$$\frac{\partial h'_i}{\partial t} + u'_k \frac{\partial h'_i}{\partial x'_k} - h'_k \frac{\partial u'_i}{\partial x'_k} = \frac{\nu}{p_M} \frac{\partial^2 h'_i}{\partial x'_k \partial x'_k} \quad (2)$$

$$\frac{\partial h''_j}{\partial t} + u'' \frac{\partial h''_j}{\partial x''_k} - h'' \frac{\partial u''_j}{\partial x''_k} = \frac{\nu}{p_M} \frac{\partial^2 h''_j}{\partial x''_k \partial x''_k} \quad (3)$$

$$\frac{\partial h_m'''}{\partial t} + u_k''' \frac{\partial h_m'''}{\partial x_k'''} - h_k''' \frac{\partial u_m'''}{\partial x_k'''} = \frac{\nu}{P_M} \frac{\partial^2 h_m'''}{\partial x_k'''} \quad (4)$$

where  $\omega = \frac{P}{\rho} + \frac{1}{2} \overline{h}^2$  is the total MHD pressure  $\rho(x,t)$  is the hydrodynamic pressure,  $\rho$  is the fluid density,

$P_M = \frac{\nu}{\lambda}$  is the Magnetic Prandtl number,  $\nu$  is the kinematics viscosity,  $\lambda$  is the magnetic diffusivity,  $h_i(x,t)$  is the

magnetic field fluctuation,  $u_k(x,t)$  is the turbulent velocity,  $t$  is the time,  $x_k$  is the space co-ordinate and repeated subscripts are summed from 1 to 3.

Multiplying [Equ. \(1\)](#) by  $h_i' h_j'' h_m'''$  (2) by  $u_l h_j'' h_m'''$  (3) by  $u_l h_i' h_m'''$  (4) by  $u_l h_i' h_j''$  and adding the four equations, we then taking the space or time averages and they are denoted by  $(\overline{\dots})$  or  $\langle \dots \rangle$ . We get

$$\begin{aligned} & \frac{\partial}{\partial t} (\overline{u_l h_i' h_j'' h_m'''}) + \frac{\partial}{\partial x_k} (\overline{u_l u_k h_i' h_j'' h_m'''}) - \frac{\partial}{\partial x_k} (\overline{h_k h_l h_i' h_j'' h_m'''}) + \\ & \frac{\partial}{\partial x_k'} (\overline{u_l u_k h_i' h_j'' h_m'''}) - \frac{\partial}{\partial x_k'} (\overline{u_l u_i' h_k' h_j'' h_m'''}) + \frac{\partial}{\partial x_k''} (\overline{u_l u_k'' h_i' h_j'' h_m'''}) - \\ & \frac{\partial}{\partial x_k''} (\overline{u_l u_j'' h_i' h_k'' h_m'''}) + \frac{\partial}{\partial x_k''} (\overline{u_l u_k'' h_i' h_j'' h_m'''}) - \frac{\partial}{\partial x_k''} (\overline{u_l u_j'' h_i' h_j'' h_m'''}) = \\ & - \frac{\partial}{\partial x_l} (\overline{w h_i' h_j'' h_m'''}) + \frac{\partial^2}{\partial x_k \partial x_k} (\overline{u_l h_i' h_j'' h_m'''}) + \frac{\nu}{P_M} \left[ \frac{\partial^2}{\partial x_k' \partial x_k'} (\overline{u_l h_i' h_j'' h_m'''}) + \right. \\ & \left. \frac{\partial^2}{\partial x_k'' \partial x_k''} (\overline{u_l h_i' h_j'' h_m'''}) + \frac{\partial^2}{\partial x_k'' \partial x_k''} (\overline{u_l h_i' h_j'' h_m'''}) \right] \quad (5) \end{aligned}$$

Using the transformations

$$\frac{\partial}{\partial x_k''} = \frac{\partial}{\partial r_k'}, \quad \frac{\partial}{\partial x_k'} = \frac{\partial}{\partial r_k'}, \quad \frac{\partial}{\partial x_k} = -\left( \frac{\partial}{\partial r_k'} + \frac{\partial}{\partial r_k''} + \frac{\partial}{\partial r_k'''} \right)$$

into [Equ. \(5\)](#) we get,

$$\begin{aligned} & \frac{\partial (\overline{u_l h_i' h_j'' h_m'''})}{\partial t} + (1 + P_M) \frac{\partial^2}{\partial r_k' \partial r_k'} (\overline{u_l h_i' h_j'' h_m'''}) + (1 + P_M) \frac{\partial^2}{\partial r_k'' \partial r_k''} (\overline{u_l h_i' h_j'' h_m'''}) + 2P_M \frac{\partial^2}{\partial r_k' \partial r_k''} (\overline{u_l h_i' h_j'' h_m'''}) \\ & + 2P_M \frac{\partial^2}{\partial r_k' \partial r_k''} (\overline{u_l h_i' h_j'' h_m'''}) + 2P_M \frac{\partial^2}{\partial r_k' \partial r_k''} (\overline{u_l h_i' h_j'' h_m'''}) = \frac{\partial}{\partial r_k} (\overline{u_l u_k h_i' h_j'' h_m'''}) + \frac{\partial}{\partial r_k'} (\overline{u_l u_k h_i' h_j'' h_m'''}) + \end{aligned}$$

$$\begin{aligned}
& \frac{\partial}{\partial r_k''} (\overline{u_l u_k h_i' h_j'' h_m''''}) - \frac{\partial}{\partial r_k} (\overline{h_l h_k h_i' h_j'' h_m''''}) - \frac{\partial}{\partial r_k'} (\overline{h_l h_k h_i' h_j'' h_m''''}) - \frac{\partial}{\partial r_k''} (\overline{h_l h_k h_i' h_j'' h_m''''}) - \frac{\partial}{\partial r_k} (\overline{u_l u_k' h_i' h_j'' h_m''''}) + \\
& \frac{\partial}{\partial r_k} (\overline{u_l u_k' h_i' h_j'' h_m''''}) - \frac{\partial}{\partial r_k'} (\overline{u_l u_k'' h_i' h_j'' h_m''''}) + \frac{\partial}{\partial r_k'} (\overline{u_l u_j'' h_i' h_k'' h_m''''}) - \frac{\partial}{\partial r_k''} (\overline{u_l u_k'' h_i' h_j'' h_m''''}) + \frac{\partial}{\partial r_k''} (\overline{u_l u_m'' h_i' h_j'' h_m''''}) + \\
& \frac{\partial}{\partial r_l} (\overline{w h_i' h_j'' h_m''''}) + \frac{\partial}{\partial r_l'} (\overline{w h_i' h_j'' h_m''''}) + \frac{\partial}{\partial r_l''} (\overline{w h_i' h_j'' h_m''''}) \quad (6)
\end{aligned}$$

In order to write the [Equ. \(6\)](#) to spectral form, we can define the following [nine-dimensional](#) Fourier transforms

$$\langle u_l h_i'(\hat{r}) h_j''(\hat{r}') h_m''''(\hat{r}'') \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \gamma_i'(\hat{k}) \gamma_j''(\hat{k}') \gamma_m''''(\hat{k}'') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}' + \hat{k}'' \cdot \hat{r}'')] d\hat{k} d\hat{k}' d\hat{k}'' \quad (7)$$

$$\langle u_l u_k' h_i'(\hat{r}) h_j''(\hat{r}') h_m''''(\hat{r}'') \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \phi_k'(\hat{k}) \gamma_i'(\hat{k}) \gamma_j''(\hat{k}') \gamma_m''''(\hat{k}'') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}' + \hat{k}'' \cdot \hat{r}'')] d\hat{k} d\hat{k}' d\hat{k}'' \quad (8)$$

$$\begin{aligned}
& \langle u_l u_i' h_i'(\hat{r}) h_j''(\hat{r}') h_m''''(\hat{r}'') \rangle \\
& = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \phi_i'(\hat{k}) \gamma_i'(\hat{k}) \gamma_j''(\hat{k}') \gamma_m''''(\hat{k}'') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}' + \hat{k}'' \cdot \hat{r}'')] d\hat{k} d\hat{k}' d\hat{k}'' \quad (9)
\end{aligned}$$

$$\begin{aligned}
& \langle u_l u_k' h_i'(\hat{r}) h_j''(\hat{r}') h_m''''(\hat{r}'') \rangle \\
& = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \phi_k''(\hat{k}') \gamma_i'(\hat{k}) \gamma_j''(\hat{k}') \gamma_m''''(\hat{k}'') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}' + \hat{k}'' \cdot \hat{r}'')] d\hat{k} d\hat{k}' d\hat{k}'' \quad (10)
\end{aligned}$$

$$\begin{aligned}
& \langle u_l u_j'' h_i'(\hat{r}) h_k''(\hat{r}') h_m''''(\hat{r}'') \rangle \\
& = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \phi_j''(\hat{k}') \gamma_i'(\hat{k}) \gamma_j''(\hat{k}') \gamma_m''''(\hat{k}'') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}' + \hat{k}'' \cdot \hat{r}'')] d\hat{k} d\hat{k}' d\hat{k}'' \quad (11)
\end{aligned}$$

$$\begin{aligned}
& \langle u_l u_k h_i'(\hat{r}) h_j''(\hat{r}') h_m''''(\hat{r}'') \rangle \\
& = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \phi_k \gamma_i'(\hat{k}) \gamma_j''(\hat{k}') \gamma_m''''(\hat{k}'') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}' + \hat{k}'' \cdot \hat{r}'')] d\hat{k} d\hat{k}' d\hat{k}'' \quad (12)
\end{aligned}$$

$$\begin{aligned}
& \langle u_l u_i' h_i'(\hat{r}) h_j''(\hat{r}') h_m''''(\hat{r}'') \rangle \\
& = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \phi_i'(\hat{k}') \gamma_i'(\hat{k}) \gamma_j''(\hat{k}') \gamma_m''''(\hat{k}'') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}' + \hat{k}'' \cdot \hat{r}'')] d\hat{k} d\hat{k}' d\hat{k}'' \quad (13)
\end{aligned}$$

$$\begin{aligned}
& \langle w h_i'(\hat{r}) h_j''(\hat{r}') h_m''''(\hat{r}'') \rangle \\
& = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \delta \gamma_i'(\hat{k}) \gamma_j''(\hat{k}') \gamma_m''''(\hat{k}'') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}' + \hat{k}'' \cdot \hat{r}'')] d\hat{k} d\hat{k}' d\hat{k}'' \quad (14)
\end{aligned}$$

Interchange of points  $p'$  and  $p''$ ,  $p'$  and  $p'''$  the subscripts i and k; i and j results in the relations

$$\overline{u_l u_k''' h_i' h_j'' h_m'''} = \overline{u_l u_k' h_i' h_j'' h_m'''}; \overline{u_l u_k''' h_i' h_j'' h_m'''} = \overline{u_l u_k' h_i' h_j'' h_m'''};$$

$$\overline{u_l u_m''' h_i' h_j'' h_m'''} = \overline{u_l u_i' h_i' h_k'' h_j'' h_m'''}; \overline{u_l u_j''' h_i' h_k'' h_m'''} = \overline{u_l u_i' h_i' h_k'' h_j'' h_m'''};$$

By use of these facts and Equ. (7) to (14), we can write Equ. (6) in the form

$$\begin{aligned} & \frac{\partial}{\partial t} \overline{(\phi_l \gamma_i' \gamma_j'' \gamma_m''')} + \frac{\nu}{p_M} [(1 + P_M) K^2 + (1 + p_M) K'^2 + (1 + p_M) K''^2 + 2p_M K K' + 2p_M K K'' + 2p_M K K'''] \\ & \overline{(\phi_l \gamma_i' \gamma_j'' \gamma_m''')} = i(K_k + K_k' + K_k'') \overline{(\phi_l \phi_k \gamma_i' \gamma_j'' \gamma_m''')} - i(K_k + K_k' + K_k'') \overline{(\gamma_l \gamma_k \gamma_i' \gamma_j'' \gamma_m''')} - \\ & \overline{(\phi_l \phi_k' \gamma_i' \gamma_j'' \gamma_m''')} + i(K_k + K_k' + K_k'') \overline{(\phi_l \phi_k' \gamma_i' \gamma_j'' \gamma_m''')} + i(K_k + K_k' + K_k'') \overline{(\delta \gamma_i' \gamma_j'' \gamma_m''')} \end{aligned} \quad (15)$$

The tensor Equ (15) can be converted to the scalar equation by contraction of the indices  $i$  and  $j$ ;  $\frac{\partial}{\partial t} \overline{(\phi_l \gamma_i' \gamma_i'' \gamma_m''')} +$

$$\frac{\nu}{p_M} [(1 + P_M) K^2 + (1 + p_M) K'^2 + (1 + p_M) K''^2 + 2p_M K K' + 2p_M K K'' + 2p_M K K''']$$

$$\begin{aligned} & \overline{(\phi_l \gamma_i' \gamma_i'' \gamma_m''')} = i(K_k + K_k' + K_k'') \overline{(\phi_l \phi_k \gamma_i' \gamma_i'' \gamma_m''')} - \\ & i(K_k + K_k' + K_k'') \overline{(\gamma_l \gamma_k \gamma_i' \gamma_i'' \gamma_m''')} - i(K_k + K_k' + K_k'') \overline{(\phi_l \phi_k' \gamma_i' \gamma_i'' \gamma_m''')} + \\ & i(K_k + K_k' + K_k'') \overline{(\phi_l \phi_k' \gamma_i' \gamma_i'' \gamma_m''')} + i(K_k + K_k' + K_k'') \overline{(\delta \gamma_i' \gamma_i'' \gamma_m''')} \end{aligned} \quad (16)$$

If we take the derivative with respect to  $x_l$  of the momentum Equ (1) at p, we have,

$$-\frac{\partial^2 w}{\partial x_l \partial x_l} = \frac{\partial^2}{\partial x_l \partial x_l} (u_l u_k - h_l h_k) \quad (17)$$

Multiplying Equ. (17) by  $h_i' h_j'' h_m'''$ , taking time averages and writing the equation in terms of the independent variables  $\vec{r}$ ,  $\vec{r}'$ ,  $\vec{r}''$  we have,

$$-\left[ \frac{\partial^2}{\partial r_l \partial r_l} + \frac{\partial^2}{\partial r_l' \partial r_l'} + \frac{\partial^2}{\partial r_l'' \partial r_l''} + 2 \frac{\partial^2}{\partial r_l \partial r_l'} + 2 \frac{\partial^2}{\partial r_l' \partial r_l''} + 2 \frac{\partial^2}{\partial r_l \partial r_l''} \right] \overline{(w h_i' h_j'' h_m''')} =$$

$$\left[ \frac{\partial^2}{\partial r_l \partial r_k} + \frac{\partial^2}{\partial r_l \partial r'_k} + \frac{\partial^2}{\partial r'_l \partial r_k} + \frac{\partial^2}{\partial r'_l \partial r'_k} + \frac{\partial^2}{\partial r'_l \partial r''_k} + \frac{\partial^2}{\partial r'_l \partial r''_k} + \frac{\partial^2}{\partial r'_l \partial r''_k} + \frac{\partial^2}{\partial r'_l \partial r''_k} \right] \left( \overline{u_l u_k h'_i h''_j h'''_m} - \overline{h_l h_k h'_i h''_j h'''_m} \right) \quad (18)$$

$$-(\overline{\delta \gamma'_i \gamma''_j \gamma'''_m}) = \frac{(K_l K_k + K_l K'_k + K_l K''_k + K'_l K_k + K'_l K'_k + K'_l K''_k + K''_l K_k + K''_l K'_k + K''_l K''_k)}{K_l K_l + K'_l K'_l + K''_l K''_l + 2K_l K'_l + 2K'_l K''_l + 2K_l K''_l}$$

$$(\overline{\phi_l \phi_k \gamma'_i \gamma''_j \gamma'''_m} - \overline{\gamma_l \gamma_k \gamma'_i \gamma''_j \gamma'''_m}) \quad (19)$$

Equation (19) can be used to eliminate  $(\overline{\delta \gamma'_i \gamma''_j \gamma'''_m})$  from Equ (16) if we take contraction.

### Three-point Correlation and Spectral Equations

The spectral equations corresponding to the three-point correlation equations by contraction of the indices  $i$  and  $j$  are

$$\frac{\partial}{\partial t} (\overline{\phi_l \beta'_i \beta''_i}) + \frac{\nu}{P_M} [(1 + P_M)(K^2 + K'^2) + 2P_M K K'] (\overline{\phi_l \beta'_i \beta''_i}) = i(K_k + K'_k) (\overline{\phi_l \phi_k \beta'_i \beta''_i}) - i(K_k + K'_k) (\overline{\beta_l \beta_k \beta'_i \beta''_i}) - i(K_k + K'_k) (\overline{\phi_l \phi'_k \beta'_i \beta''_i}) + i(K_k + K'_k) (\overline{\phi_l \phi'_i \beta'_k \beta''_k}) + i(k_l + k'_l) \gamma \beta'_i \beta''_i$$

and

$$-(\gamma \overline{\beta'_i \beta''_j}) = \frac{(K_l K_k + K'_l K_k + K_l k'_k + K'_l K'_k)}{(K_l^2 + K_l'^2 + 2K_l K'_l)} (\overline{\phi_l \phi_k \beta'_i \beta''_i} - \overline{\beta_l \beta_k \beta'_i \beta''_j})$$

Here the spectral tensors are defined by

$$\langle u_l h'_i(\hat{r}) h'_j(\hat{r}') \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \beta'_i(\hat{k}) \beta''_j(\hat{k}') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}')] d\hat{k} d\hat{k}'$$

$$\langle u_l u'_k(\hat{r}) h'_i(\hat{r}) h''_j(\hat{r}') \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \phi'_k(\hat{k}) \beta'_i(\hat{k}) \beta''_j(\hat{k}') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}')] d\hat{k} d\hat{k}'$$

$$\langle u_l u'_i(\hat{r}) h'_i(\hat{r}) h''_j(\hat{r}') \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_l \phi'_i(\hat{k}) \beta'_i(\hat{k}) \beta''_j(\hat{k}') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}')] d\hat{k} d\hat{k}'$$

$$\langle u_i h'_i(\hat{r}) h''_j(\hat{r}') \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_i \beta'_i(\hat{k}) \beta''_j(\hat{k}') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}')] d\hat{k} d\hat{k}'$$

$$\langle u_i h_k(\hat{r}) h'_i(\hat{r}) h''_j(\hat{r}') \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \phi_i \beta_k(\hat{k}) \beta'_i(\hat{k}) \beta''_j(\hat{k}') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}')] d\hat{k} d\hat{k}'$$

$$\langle w h'_i(\hat{r}) h''_j(\hat{r}') \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \langle \gamma \beta'_i(\hat{k}) \beta''_j(\hat{k}') \rangle \exp[i(\hat{k} \cdot \hat{r} + \hat{k}' \cdot \hat{r}')] d\hat{k} d\hat{k}'$$

### Solution Neglecting Quintuple Correlations

Neglecting all the terms on the right side of Equ. (16), the equation can be integrated between  $t_1$  and  $t$  to give

$$\langle \phi_i \gamma'_i \gamma''_j \gamma'''_m \rangle = \langle \phi_i \gamma'_i \gamma''_j \gamma'''_m \rangle_1 \exp \left[ \left\{ \frac{-\nu}{p_M} (1 + p_M) (k^2 + k'^2 + k''^2 + 2kk' + 2k'k'' + 2kk'') \right\} (t - t_1) \right] \quad (20)$$

where  $\langle \phi_i \gamma'_i \gamma''_j \gamma'''_m \rangle_1$  is the value of  $\langle \phi_i \gamma'_i \gamma''_j \gamma'''_m \rangle$  at  $t = t_1$  that is stationary value for small values of  $k$ ,  $k'$  and  $k''$  when the quintuple correlations are negligible.

$$\frac{\partial}{\partial t} \overline{(k_k \phi_i \beta'_i \beta''_i)} + \frac{\nu}{p_M} [(1 + p_M)(K^2 + K'^2) + 2p_M KK'] \overline{(k_k \phi_i \beta'_i \beta''_i)} =$$

$$[a]_1 \int_{-\infty}^{\infty} \exp \left[ -\frac{\nu}{p_M} (t - t_1) \{ (1 + p_M)(k^2 + k'^2 + k''^2) + 2p_M(kk' + k'k'' + k''k) \} \right] dk'' +$$

$$[b]_1 \int_{-\infty}^{\infty} \exp \left[ -\frac{\nu}{p_M} (t - t_1) \{ (1 + p_M)(k^2 + k'^2 + k''^2) + 2p_M kk' - 2p_M k''k \} \right] dk'' +$$

$$[c]_1 \int_{-\infty}^{\infty} \exp \left[ -\frac{\nu}{p_M} (t - t_1) \{ (1 + p_M)(k^2 + k'^2 + k''^2) + 2p_M kk' - 2p_M k''k \} \right] dk'' \quad (21)$$

At  $t_1$ ,  $\gamma^{(s)}$  have been assumed independent of; that assumption is not, made for other times. This is one of several assumptions made concerning the initial conditions, although continuity equation satisfied the conditions. The complete specification of initial turbulence is difficult; the assumptions for the initial conditions made herein are partially on the basis of simplicity. Substituting  $dk'' = dk''_1 dk''_2 dk''_3$  and integrating with respect to  $k''_1, k''_2, k''_3$  and we get,

$$\frac{\partial}{\partial t} \overline{(k_k \phi_l \beta'_i \beta''_i)} + \frac{\nu}{p_M} [(1 + p_M)(K^2 + K'^2) + 2p_M KK'] \overline{(k_k \phi_l \beta'_i \beta''_i)} =$$

$$\left( \frac{\pi p_M}{\nu(t-t_1)(1+p_M)} \right)^{\frac{3}{2}} [a]_1 \exp \left[ \frac{\nu(t-t_1)(1+p_M)}{p_M} \left\{ \frac{(1+2p_M)(k^2 + k'^2)}{(1+p_M)^2} + \frac{2p_M kk'}{(1+p_M)^2} \right\} \right] +$$

$$\left( \frac{\pi p_M}{\nu(t-t_1)(1+p_M)} \right)^{\frac{3}{2}} [b]_1 \exp \left[ \frac{\nu(t-t_1)(1+p_M)}{p_M} \left\{ \frac{(1+2p_M)(k^2)}{(1+p_M)^2} + \frac{2p_M kk'}{(1+p_M)} + k'^2 \right\} \right]$$

$$+ \left( \frac{\pi p_M}{\nu(t-t_1)(1+p_M)} \right)^{\frac{3}{2}} [c]_1 \exp \left[ -\frac{\nu(t-t_1)(1+p_M)}{p_M} \left\{ k^2 + \frac{(1+2p_M)(k'^2)}{(1+p_M)^2} + \frac{2p_M kk'}{(1+p_M)} \right\} \right] \quad (22)$$

Which result in

$$\frac{\partial H}{\partial t} + \frac{2\nu k^2}{p_M} H = G$$

where

$$G = k^2 \int_{-\infty}^{\infty} 2\pi i \left[ \left\langle k_k \phi_l \beta'_i \beta''_i(\hat{k}, \hat{k}') \right\rangle - \left\langle k_k \phi_l \beta'_i \beta''_i(-\hat{k}, -\hat{k}') \right\rangle \right]_0.$$

$$\exp \left[ -\frac{\nu}{p_M} (t-t_0) \{ (1+p_M)(k^2 + k'^2) + 2p_M kk' \} \right] dk' + k^2 \int_{-\infty}^{\infty} \frac{2p_M \pi^{\frac{5}{2}}}{\nu} i \left[ b(\hat{k}, \hat{k}') - b(-\hat{k}, -\hat{k}') \right] .$$

$$\{ -\omega^{-1} \exp \left[ \left( -\omega^2 \right) \left\{ \frac{(1+2p_M)(k^2)}{(1+p_M)^2} + \frac{2p_M kk'}{(1+p_M)} + k'^2 \right\} \right] \right]$$

$$+ k \cdot \exp \left[ (-\omega^2) \left\{ (1+p_M)(k^2 + k'^2) + 2p_M kk' \right\} \right] - 1 \int_0^{\frac{\omega k}{2}} \exp(x^2) dx \} dk' +$$

$$k^2 \int_{-\infty}^{\infty} \frac{2p_M \pi^{\frac{5}{2}}}{\nu} i \left[ c(\hat{k}, \hat{k}') - c(-\hat{k}, -\hat{k}') \right] .$$



$$\left\{ \omega^{-1} \exp \left[ (-\omega^2) \left\{ k^2 + \frac{(1+2p_M)(k'^2)}{(1+p_M)^2} + \frac{2p_M kk'}{(1+p_M)} \right\} \right] + \right. \\ \left. k' \exp \left[ -\omega^2 \left( (1+p_M)(k^2 + k'^2) + 2p_M kk' \right) \right] \cdot \int_0^{\frac{\omega k'}{2}} \exp(x^2) dx \right\} dk' \quad (23)$$

Here  $H$  is the magnetic energy spectrum function, which represents contributions from various wave numbers (or eddy sizes) to the energy and  $G$  is the energy transfer function, which is responsible for the transfer of energy between wave numbers, [Equ \(23\)](#) which depends on the initial conditions.

$$(2\pi)^2 \left[ \left\langle k_k \phi_l \beta'_i \beta''_i(\hat{k}, \hat{k}') \right\rangle - \left\langle k_k \phi_l \beta'_i \beta''_i(-\hat{k}, -\hat{k}') \right\rangle \right]_0 = -\xi_0 (k^2 k'^4 - k^4 k'^2) \quad (24)$$

where  $\xi_0$  is a constant depending on the initial conditions. For the other bracketed quantities above equation is

$$\frac{4p_M \cdot \pi^{\frac{7}{2}}}{\nu} i \left[ b(\hat{k} \cdot \hat{k}') - b(-\hat{k} \cdot -\hat{k}') \right]_{\parallel} = \frac{4p_M \cdot \pi^{\frac{7}{2}}}{\nu} i \left[ c(\hat{k} \cdot \hat{k}') - c(-\hat{k} \cdot -\hat{k}') \right]_{\parallel} \quad (25) \\ = -2\xi_1 (k^4 k'^6 - k^6 k'^4)$$

Remembering that  $d\hat{k}' = -2\pi \cdot \hat{k}'^2 d(\cos \theta)$  and  $kk' = kk' \cos \theta$ ,  $\theta$  is the angle between  $\hat{k}$  and  $\hat{k}'$  and carrying out the integration with respect to  $\theta$ , we get,

$$G = - \int_0^{\infty} \left[ \frac{\xi_0 (k^2 k'^4 - k^4 k'^2) kk'}{\nu(t-t_0)} \left\{ \exp \left[ -\frac{\nu}{p_M} (t-t_0) \{ (1+p_M)(k^2 + k'^2) - 2p_M kk' \} \right] - \right. \right. \\ \left. \exp \left[ -\frac{\nu}{p_M} (t-t_0) \{ (1+p_M)(k^2 + k'^2) + 2p_M kk' \} \right] \right\} \\ + \frac{\xi_1 (k^4 k'^6 - k^6 k'^4) kk'}{\nu(t-t_0)} \left( \omega^{-1} \exp \left[ -\omega^2 \left( \frac{(1+2p_M)k^2}{(1+p_M)^2} - \frac{2p_M kk'}{1+p_M} + k'^2 \right) \right] - \right. \\ \left. \omega^{-1} \exp \left[ -\omega^2 \left( \frac{(1+2p_M)k^2}{(1+p_M)^2} + \frac{2p_M kk'}{1+p_M} + k'^2 \right) \right] + \omega^{-1} \right. \\ \left. \exp \left[ -\omega^2 \left( k^2 - \frac{2p_M kk'}{1+p_M} + \frac{(1+2p_M)k'^2}{(1+p_M)^2} \right) \right] \right) \right] dt$$

$$\begin{aligned}
& -\omega^{-1} \exp\left[-\omega^2 \left(k^2 + \frac{2P_M k k'}{1+p_M} + \frac{(1+2p_M)k'^2}{(1+p_M)^2}\right)\right] \\
& + \{k \exp[-\omega^2 (1+p_M)(k^2 + k'^2) - 2p_M k k']\} \\
& -k \exp[-\omega^2 (1+p_M)(k^2 + k'^2) + 2p_M k k'] \} \int_0^{\frac{\omega k}{2}} \exp(x^2) dx + \{k' \exp[-\omega^2 (1+p_M)(k^2 + k'^2) - 2p_M k k'] \\
& -k' \exp[-\omega^2 (1+p_M)(k^2 + k'^2) + 2p_M k k'] \} \int_0^{\frac{\omega k'}{2}} \exp(x^2) dx \} dk' \quad (26)
\end{aligned}$$

$$\text{where } \omega = \left( \frac{\nu(t-t_1)(1+p_M)}{p_M} \right)^{\frac{1}{2}}.$$

Integrating Equ. (26) with respect to  $k'$ . We has

$$G = G_\beta + G_\gamma \quad (27)$$

where,

$$\begin{aligned}
G_\beta = & -\frac{\pi^{\frac{1}{2}} \xi_0 p_M^{\frac{5}{2}}}{\nu^{\frac{3}{2}} (t-t_0)^{\frac{3}{2}} (1+p_M)^{\frac{5}{2}}} \exp\left\{-\frac{\nu(t-t_0)(1+2p_M)k^2}{p_M(1+p_M)}\right\} \\
& \left[ \frac{15p_M k^4}{4\nu^2(t-t_0)^2(1+p_M)} + \left\{ \frac{5p_M^2}{(1+p_M)^2 \nu(t-t_0)} - \frac{3}{2\nu(t-t_0)} \right\} k^6 + \frac{p_M}{1+p_M} \left\{ \frac{p_M^2}{(1+p_M)^2} - 1 \right\} k^8 \right]
\end{aligned} \quad (28)$$

$$\text{and, } G_\gamma = G_{\gamma_1} + G_{\gamma_2} + G_{\gamma_3} + G_{\gamma_4}$$

The quantity  $G_\beta$  represents the transfer function arising owing to consideration of magnetic field at three point correlation equation;  $G_\gamma$  arises from consideration of the four –point equation. Integration over all wave number shows that

$$\int_0^\infty G.d\vec{k} = 0 \quad (29)$$

Indicating that the expression for  $G$  satisfies the conditions of continuity and homogeneity, physically, it was to be expected, since  $G$  is a measure of transfer of energy and the numbers must be zero. Hence

$$H = \exp\left[-\frac{2\nu k^2(t-t_0)}{p_M}\right] \int G \exp\left[-\frac{2\nu k^2(t-t_0)}{p_M}\right] dt + J(k) \exp\left[-\frac{2\nu k^2(t-t_0)}{p_M}\right]$$

where  $J(k) = \frac{N_0 k^2}{\pi}$  is a constant of integration and can be obtained as by Corrsin [2]

$$H = \frac{N_0 k^2}{\pi} \exp\left[-\frac{2\nu k^2(t-t_0)}{p_M}\right] + \exp\left[-\frac{2\nu k^2(t-t_0)}{p_M}\right] \int [G_\beta + (G_{\gamma_1} + G_{\gamma_2} + G_{\gamma_3} + G_{\gamma_4})] \exp\left[-\frac{2\nu k^2(t-t_0)}{p_M}\right] dt$$

where,  $G = G_\beta + G_{\gamma_1} + G_{\gamma_2} + G_{\gamma_3} + G_{\gamma_4}$

Then after integration equation

$$H = \frac{N_0 k^2}{\pi} \exp\left[-\frac{2\nu k^2(t-t_0)}{p_M}\right] + H_\beta + [H_{\gamma_1} + H_{\gamma_2} + H_{\gamma_3} + H_{\gamma_4}] \quad (30)$$

The terms  $H_\beta, H_{\gamma_1}, H_{\gamma_2}, H_{\gamma_3}$  and  $H_{\gamma_4}$  can be expressed as follows:

$$H_\beta = \frac{\xi_0 \pi^{\frac{1}{2}} p_M^{\frac{5}{2}}}{8\nu^{\frac{3}{2}} (1+p_M)^{\frac{7}{2}}} \exp\left(\frac{-\nu(t-t_0)(1+2p_M)}{p_M(1+p_M)}\right) k^2$$

$$\left[ \frac{3p_M k^4}{2\nu^2(t-t_0)^{5/2}} + \left( \frac{(7p_M^2 - 6p_M)}{3\nu(1+p_M)(t-t_0)^{3/2}} \right) k^6 - \left( \frac{(3p_M^2 - 2p_M + 3)}{3(1+p_M^2)(t-t_0)^{1/2}} \right) k^8 \right.$$

$$\left. + \left( \frac{8\nu^{1/2}(3p_M^2 - 2p_M + 3)}{3(1+p_M)^{5/2} p^{1/2}_M} \right) k^9 F(\omega) \right],$$

$$F(\omega) = \exp(-\omega^2) \int_0^\omega \exp(x^2) dx, \omega = \left[ \frac{\nu(t-t_0)}{p_M(1+p_M)} \right]^{1/2} k,$$

Here  $H_1$  and  $H_2$  magnetic energy spectrum arising from consideration of the three and four –point correlation equations respectively. The total magnetic turbulent energy is

$$\frac{\langle h_i h'_i \rangle}{2} = \int_0^\infty H dk \quad (31)$$

here

$$\int_0^{\infty} H_1 dk = \frac{N_0 p^{\frac{3}{2}} M v^{\frac{-3}{2}} (t-t_0)^{\frac{-3}{2}}}{8\sqrt{2\pi}} + \xi_0 Q v^{-6} (t-t_0)^{-5},$$

$$\int_0^{\infty} H_2 dk = \xi_1 [R v^{\frac{-17}{2}} (t-t_1)^{\frac{-15}{2}} + S v^{\frac{-19}{2}} (t-t_1)^{\frac{-17}{2}}], L_1 = Q_2 + Q_4 + Q_6 + Q_7, L_2 = Q_1 + Q_3 + Q_5$$

By using above values, Equ. (31) we get

$$\frac{\langle h_i h'_i \rangle}{2} = \frac{N_0 p^{\frac{3}{2}} M v^{\frac{-3}{2}} (t-t_0)^{\frac{-3}{2}}}{8\sqrt{2\pi}} + \xi_0 \cdot Q \cdot v^{-6} \cdot (t-t_0)^{-5}$$

$$+ [\xi_1 L_1 v^{\frac{-17}{2}} (t-t_1)^{\frac{-15}{2}} + \xi_1 L_2 v^{\frac{-19}{2}} (t-t_1)^{\frac{-17}{2}}] \quad (32)$$

$$<T^2> = \langle h^2 \rangle = A(t-t_0)^{-3/2} + B(t-t_0)^{-5} + C(t-t_1)^{-15/2} + D(t-t_1)^{-17/2}, \quad (33)$$

This is the energy decay law of MHD turbulence for four point correlations. where,

$$<T^2> = \langle h^2 \rangle = \langle h_i h'_i \rangle, A = \frac{N_0 p^{\frac{3}{2}} M v^{\frac{-3}{2}}}{4\sqrt{2\pi}}, B = 2 \xi_0 Q v^{-6}, C = 2 \xi_1 L_1 v^{\frac{-17}{2}} \text{ and } D = 2 \xi_1 L_2 v^{\frac{-19}{2}}.$$

If  $L_1=0$  and  $L_2=0$  that is  $C=0$  and  $D=0$  in Equ. (33) than we get,

$$<T^2> = \langle h^2 \rangle = A_1 (t-t_0)^{-3/2} + B_1 (t-t_0)^{-5} \quad (34)$$

This is the energy decay of MHD turbulence in three- point correlations which was obtained earlier by Sarker and Kishore [12]

**Table-1:**The value of the constants and parameter used in Equ. (33)

Fluid	$P_M$	$v$	$N_0$	$\xi_0$	$\xi_1$	A	B	C	D
Mercury	0.015	0.10	.1	.01	.02	.00058	$4.18 \times 10^{-7}$	$3.69 \times 10^{-13}$	5.87
	0.015	0.08	.1	.01	.02	.00081	$1.6 \times 10^{-6}$	$-1.01 \times 10^{-12}$	20.03

Mix Gas	0.2	80	.1	.01	.02	$1.15 \times 10^{-6}$	$5.75 \times 10^{-18}$	$3.78 \times 10^{-16}$	$9.95 \times 10^{-13}$
	0.2	200	.1	.01	.02	$3.15 \times 10^{-7}$	$2.36 \times 10^{-20}$	$6.12 \times 10^{-18}$	$6.44 \times 10^{-15}$
Hyd Gas	.04	100	.1	.01	.02	$2.5 \times 10^{-6}$	$6.8 \times 10^{-17}$	$2.7 \times 10^{-14}$	$9.79 \times 10^{-13}$
	0.4	300	.1	.01	.02	$4.86 \times 10^{-7}$	$9.4 \times 10^{-20}$	$1.9 \times 10^{-16}$	$2.3 \times 10^{-15}$
Hel Gas	0.7	120	.1	.01	.02	$4.6 \times 10^{-6}$	$4.8 \times 10^{-16}$	$7.4 \times 10^{-13}$	$9.4 \times 10^{-23}$
	0.7	400	.1	.01	.02	$7.6 \times 10^{-7}$	$3.4 \times 10^{-19}$	$3.3 \times 10^{-15}$	$1.2 \times 10^{-15}$

### The graphical representations and explanations

Figure of Equ.(33) for  $P_M = 0.015$ :




Figure1 (a):Sketch of Equ.(33).Figure1 (b):Sketch of Equ.(33)

Figure 1(a), Figure 1(b) represents the energy decay curve for four-point correlations of Equ (33). When the Prandtl no. is small as of mercury  $P_M = 0.015$  and It is observed that the energy decreases more rapidly as viscosity decreases.

Figure of Equ.(33) for  $P_M = 0.2$ :





Figure-(2a): Sketch of Equ.(33)

Figure-(2b): Sketch of Equ.(33)

Figure-(2a) and Figure-(2b) are the energy curve of Equ.(33) when the Prandtl no. is as of mixture of gas for  $P_M = 0.2$  and  $\nu = 80$  in fig. (2a) and  $\nu = 200$  in fig. (2b). In this case, energy decreases rapidly as viscosity decreases.

Figure of Equ.(33) for  $P_M = 0.4$ :





Figure-(3a):Sketch of Equ.(33)

Figure-(3b):Sketch of Equ.(33)

Figure-(3a), Figure-(3b) indicate the curve of energy Equ.(33). When the Prandtl no. is as of Hydrogen gas,  $P_M = 0.4$  and  $\nu = 100$  and  $\nu = 300$ . Result: Energy decreases as well as viscosity decresies.

Figure of Equ.(33) for  $P_M = 0.7$ :




Figure-(4a):Sketch of Equ (33)

Figure-(4b):Sketch of Equ (33)

Figure-(4a) and Figure-(4b) are the energy curve of Equ (33). When the Prandtl No. is as of Helium gas  $P_r=0.7$  and  $\nu=120$  and  $\nu=400$ . Energy decreases rapidly as viscosity decreases from 400 to 120.

Comparing fig (1a)-(4b): we see that Energy changes rapidly as Prandtl no. changes. Figure (1)-(4):  $y_1, y_2, y_3, y_4, y_5$  and  $y_6$  are represented the energy decay curves of MHD turbulence for four-point correlations of Equ (33) at several times. From figure 1 and Figure 4, we see that, in four-point correlations system energy die out faster than the three-point correlations system in MHD turbulent flow.

#### Comparison between four-point and three point correlations of equation:




Fig (5a): Energy curves of Equ (33) Fig (5b): Energy curves of Equ (34)

Fig-(5a) and Fig (5b) represents the energy decay curve for four-point and three-point correlations of equation. When the Prandtl no. is small as of mercury  $P_M=0.015$ . It is clear that, in four-point correlations energy decreases more rapidly than three point correlations.

Figure of Equ.(33) and (34) for large Prandtl No.




Fig (6a): Energy curves of Equ.(33) Fig (6b): Energy curves of Equ.(34)

When the Prandtl no. is as of mixture of gas  $P_M = 0.2$  i.e. for large Prandtl no. we conclude that, energy at four- point correlations and three -point correlations has no change significantly.

Figure of Equ.(33) and (34) for Hydrogen gas:




Fig (7a) Energy curves of equ.(33)

Fig (7b) Energy curves of equ.(34)

Fig-(7a) and Fig-(7b) indicate the energy curve equation (33) and (34). When the Prandtl No. as of Hydrogen gas  $P_M = 0.4$ .

We observed that there is no change in energy for four point and three point correlations as for same viscosity

## Conclusion

- For mercury, I observed that the energy decreases more rapidly as viscosity decreases.
- In Helium gas for  $P_r = 0.7$  and  $\nu = 120$  and  $\nu = 400$ . Energy decreases rapidly as viscosity decreases.



- It is observed that the decay law for four-point correlations systems energy decreases rapidly more and more by exponential manner than the decreases of three point correlation systems.
- We observed that there is no change in energy for four point and three point correlations as for same viscosity.
- If the time increases than energy decay also increases.
- We finally conclude that from all above the figures that energy decreases as viscosity and Prandtl number decrease.

## References

- [1] Boussinesq, J. 1877: Mem. press. pardiv. Savants al' Acad. des. Sci. Paris 23, 46.
- [2] Corrsin, S. (1951b): J. Applied Physics, 22, 469.
- [3] Ferit, J. Kampe de (1948): La Houille Blanch, Grand Rub, Grenoble, France, 23, 1.
- [4] Hopf, E. (1957): In Proc. Symp. Appl. Math, 7, American Mathematical Society, 41.
- [5] Hopf, E. (1962: In Proc, Symp. Appl. Math.13, American Mathematical society, 157.
- [6] Kolmogoroff, A. N. (1941): Academia de Science de URSS xxx No. 4.
- [7] Kolmogoroff, A. N. (1962): Fluid Mech., 13, Part-1.
- [8] Monin, A.S. and Yaglom A.M. (1975): Statistical Fluid Mechanics, 11, M.I.T. Press.
- [9] Monin. A.S. and yaglom, A.M. (1973): Statistical Fluid Mechanics, 1. M.I.T. Press.
- [10] Prandtl, L. (1925): Z. Angew. Math. u. Mech., 5, 136.
- [11] Renolds, O. (1883): Phil. Trans. Roy. Soc., London, 174, 935.
- [12] Sarker, M.S.A. and Kishore, N. (1991b): Int. J. Engng. Sci., 29, 1479.
- [13] Taylor, G. I. (1935a): Proc. Roy. Soc., London, A151, 421..
- [14] Taylor, G. I. (1935b): Z. Angew. Math. Mech., 15, 91.
- [15] Taylor, G. I. (1921): Proc. Lond. Math. Soc., 20, 196.
- [16] Taylor, G. I. (1938b): Proc. Roy. Soc., London, A164, 476.
- [17] Von Karman, T. (1937a): Proc. U. S. Nat. Acad., Wash, 23, 98.
- [18] Von Karman, T. (1937b): J. Aero. Sci., 4, 134.
- [19] Kraichnan, R. H. (1959): J. Fluid Mech., 5, 497.

- [20] Deissler, R. G. (1960): Physics of Fluids, 3, 176.
- [21] Deissler, R. G. (1961): Technical Rept R-96, NASA.
- [22] M. A. Bkar. Pk, M. A. K. Azad and M.S.Alam Sarker, "Decay of energy of MHD turbulence for four-point correlation. International Journal of Engineering Research and Technology". Vol. 1 Issue 9, November-2012, pp 1-13.
- [23] M.A.BkarPk, M.S.AlamSarker and M.A.K.Azad, "Decay of MHD turbulence prior to the ultimate phase in presence of dust particle for four-point correlation", International Journal of Applied Mathematics and Mechanics. Int. **J.** of Appl. Math and Mech. **9**(10):34-57, 2013.
- [24] M. A. Bkar. Pk, M. A. K. Azad and M.S.Alam Sarker, "Decay of dusty fluid MHD turbulence for four-point correlation in a rotating system", **J. Sci. Res.** **5**(1), 77-90(2013).
- [25] M.AbuBkarPk, Abdul Malek, M.AbulKalamAzad, "Effects of first-order reactant on MHD turbulence at four-point correlation. Applied and Computational Mathematics", 4(1): 11-19; 2015.
- [26] Abu BkarPk, Abdul Malek, M.AbulKalam Azad , " *4-Point Correlations of Dusty Fluid MHD Turbulent Flow in a 1st order Chemical-Reaction*", Global Journal of Science Frontier Research, *GJSFR-F (2015) Volume 15 Issue 2: 53-69.*
- [27]M. A. Bkar**Pk** , M.A.K. Azad."Decay of Energy of MHD Turbulence before the Final Period for Four-point Correlation in a Rotating System Undergoing a First Order Chemical Reaction", International Journal of Scholarly Research Gate, 3(4):PP: 143-160, 2015.
- [28] M. MonuarHossain,M. Abu BkarPk, M .S. AlamSarker "Homogeneous fluid turbulence before the final period of decay for four-point correlation in a rotating system for first-order reactant", American Journal of Theoretical and Applied Statistics 3(4): 81-89, 2014.
- [ 29]M. AbulKalam Azad, M. Abu BkarPk, Abdul Malek, "Effect of Chemical Reaction on Statistical Theory of Dusty Fluid MHD Turbulent Flow for Certain Variables at Three- Point Distribution Functions," Science Journal of Applied Mathematics and Statistics. Volume 3 Issue 3: 75-98.2015.
- [30] M. A. BkarPk&RipanRoy,"Effect of First Order Chemical Reaction for Coriolis Force and Dust Particles for Small Reynolds Number in the Atmosphere Over Territory," Global Journal of Science Frontier Research: H Environment & Earth Science. Volume 16 Issue 1, PP:39-50.2016.
- [31] M. AbulKalam Azad, M. Abu BkarPk, Abdul Malek,Effect of Chemical Reaction on Statistical Theory of Dusty Fluid MHD Turbulent Flow for Certain Variables at Three- Point Distribution Functions. Science Journal of Applied Mathematics and Statistics. 2015; 3(3): 75-98.
- [32]Shimin Yu , Tingting Tang , Jianhui Li and Peng Yu, "Effect of Prandtl Number on Mixed Convective HeatTransfer from a Porous Cylinder in the Steady Flow Regime." Entropy **2020**, 22, 184;
- [33] Abdul MalekPh.DThesis analytical investigations on some problems of turbulent and magneto-hydrodynamic turbulent. Awarded in 2018, Dept of Applied Mathematica, R.U