

Effect of different amendments on nutrients availability of sodic soils

Abstract

A pot house ~~study~~experiment was conducted at the Net House, Regional Research Station, AAU, Anand during *rabi* season of 2018-19 on Wheat (GW-496) as a indicator crop. Total four amendments i.e. A₀(control), A₁(Gypsum @ 50% GR), A₂(Vermi-compost @ 4.0 t ha⁻¹)&and A₃ (sulphur @ 50 kg ha⁻¹) and six soils type (S1 to S6) were selected under Factorial CRD. Pot study results indicated that the soil amendments significantly decreased the pH, EC and ESP of sodic soils. The order of decrease in pH of sodic soils from 8.75 to 7.89 by application of amendments remained as: vermi-compost > sulphur > gypsum. The EC of sodic soils significantly decreased due to application of vermi-compost and gypsum by 20.48% and 10.84%~~-per-cent~~, respectively over control. The application of vermi-compost, gypsum and sulphur significantly lowered down ESP from 18.45 to 12.44, 13.00 and 13.57, respectively, and the effect of gypsum was found at par with sulphur in decreasing soil ESP. Application of vermi-compost (4.0 t ha⁻¹) significantly influenced CEC (28.28 Cmol (P⁺) kg⁻¹) over control (no amendment). Irrespective of amendments, soil S5 recorded the highest pH (8.39), EC (0.92 dSm⁻¹), ESP (16.48) and the lowest was in soil S6 (pH 7.99, EC 0.64 dSm⁻¹). The interaction of soil x amendment treatments showed significant effects on pH, EC and ESP of sodic soils. The application of amendments showed significant effect on available nutrients in soil after harvest of wheat and the highest OC (0.73%), available P₂O₅ (65.71 kg ha⁻¹), available K₂O (207.1 kg ha⁻¹) content in soil was recorded when vermi-compost applied at 4.0 t ha⁻¹ followed by sulphur at 50 kg ha⁻¹ and gypsum at 50% GR but available S (16.33 mg kg⁻¹) was recorded the highest due to sulphur application at 50 kg ha⁻¹ followed by gypsum and vermi-compost.

Key words: amelioration, amendment, sodic soil, ESP, CEC

Introduction:

The ever increase in the world's *population* needs food productivity to step up within a few decades. Unfortunately, extensive areas of irrigated lands are unproductive, due to the sodicity and salinity problems in the soil profile occupied by root systems. According to ~~one~~ estimate study, an area of 6.74 mha in India suffers from salt accumulation out of which 3.78 mha are sodic, while 2.96 mha are saline soils. Among different states, Gujarat rank first (22 lakh ha.) in area of salt affected soil, while alkali soils is 5.41 lakh ha (Anon., 2021). These soils are distributed in the districts of Kutch, Patan and Mahesana in north Gujarat, Anand, Ahmadabad and Kheda in middle Gujarat, Surendranagar, Rajkot, Amreli, Bhavnagar, Porbandar, Junagadh and Jamnagar in Saurashtra and Valsad, Surat and Bharuch districts of south Gujarat. (Anon., 2021). Excess salt accumulations adversely affect soil physical, chemical and biological properties of soil results poor fertility status of salt affected soil. The sodic soils have $pH_e > 8.5$, $EC_e < 4 \text{ dSm}^{-1}$ at 25 °C, $ESP > 15$ and $SAR > 13$. So, for

Comment [N1]: Reference

increasing food production we have to utilize the salt affected soil which is increasing day by day. Reclamation has been proved to be an effective way to improve soil quality of salt affected soils. Theoretically, two steps are needed to perform sodic soil reclamation. The first step is remove Na^+ from the cation exchange sites in the soil colloid, and the second step is leaching out the replaced Na^+ in the plant root zone (Tripathi *et al.*, 2016). Amelioration of sodic and alkali soil primarily involves increasing Ca^{2+} on the cation exchange complex at the expense of Na^+ . The replaced Na^+ is removed from the root zone through infiltrating water resulting from excessive irrigation(s). The combine application of inorganic-organic amendments like farm yard manure, vermi-compost, castor cake, gypsum, elemental sulphur, sulphuric acid, etc. improve physical, chemical, and biological properties of soil. The biological amelioration methods which is using living or dead organic matter (crops, stems, straw, green manure, barnyard manure, compost, sewage sludge) have two beneficial effects on the— alkali soils reclamation,— first is the improvement of the soil structure and permeability, thus enhancing salt leaching, reducing surface evaporation, and inhibiting salt accumulation in the surface layers; and the second is release of carbon dioxide during respiration and decomposition. For saline or sodic soils, the addition of organic matter (OM) can accelerate the leaching of Na, decrease the ESP and electrical conductivity (EC), and increase water infiltration, water-holding capacity, and aggregate stability (Lax *et al.* 1994; Qadir *et al.* 2001). So here in this study we have used different amendments to know the effect on reclamation of sodic soils.

Materials and methods

The survey work was carried out in Anand and Kheda districts by Patel (2016) and Vaghela (2016), respectively. On the basis of 160 surface soil samples analysis, pot study was carried out on selected six soils categories to study the amelioration of sodic soils by using different amendments and its effect on soil properties and yield of wheat (*Triticum aestivum* L.). Results indicated that 20 and 24 % soil samples categorised under sodic soils i.e. soil having $\text{pH} > 8.5$ and $\text{ESP} > 15.0\%$ and $\text{ECe} < 4.0 \text{ dSm}^{-1}$. Six soils (3 from each district) were selected i.e. from Anand and Kheda district of middle Gujarat region. In survey work, Soil analysed for chemical parameters viz., pH, EC, SOC, available N, P, K, S, micronutrients (Fe, Mn, Zn, Cu), CEC, ESP, exchangeable Ca, Mg, K, Na, CO_3^{2-} , HCO_3^- , Cl^- ~~and~~ gypsum requirements. Experiment was designed under Factorial CRD with 24 treatments and 3 replications. After harvest of the crop, pot wise soil samples were collected from approximately 20cm depth and air dried in laboratory. The air dried soil samples were pounded with wooden mortar and pestle. After pounding, the soils were sieved through 2 mm

Comment [N2]: Rewrite this sentence

sieve and preserved in polythene bags properly ~~labeled~~labelled. The soil sample was analysed for the pH, EC, CEC, ESP, GR, organic C, available N, P₂O₅, K₂O and Micronutrients (Fe, Mn, Zn, Cu) as per standard methods.

Table 1: The chemical properties of initial soil sample

Chemical parameters	Soil types (Initial value)					
	S1	S2	S3	S4	S5	S6
pH (1:2.5)	8.78	8.88	8.90	8.84	9.05	8.65
EC (1:2.5) dSm ⁻¹	0.80	0.82	0.98	0.85	1.01	0.68
Organic Carbon (%)	0.19	0.24	0.57	0.75	0.49	0.60
Available P ₂ O ₅ (kg ha ⁻¹)	19.31	64.50	61.84	85.87	49.12	74.58
Available K ₂ O (kg ha ⁻¹)	112.51	116.20	320.14	165.10	185.30	130.32
Available S (mg kg ⁻¹)	8.13	12.54	10.64	11.85	6.60	10.50
CEC (Cmol (P ⁺)kg ⁻¹)	22.40	29.05	22.45	32.10	24.55	28.30
ESP	16.67	20.00	22.25	18.53	22.97	17.85
GR (t ha ⁻¹)	7.16	11.15	9.92	11.40	10.78	9.69
DTPA- Fe (ppm)	9.10	10.15	8.00	9.20	12.85	11.14
DTPA- Mn (ppm)	7.10	12.90	9.60	11.80	9.30	6.00
DTPA- Zn (ppm)	0.80	1.00	0.99	0.90	0.42	0.85
DTPA- Cu (ppm)	2.00	2.10	1.25	1.00	2.63	2.14

DTPA= Diethylenetriamine pentaacetate, ESP= exchangeable sodium percentage

Table 2: Chemical Compositions of Amendments:

Amendments	N (%)	P (%)	K (%)	Ca %	Mg %	S (%)	Fe %	Mn %	Zn %	Cu %
Vermicompost	1.12	0.33	0.80	2.28	0.41	0.20	0.45	0.02	0.06	0.007
Gypsum	-	-	-	24	-	16	-	-	-	-
Sulphur	-	-	-	-	-	90	-	-	-	-

Results and discussion:

Effect on pH

~~A figure.1 shows that a~~All the amendments significantly decreased the pH of soil (Fig.1). A vermi_compost (4.0 t ha⁻¹) application significantly decreased the pH (7.89), ~~followed~~followed by sulphur (50 kg ha⁻¹) and gypsum (50% GR)(Table 3). The order of decrease in pH of soil from 8.75 to 7.89 by application of amendments remained as: vermi_compost>sulphur>

gypsum. ~~The significantly h~~ Highest pH value was observed due to interaction effect of S₅A₀ treatments (no amendment) which was significantly decreased with application of amendments. Formation of carbonic acid from CO₂ produced during the decomposition of organic residues in soils is the main component responsible for decreasing pH (M. Jalali *et al.*, 2020). ~~The a~~ Application of organic amendments leads to increased CO₂ partial pressure within soil profile, lowered soil pH value in soil solution and subsequently increased native CaCO₃ mineral dissolution and reduces the soil sodicity (Li and Keren, 2009).

Effect on EC

~~A v~~ Vermicompost and gypsum significantly decreased the EC of sodic soils by 20.48% and 10.84% ~~per cent~~, respectively over control. There was no practically change in EC due to sulphur application (Fig. 2). ~~As per data in Table 3~~, Irrespective of amendments, soil S₅ recorded the highest EC (0.92 dSm⁻¹), while it was lowest in S₆ (0.62 dSm⁻¹), being at par with S₁ (0.64 dSm⁻¹). Interaction effect of S₃A₃ and S5A3 combinations found significantly superior over others but was at par with S3A0 and S5A0. For saline or sodic soils, the addition of organic matter (OM) can accelerate the leaching of Na, decrease the ESP and electrical conductivity (EC), and increase water infiltration, water-holding capacity, and aggregate stability (Lax *et al.*, 1994; Qadir *et al.*, 2001). However, overall evaluation indicated that EC values of soils treated with organic fertilizers were below the EC value of soil with chemical fertilizer. This is possibly due to the fact that, unlike chemical fertilizers, organic fertilizers release nutrients more gradually as a result of degradation process and their chelating effect (Gerke *et al.*, 1999; Sharma and Banik, 2014).

Effect on ESP

Application of amendments significantly decreased the ESP of soils. The application of vermicompost, gypsum and sulphur significantly lowered down ESP from 18.45 to 12.44, 13.00 and 13.57, respectively, the effect of gypsum was found at par with sulphur in decreasing soil ESP. However, all the amendments reclaimed soil by reducing the ESP below 15 (Fig. 3). ~~With regards to different soils, data given in the Table 3 indicated it shows~~ that soil S5 recorded with highest ESP (16.48) (Table 3). Almost in all soils, the ameliorating effect of amendments were observed by reducing the ESP below 15 except soil ~~soil~~ S5. Among above amendments vermicompost was most efficient to decrease soil ESP, might be due to amendments alleviating the harmful effects of sodicity by replacing the Na⁺ from exchange site, also after leaching of Na⁺ from root zone, crop might also benefited by the improved physical properties of soil leading to more crop growth in these treatments (Hussain *et al.*, 2001; Tzanakakis *et al.*, 2011; Mohamed *et al.*, 2012).

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Effect on CEC

Application of vermi-compost (4.0 t ha^{-1}) significantly influenced CEC ($28.28 \text{ Cmol (P}^+) \text{ kg}^{-1}$) over control (no amendment). There were no significant effect of sulphur, gypsum was found at par with control indicating there were no effect of gypsum and sulphur on CEC of soils (Table 3). The increased in CEC of soil due to vermi-compost might be due to increased in fine organic particle which increased the total exchange sites of cations. The highest CEC was found in soil S4 ($32.37 \text{ Cmol (P}^+) \text{ kg}^{-1}$), while the lowest in S3 soil. The organic amendments improve soil cation exchange capacity (CEC) through humus formation and increase concentrations of nutrients such as Ca and Mg as a result of decomposition (Jenkinson, 1990; Johnston, 1997). Havlin *et al.* (1999) indicated that the addition of different types of compost into the soil increased CEC varying from 20% to 70% of initial values. The increase in CEC after the application of biological amendments has also been reported in previous researches (Seenivasan *et al.* 2015; Nisha *et al.* 2018).

Comment [N3]: value insert




Fig. 1: Effect of amendments on pH of soils




Fig. 2: Effect of amendments on electrical conductivity of soils




Fig. 3: Effect of amendments on ESP of soils




Fig. 4: Effect of amendments on CEC of soils

Table 3: Amelioration effect of amendments on chemical properties of soils.

Sr. No.	Treatments	pH _(1:2.5)	EC _(1:2.5) (dSm ⁻¹)	ESP	CEC(Cmol (P ⁺) kg ⁻¹)
Types of Soils					
1	S1	8.16	0.64	12.90	23.22
2	S2	8.29	0.74	14.73	29.40
3	S3	8.31	0.88	14.69	23.32
4	S4	8.12	0.78	13.15	32.37
5	S5	8.39	0.92	16.48	24.41
6	S6	7.99	0.62	14.25	28.86
CD (P=0.05)		0.07	0.03	1.37	1.06
Types of amendments					
1	A0 (control)	8.75	0.83	18.45	26.60
2	A1 (Gypsum @ 50% GR)	8.17	0.74	13.00	26.65
3	A2 (Vermicompost @ 4.0 t ha⁻¹)	7.89	0.66	12.44	28.28
4	A3 (sulphur @ 50 kg ha⁻¹).	8.02	0.84	13.57	26.20
CD (P=0.05)		0.05	0.02	0.97	0.75
Interaction (S × A)		0.14	0.06	NS	NS
CV%		1.04	4.82	11.67	4.85

Table 4: Amelioration effect of amendments on nutrient status -of soils.

Sr. No.	Treatments	OC %	Avail. P ₂ O ₅ (kg ha ⁻¹)	Avail. K ₂ O (kg ha ⁻¹)	Avail. S (mg kg ⁻¹)	Avail. Fe (mg kg ⁻¹)	Avail. Zn (mg kg ⁻¹)	Avail. Mn (mg kg ⁻¹)	Avail. Cu (mg kg ⁻¹)
Types of soil									
1	S1	0.41	26.53	126.9	12.54	9.73	0.95	8.03	2.17
2	S2	0.32	68.94	150.9	16.76	11.27	1.23	13.99	2.24
3	S3	0.67	63.90	326.6	14.07	8.80	1.09	10.50	1.35
4	S4	0.89	88.87	181.5	13.19	10.20	1.04	12.96	1.34
5	S5	0.63	51.51	197.1	9.34	13.85	0.57	9.98	2.73
6	S6	0.85	77.77	143.3	10.77	12.64	1.02	7.21	2.29
CD (P=0.05)		0.05	2.07	11.6	1.19	0.24	0.05	0.17	0.08
Types of amendments									
1	A0 (control)	0.54	60.87	172.2	9.22	10.38	0.88	9.81	1.96
2	A1 (Gypsum @ 50% GR)	0.62	62.93	183.2	13.77	11.29	1.00	10.65	2.02
3	A2 (Vermicom post @ 4.0 t ha⁻¹)	0.73	65.71	207.1	11.80	11.41	1.03	10.68	2.06
4	A3 (sulphur@ 50 kg ha⁻¹).	0.62	62.19	188.4	16.33	11.25	1.00	10.65	2.03
CD (P=0.05)		0.03	1.46	8.1	0.85	0.17	0.03	0.12	0.06
Interaction (S × A)		NS	NS	NS	NS	NS	NS	NS	NS
CV%		9.14	4.01	7.52	11.42	2.61	6.49	1.98	1.60

Effect on OC, P, K & S

The amendments showed significant effect on soil organic carbon content after harvest of wheat crop (Table 4). The results indicated that the highest OC was found due to vermi-compost application at 4.0 t ha⁻¹ (0.73%) followed by sulphur at 50 kg ha⁻¹ (0.62%) and gypsum at 50% GR (0.62%). A vermi-compost at 4.0 t ha⁻¹ increased soil OC by 35.18 per cent more over control treatment. The organic matter addition through vermi-compost increased OC content in soil. Singh *et al.*, (2009) reported that addition of pressmud increased the soil organic carbon due to increase in yield and root biomass. The significant effect of vermi-compost and gypsum was observed over control on soil available P₂O₅ in soil after harvest of wheat. The significant highest available P₂O₅ was recorded due to Vermicompost at 4.0 t ha⁻¹ (65.71 kg ha⁻¹) but the effect of sulphur and gypsum on available P₂O₅ was found at par. The available P₂O₅ was significantly influenced in soils S2 to S6 over soil S1 and the highest available P₂O₅ was recorded in soil S4 (88.87 kg ha⁻¹). The interaction effect of different combination was found non- significant on available P₂O₅ in soil. The perusal of data (Table 4) It indicated that amendments showed significant effect on available K₂O of soil over control after harvest of wheat crop (Table 4). The highest available K₂O was

found with vermi-compost application at 4.0 t ha^{-1} i.e., 207.1 kg ha^{-1} . The available K_2O was also increased under sulphur application next to vermi-compost, being at par with gypsum. The highest available K_2O was recorded in soil S3 (326.6 kg ha^{-1}) and the lowest in S (126.9 kg ha^{-1}). When organic materials are applied to soils it improve organic matter status, and also supply soils with nutrients including N, P and K (Tiwari *et al.*, 2004; Zakir *et al.*, 2012); ~~Also supported by~~ Xu *et al.*, (2019). Reports also suggested that especially phosphorus in vermi-compost is released more gradually in available form in soil (Kale *et al.*, 1987; Doube and Brown, 1998; Nethra *et al.*, 1999; Lazcano *et al.*, 2008). Azarmi *et al.* (2008) showed that addition of vermi-compost at 15 t ha^{-1} significantly ($P < 0.05$) increased contents of soil total organic carbon, total N, P, K, Ca, Zn and Mn substantially compared with control plots. The addition of vermi-compost in soil resulted in decrease of soil pH. The physical properties such as bulk density and total porosity in soil amended with vermi-compost were improved. The results of this experiment revealed that addition of vermi-compost had significant ($P < 0.05$) positive effects on the soil chemical ~~and~~ physical properties.

The data presented in Table 4, indicated that amendments significantly increased the available S content of soil. The application of sulphur (50 kg ha^{-1}) showed the highest available S (16.33 mg kg^{-1}), followed by gypsum at 50% GR (13.77) and vermi-compost at 4.0 t ha^{-1} (11.80 mg kg^{-1}). The available S was recorded the highest in soil S2 (16.76 mg kg^{-1}), while the lowest in S5 (9.34 mg kg^{-1}). Application of sulphur increased the available S might be due to sulphur containing compounds are acid forming materials which release sulphuric acid on oxidation in soil resulting a favourable condition for crop growth particularly in alkaline soils (Singh *et al.*, 1981; Ghafoor and Muhammad, 1981).

Effect on micronutrients:

The amendments showed significant effect on DTPA extractable micronutrients (Fe, Mn, Zn, Cu) after harvest of wheat in sodic soils. The application of vermi-compost (4.0 t ha^{-1}) recorded numerically high available Fe, Mn, Zn ~~&and~~ Cu i.e., 11.41, 10.68, 1.03 ~~&and~~ 2.06 mg kg^{-1} respectively, being at par with gypsum (50% GR) and sulphur (50 kg ha^{-1}). Which supported by decrease in soil pH. Irrespective of amendments, soils S5, S4, S2 ~~&and~~ S5 recorded the highest available Fe, Mn, Zn ~~&and~~ Cu, while the lowest in S3, S6, S5 ~~&and~~ S4, respectively. In study, the effect of vermi-compost found better over other amendments in increasing micronutrient availability of soil. This might be due to the micro-organisms in the vermi-products play a significant role in altering the soil micronutrient content. The microbial activity induced by the bio-fertilizers increase the soil microorganism activity by availing

additional substances that are not found in chemical fertilizers. Azarmi *et al.* (2008) showed that addition of vermi-compost application at 15 t ha⁻¹ significantly ($P < 0.05$) increased contents of soil total organic carbon, total N, P, K, Ca, Zn and Mn substantially compared with control plots.

Conclusion:

The amendments significantly decreased the pH, EC and ESP of soil. The order of decrease in pH of sodic soils from 8.75 to 7.89 by application of amendments remained as: vermi-compost > sulphur > gypsum. The amendments also showed significant effect on soil OC, available P₂O₅, K₂O, SO₄-S, DTPA extractable Fe, Mn, Zn, Cu after harvest of wheat and the highest OC (0.73%), available P₂O₅ (65.71 kg ha⁻¹), available K₂O (207.1 kg ha⁻¹) was recorded when vermicompost applied at 4.0 t ha⁻¹ followed by sulphur at 50 kg ha⁻¹ and gypsum at 50% GR but the highest available S (16.33 mg kg⁻¹) was recorded due to sulphur application @ 50 kg ha⁻¹ followed by gypsum and vermicompost.

Competing interest:

Authors contribution:

Acknowledgement:

References:

- Anonymous (2016-17). Area and production of wheat in India, 2nd advance estimates, Ministry of Agriculture & Farmers Welfare, Govt. of India. [page??](#)
- Anonymous (2021). Central Soil Salinity Research Institute, Karnal. [page??](#)
- Azarmi, R., Giglou, M. T. & Taleshmikail, R. D. (2008). Influence of vermicompost on soil chemical and physical properties in tomato (*Lycopersicon esculentum*) field. *African Journal of Biotechnology*, 7 (14), 2397-2401.
- Doube, B. M. & Brown, G. G. (1998). Life in a complex community: Functional interactions between earthworms, organic matter, microorganisms, and plants. In: Edwards C (Ed). *Earthworm ecology*, St Lucie Press, London, 179-211.
- Gerke, H. H., Arning, M. & Stoppler-Zimmer, H. (1999). Modeling long-term compost application effects on nitrate leaching. *Plant and Soil*, 213, 75-92.
- Ghafoor, A. & Muhammed, S. (1981). Comparison of H₂SO₄, HCl, HNO₃ and gypsum for reclaiming calcareous saline-sodic soil and for plant growth. *Bull. (J) Irrigation, Drainage and Flood Control Research*. (Pakistan), 11, 69-75.
- Havlin, L., Beaton, J.D., Tisdale, S.L. & Nelson, W.L. (1999). Soil Fertility and Fertilizers: An Introduction to Nutrient Management. Prentice Hall, Englewood Cliffs. [page?](#)
- Jalali, M., Lotf, M. S. & Ranjbar, F. (2020). Changes in some chemical properties of saline-sodic soils over time as affected by organic residues: an incubation study. *Polish Journal of Soil Science*, 53, 79.

Comment [N4]: Please write this section again

Comment [N5]: Please refer according to journal rules

- Kale, R. D., Bano, K., Sreenivasa, M. N., Vinayak, K. ~~&~~ Bagyaraj, D. J. (1987). Incidence of cellulolytic and lignolytic organisms in the earthworm worked soils. In: Julka JM (Ed). *Eartworm resources and vermiculture*, Oxford University Press, New Delhi, India, 49-53.
- Lax, A., Diaz, E., Castillo, V. ~~&~~ Albaladejo, J. (1994). Reclamation of physical and chemical properties of a salinized soil by organic amendment. *Arid Soil Research and Rehabilitation*, 8, 9-17.
- Lazcano, C., Gomez-Brandon, M., ~~and~~ Dominguez, J. (2008). Comparison of the effectiveness of composting and vermicomposting for the biological stabilization of cattle manure. *Chemosphere*, 72, 1013-1019.
- Li, F. ~~&~~ Keren, R. (2009). Calcareous sodic soil reclamation as affected by corn stalk application and incubation: a laboratory study. *Pedosphere*, 9, 465-475.
- Nethra, N.N., Jayaprasad, K. V. ~~&~~ Kale, R. D. (1999). China aster [*Callistephus chinensis* (L)] cultivation using vermicompost as organic amendment. *Crop Research* (Hisar), 17, 209-215.
- Nisha, R., Kiran, B., Kaushik, A. ~~&~~ Kaushik, C.P. (2018). Bioremediation of salt affected soils using cyanobacteria in terms of physical structures, nutrient status and microbial activity. *Int. ernational J. ournal of Environ. mental Sci. ence and Tech. nology*, 15, 571–580.
- Qadir, M., Schubert, S., Ghafoor, A. ~~&~~ Murtaza, G. (2001). Amelioration strategies for sodic soils: A review. *Land Degradation and Development*, 12, 357-386.
- Seenivasan, R., Prasath, V. ~~&~~ Mohanraj, R. (2015). Restoration of sodic soils involving chemical and biological amendments and phytoremediation by *Eucalyptus camaldulensis* in a semi-arid region. *Environ. mental Geo. chemistry and Health*, 37, 575–586.
- Sharma, R. C. ~~&~~ Banik, P. (2014). Vermicompost and fertilizer application: Effect on productivity and profitability of baby corn (*Zea Mays* L.) and soil health. *Compost Science and Utilization*, 22, 83-92.
- Singh, N.T., Hira, G.S. ~~&~~ Bajwa, M.S. (1981). Use of amendments in reclamation of alkali soils in India. *Agrokemiaes Talajtan Tom. Supplementum*, 30, 159-177.
- Singh, R., Singh, Y.P., Yaduvanshi, N.P.S ~~&~~ Sharma, D.K. (2009). eEffect of irrigation scheduling and integrated nutrient management on yield of rice-wheat system and properties of a reclaimed sodic soil. *J. ournal of the Indian Society of Soil Science*, 57, 280-286.
- Tiwari, R. C., Sharma, P. K. ~~&~~ Khandelwal, S. K. (2004). Effect of green-manuring through *Sesbaniacannabina* and *Sesbaniastrata* and nitrogen application through urea to maize (*Zea mays*) in maize-wheat (*Triticumaestivum*) cropping system. *Indian J. ournal of Agronomy*, 49, 15-17.
- Tripathi N., Singh R.S. ~~&~~ Hills C.D. (2016). Soil carbon development in rejuvenated Indian coal mine spoil. *Ecological Engineering*, 90, 482.
- Xu C., Pu L., Li J., ~~and~~ Zhu M. (2019). Effect of reclamation on C, N, and P stoichiometry in soil and soil aggregates of a coastal wetland in eastern China. *J. ournal of Soils and Sediments*, 19, 1215.

Zakir, H. M., Sultana, M. N. & Saha, K. C. (2012). Influence of commercially available organic vs inorganic fertilizers on growth yield and quality of carrot. *Journal of Environmental Science and Natural Resources*, 5, 39-45.

UNDER PEER REVIEW