

Teasing-Induced Self-Circumcision in a Teenager: A Case Report

ABSTRACT

Aim: Male circumcision in traditionally circumcising communities is often performed by an experienced traditionalist or a surgeon. Transitioning through teenage years as an uncircumcised male in these communities can be challenging. This report highlights the psychosocial pressures of uncircumcised teenagers and the need for early formal circumcision to prevent self-injurious behaviour.

Case Presentation: We present a 13-year-old male, who performed a self-circumcision using a kitchen knife after enduring repeated teasing episodes at annual sports camps. He was resuscitated and later had penile skin reconstruction done. He did not have an underlying psychological disorder.

Discussion: Self-circumcision is rare and mostly reported amongst patients with underlying psychological disorders. Cases amongst teenagers with no underlying pathologies are ever rarer. Negative peer pressure in the form of teasing and push such patients to self-injurious behaviour. Early appropriate surgical intervention following self-circumcision results in good clinical outcomes.

Conclusion: Uncircumcised teenagers in traditionally circumcising communities may suffer adverse psychological trauma from peers leading to self-harm. Parents of such children should seek early medical care to avert the risks of self-injurious behaviour.

Keywords: Self-circumcision, teasing, teenager

1. INTRODUCTION

Male circumcision is an age-old practice and performed for religious, cultural, medical and social reasons¹. Though the number of circumcised males varies considerably for each country and territory, the global prevalence is estimated to be 38.7%². Ghana is estimated to have a country-specific male circumcision prevalence of 91.6%.²

Male circumcision has in the past been performed by traditional ascribers to modern surgeons but not the individuals on themselves.³⁻⁵

A few cases of male circumcision using devices marketed over the counter or internet have been reported.^{3,4,6,7} Even rarer are reports of self-circumcisions using basic tools such as knives and razor blades.³

In many African countries where male circumcision is performed on adolescents/teenagers, it is done as a rite of passage into manhood.^{8,9}

Circumcising oneself due to pressure from peers may be a trigger to pay attention to the mental health of uncircumcised adolescents/teenagers living in traditionally poor-

circumcising societies. It is important to rule out possible underlying psychiatric causes and manage them in tandem with the complications of self-circumcision.¹⁰ This case report highlights the dangers of peer pressure and the potential to induce self-harm in the form of self-circumcision.

2. CASE PRESENTATION

The patient is a 13-year-old male athlete, who presented to the hospital 4 hours following self-circumcision (Figure 1). He complained of severe penile pain and bleeding. The patient admitted to using a home knife in an attempted circumcision.

The patient narrates he has had several teasing episodes at sports camps concerning his uncircumcised phallus. His self-circumcision was an attempt to prevent further teasing at an upcoming sports camp.

He has no previous reports of abnormal behaviour nor a family history of a psychiatric illness. He lives with both parents and is a junior high school student.

On examination, he was a young male, stable. No abnormalities were detected on systemic examination.

The initial mental assessment showed the patient had a good memory and judgement. He had no evidence of psychosis.

Status localis – He had approximately 5cm of circumferential denuded penile skin, with minimal bleeding from the raw areas.



Figure 1 – Penis after self-circumcision

2.1 Interventions.

The patient was admitted and given intravenous analgesics and antibiotics. Tetanus prophylaxis was given, and the penile wound **was** dressed in sterile gauze. An appropriately sized urethral catheter was passed.

Informed consent was obtained from the patient and parents. An assent form was signed by the parents. Permission was sought and obtained from parents for clinical photography and the use of clinical details for publication. The patient was optimized for surgical reconstruction the following day. Under spinal anaesthesia, the wound was cleaned, and debridement **was** done. Both outer and inner skin layers were cut from the attempted circumcision. The retracted skin was advanced to the corona with minimal tension. The skin was sutured using 2.0 interrupted Vicryl. (Figure 2). On the post-operative day (POD)-1, the

urethral catheter was removed, and the patient was attended to by clinical psychologists. On POD-2, the patient had a psychological assessment and was not found to have any underlying psychological disorders. He was discharged home with a clear long-term follow-up plan, with urology and the clinical psychologists. On his last review 3 months following surgery, the patient is doing well and has no complications.



Figure 2 – Penis following reconstruction.

3. DISCUSSION

There are varied reasons behind self-circumcision.¹¹ Feelings of shame, rejection or feeling odd amongst peers, have been reported.¹¹ Some authors have suggested an obsessive-compulsive behavior can be precipitated by the stigma of not being circumcised.³ Nagy and colleagues reported a sophisticated self-circumcision case due to the patients dissatisfaction with his redundant foreskin.¹² Our patient admitted to the prolonged teasing from peers as the trigger to his attempt at self-circumcision. Though there was no family

history or clinical evidence of psychiatric disorder at the time of presentation, it is necessary to have routine mental state assessment by a trained psychologist/psychiatrist. Lwanga et al.³ alluded to the indistinct line between self-circumcision as a form of self-treatment and self-genital mutilation. The former adopts simpler techniques and is organized, the latter is random, radical, and common in patients with psychosocial disorders or overt psychosis.^{3,10} Our patient did not have an underlying psychological disorder neither did the patient in the case reported by Nagy et al.¹² Various sharp objects are employed in cases of self-circumcisions and the use of households knives is not uncommon.¹³ Hemorrhage, urethral injury, extensive skin denudation, infection, delayed healing and **chordee** are all common documented complications associated with self-circumcision.^{3,8,14} The more severe complication of irreversible penile mutilation or amputation, penile necrosis and urethra-cutaneous fistulas are rare and often require **sub-specialty** referral.^{1,14,15} Our patient had a 5cm denudation of the penile skin. The proximal skin was adequate to cover the defect without tension. The urethra was intact, and the patient had not developed a urethro-cutaneous fistula. The defect was reconstructed by a general surgeon and the patient did not require a sub-specialty referral.

4. CONCLUSION

There is an emergence of cases of self-circumcision in circumcision-dominated regions. Several triggers to self-harm have been identified and teenagers may be at a particular risk from negative pressure. Parents and caretakers must be aware of the stigma attached to uncircumcision and readily support their children to access care.

Long-term psychological counselling and follow-up are necessary to unearth sub-clinical diseases and prevent further self-injury.

Ethical Approval:

As per international standard or university standards written ethical approval has been collected and preserved by the author(s).

Consent

As per international standards, parental written consent has been collected and preserved by the author(s).

REFERENCES

1. Natali A, Rossetti MA. Complications of Self- circumcision: A Case Report and Proposal. J Sex Med [Internet]. 2008 Dec 1 [cited 2023 Sep 30];5(12):2970–2. Available from: <https://doi.org/10.1111/j.1743-6109.2008.00858.x>

2. Morris BJ, Wamai RG, Henebeng EB, Tobian AA, Klausner JD, Banerjee J, et al. Estimation of country-specific and global prevalence of male circumcision. *Popul Health Metr* [Internet]. 2016 Mar 1 [cited 2023 Sep 30];14:4. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4772313/>
3. Kintu-Luwaga R. The emerging trend of self-circumcision and the need to define cause: Case report of a 21 year-old male. *Int J Surg Case Rep* [Internet]. 2016 Jun 29 [cited 2023 Sep 30];25:225–8. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4941114/>
4. Dunsmuir WD, Gordon EM. The history of circumcision. *BJU Int*. 1999 Jan;83 Suppl 1:1–12.
5. Adu-Gyamfi S, Adjei POW. Twentieth Century Wanzams among the Asante People of Ghana: A Historical Study of the Facts on Male Circumcision. *Open J Prev Med* [Internet]. 2014 Sep 15 [cited 2023 Sep 30];04(09):730. Available from: <http://www.scirp.org/journal/PaperInformation.aspx?PaperID=50154&#abstract>
6. Koo VSW, Lynn NNK. Infected circumcision ring: a trend of self-circumcision practice? *Surg J R Coll Surg Edinb Irel*. 2011 Aug;9(4):233–4.
7. Kim SJ, Chung H, Ahn HS, Chung DY, Kim YS. Strangulation of the penis by a self-circumcision device. *Urol Int*. 2002;68(3):197–8.
8. Wilcken A, Keil T, Dick B. Traditional male circumcision in eastern and southern Africa: a systematic review of prevalence and complications. *Bull World Health Organ* [Internet]. 2010 Dec 1 [cited 2023 Oct 2];88(12):907–14. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2995181/>
9. Dobanovacki D, Lucić Prostran B, Sarac D, Antić J, Petković M, Lakić T. [Prepuce in boys and adolescents: what when, and how?]. *Med Pregl*. 2012;65(7–8):295–300.
10. Gudugbe S, Asiedu I, Lamptey J, Kyei MY, Baidoo K. Self-Genital Mutilation and Attempted Suicide by Cut Throat in the Same Patient at Presentation: A Rare Event. *Open J Psychiatry* [Internet]. 2015 Sep 29 [cited 2023 Oct 5];5(4):330–3. Available from: <https://www.scirp.org/journal/paperinformation.aspx?paperid=60191>
11. Koç: A novel complication of self-circumcision:... - Google Scholar [Internet]. [cited 2023 Oct 2]. Available from: https://scholar.google.com/scholar_lookup?journal=Turk.+J.+Urol.&title=A+novel+complication+of+self-circumcision:+penile+skin+avulsion&author=Ko%C3%A7+Akif&volume=37&issue=2&publication_year=2011&pages=177-179&
12. Nagy V. Sophisticated self- circumcision: Case report. *Clin Case Rep* [Internet]. 2020 Oct 6 [cited 2023 Sep 4];8(12):3213–5. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7752402/>
13. A E, Y B, Em W, M D, A D, R A. Foreskin necrosis - Complication following self-circumcision. *Urol Case Rep* [Internet]. 2021 Apr 2 [cited 2023 Dec 13];38. Available from: <https://pubmed.ncbi.nlm.nih.gov/33898266/>

14. Sagar J, Sagar B, Shah DK. Penile skin necrosis - complication following selfcircumcision. Ann R Coll Surg Engl [Internet]. 2005 Jul [cited 2023 Oct 2];87(4):W5–7. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1963940/>
15. Krill AJ, Palmer LS, Palmer JS. Complications of Circumcision. Sci World J [Internet]. 2011 Dec 26 [cited 2023 Oct 2];11:2458–68. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3253617/>

UNDER PEER REVIEW