

# Effect of plant spacing and nutrient levels on growth and yield of red cabbage (*Brassica oleracea* var. *capitata* f. *rubra*)

## ABSTRACT

An experiment was conducted to investigate the impact of plant spacing and nutrient levels on the growth and yield of red cabbage. This experiment involved three distinct spacing configurations (45cmx30cm, 45cmx45cm, and 45cmx60cm) and nutrient levels (75% NPK, 100% NPK, and 125% NPK). The experiment was designed as a Factorial Randomized Block Design (FRCBD) and carried out at the Department of Horticulture, University of Agricultural Sciences, GKVK, Bengaluru, during the 2021-22 period. This study indicates that spacing and nutrients promote the vegetative growth of plants. The spacing configuration of 45cm X 60cm with 125% NPK resulted in the most significant outcomes, including the highest number of leaves (28.93), maximum plant height (36.71 cm), maximum fresh weight of the head (1421.80 g) and highest head volume (1360.19 cc) because wider spacing and higher nutrient doses led to proper nutrient use and larger heads. Conversely, the spacing configuration of 45cm X 30cm with 75% NPK demonstrated the early head initiation (45.60 days) and the most compact head formation (0.38). The highest yield per hectare (74.43 t) was achieved with the spacing configuration of 45cmx30cm using 125% NPK which improved greater availability of nutrients and, as a result, boosted the vegetative development, which allowed red cabbage heads to store enormous amounts of food.

*Key words: Growth, head diameter, nutrient, red cabbage, spacing, yield*

## 1. INTRODUCTION

The valuable decorative plant, red cabbage (*Brassica oleracea* var. *capitata* f. *rubra*) is a ~~very healthy~~ highly nutritious vegetable. Red cabbage, ~~belonging to the~~ is a member of the cabbage subgroup *Rubra*, ~~and~~ is a member of the Brassicaceae family. It carries the 2n=18 chromosome. It is also known as purple cabbage or crimson kraut. Its juice is used to be a treatment for poisonous mushrooms. Chemicals in red cabbage serve to regulate angiogenesis and shield DNA from oxidative damage. These pathways assist in treating neoplastic sickness while angiogenesis is suppressed alone to lessen tumor formation, Hagivara *et al.*, [4]. ~~Red cabbage contains compounds that are anti-inflammatory and help pancreatic cells excrete insulin. It is renowned for its medicinal properties as well. It has anticancer properties since indole-3-carbinol is present. It makes a distinction. Red cabbage is known for its medicinal properties, including anti-inflammatory compounds that assist pancreatic cells in insulin secretion. Furthermore, it possesses anticancer properties, attributed to the presence of indole-3-carbinol, highlighting its significant health benefits.~~

Among the various factors that contribute to red cabbage's potential yield, spacing is the most crucial one. ~~The yield is significantly impacted by maintaining the optimal plant population per square foot. Maintaining the optimal plant population per square foot significantly impacts yield.~~ Plant densities that are either too high or too low per unit of area have an effect on crop yields. ~~There has been a rise in interest in using close plant spacing and short rows to produce cabbage in recent years. In recent years, there has been a growing interest in utilizing close plant spacing and short rows for cabbage cultivation.~~ By altering inter and intra row spacings, several workers observed a greater production in crops like broccoli (Agarkar *et al.* [2]).<sup>1</sup>

The growth and development of plants depend on the three main primary plant nutrients, nitrogen, phosphorus, and potassium. ~~Both the quantity and method of fertiliser application must be taken into account in order to increase nutrient utilization efficiency.~~ To enhance nutrient utilization efficiency, it is essential to consider both the amount of fertilizer applied and the method of its application. Split fertiliser applications, which break up total fertiliser treatments into several dosages based on crop requirements, are a critical part of a nutrient management system. This method promotes optimal yields, improves nutrient utilization, and reduces losses. In addition to lowering production, unbalanced fertilizer application harms the soil health. Due to this unbalance nutrient utilization, there is a significant lag between the removal of crops and the application of fertilizer. In India, balanced NPK fertilization has created a lot of interest (Ghosh *et al.*, [3]). ~~In order to~~ To increase and maintain production, it is ~~required-necessary~~ to apply fertilizer components, especially N, P, and K through inorganic sources in the proper quantities.

Nutrition has been discovered to have a significant impact on the growth, yield, quality, and economics of cole crops among other agronomic methods. Two barriers to raising the production of these crops are their unbalanced use and the rising cost of chemical fertilisers. Efficiency in fertiliser utilisation improves yield, preserves soil health, and lowers cultivation costs.

## 2. MATERIAL AND METHODS

Experiment was conducted at Department of Horticulture, College of Agriculture, GKVK, Bengaluru during year 2021-2022. Geographically place is located in Eastern Dry Zone (Zone-5) of Karnataka state at 12° 58" at north latitude and 77° 35" East longitude with an elevation of about 830 meters above mean sea level. The soil is red sandy loam and well drained with uniform texture. The experiment was laid out in factorial randomized block design (FRCBD) with three replications, considering spacing as first factor and different nutrient levels as second factor. Spacing levels ~~are were~~  $S_1$  (45cmx30cm),  $S_2$  (45cmx45cm) and  $S_3$  (45cmx60cm). ~~N~~ Nutrient levels ~~are were~~  $N_1$  (75% NPK),  $N_2$  (100% NPK) and  $N_3$  (125% NPK). Total 9 treatments ~~are were~~  $S_1N_1$ ,  $S_1N_2$ ,  $S_1N_3$ ,  $S_2N_1$ ,  $S_2N_2$ ,  $S_2N_3$ ,  $S_3N_1$ ,  $S_3N_2$  and  $S_3N_3$ . Nitrogen was applied in 2 splits in the form of urea as per the treatment. Half ~~dose~~ of the total nitrogen and full dose of phosphorus and potassium was applied as basal in the form of Diammonium phosphate (DAP) and Muriate of potash (MOP) respectively, at the time of field preparation along with farm yard manure (FYM). Harvesting was done in the morning hours when red cabbage heads were at solidity, ~~that is~~ at the full mature stage, and marketable size. Five plants in each plot were selected randomly and the data were averaged and expressed per plant from the net plot of each replication in each treatment. Observations were recorded on growth parameters like plant height, plant spread, number of leaves per plant, number of days taken for head initiation and days to harvest. ~~The~~ yield parameters ~~recorded were~~ head volume, head circumference, head weight, head compactness, head diameter and head height. Yield per plot and ~~Yield-yield~~ per hectare, were ~~also~~ recorded. Head volume was calculated by using the formula :

$$\text{Head volume} = \frac{4}{3} \times \pi \times \left(\frac{1}{2}MD\right)^3$$

MD = Mean diameter calculated from head polar (PD) and equatorial diameter (ED)

$$MD = (PD + ED) \div 2$$

Head compactness was calculated using the following formula:

$$\text{Compactness rate} = \frac{\text{head volume} \left(\frac{3}{4} \text{radius}^3\right)}{\text{head weight (g)}}$$

### 2.1 Soil analysis

Soil samples were taken down to a depth of 0–15 cm. Composite soil samples were taken from each plot (three replications) at a depth of 0–15 cm prior to the start and end of the studies. For every plot, two sets of sub-samples were created from the three auger samples that were obtained. For physical and chemical tests, the collected materials were air dried, powdered, and placed in a clean plastic container after passing through a 2 mm (10 mesh) screen. Next, a soil sample was taken from every

plot following the cabbage harvest. A composite soil sample was created based on treatment after all debris had been removed, and the soil was given the designation post soil. In the lab, the soil was air dried at ambient temperature. The initial soil's physical and chemical characteristics (Table 1) were then examined using the flame photometer, Brays No. 1 method, and Kjeldahl ~~titration~~ [titration](#) method as standard techniques.

**Table 01. Initial soil properties of the experimental site**

Chemical properties	
Parameter	Value
Ph	6.2-6.4
EC (ds/m)	0.67
N (kg/ha)	302.05
P <sub>2</sub> O <sub>5</sub> (kg/ha)	80
K <sub>2</sub> O (kg/ha)	280.30

Nitrogen uptake (kg/ha) by the plants estimated by micro- Kjeldahl method, Phosphorus ~~will be~~ [was](#) analyzed by Vanado- molybdo phosphoric acid yellow colour method as described by Jackson (1973) [7] and potassium was determined by using Systronics flame photometer.

## 2.2 Benefit: cost ratio

It was obtained by dividing gross returns with cost of cultivation/ha.

$$B:C \text{ ratio} = \frac{\text{Gross return}}{\text{cost of cultivation}}$$

The data obtained from this investigation were appropriately computed, tabulated and analysed using Factorial Randomized Block Design. The statistical analysis of data was done by using OPSTAT online statistical analysis software.

## 3. RESULTS AND DISCUSSION

### 3.1 Growth parameters

Plant spacing and nutrient levels showed significant results on growth parameters (Table 2). Maximum plant height (21.18 cm, 25.90 cm and 33.47 cm) and Highest plant spread (23.02 cm, 49.71 cm and 62.64 cm) was observed in spacing 45 cm x 60 cm at 30, 60 DAT and at harvest respectively, in comparison with spacing 45 cm x 30 cm. More number of leaves was recorded in spacing 45 cm x 60 cm (13.51, 20.76 and 25.43) at 30, 60 DAT and at harvest respectively. Whereas, ~~less~~ ~~minimum~~ number of leaves ((11.98, 16.79 and 21.99) was observed under spacing level 45 cm x 30 cm at 30, 60 DAT and at harvest respectively. These findings matched [with the results these](#) of Sarker *et al.* [12] and Haque *et al.* [6]. This may be due to the fact that when plants are separated from one another by a greater distance, they compete less ferociously for resources and sunlight. Due to the additional sunlight and space provided by wider spacing, the crop may have produced highest plant height, plant spread and more number leaves per plant. Growth parameters were also increased significantly due to nutrient levels. Maximum plant height (21.55 cm, 26.54 cm and 34.56 cm) and plant spread (22.22 cm, 50.29 cm and 63.28 cm at 30, 60 DAT and at harvest respectively) was recorded in 125% NPK. Whereas, 75% of NPK recorded minimum plant height and plant spread. Maximum number of leaves (13.87, 21.18 and 26.06 at 30, 60 DAT and at harvest respectively) were observed in 125% NPK and minimum number of leaves (11.93, 16.71 and 21.65 at 30, 60 DAT and at harvest respectively) were recorded in 75% NPK. Similar results were found in Singh *et al.* [16] and Haque *et al.* [5]. That might be due to higher nutritional levels because more

nutrients are available for growth and development, which causes higher nutrient uptake, and more growth. Interaction effect of wider plant spacing and higher nutrients levels ( $S_3N_3$ ) showed significantly higher plant height, plant spread and number of leaves per plant as compare to other treatments. It was caused by the plants widest spacing receiving an adequate amount of nutrients.

Number of days from transplanting to head initiation was significantly affected by plant spacing and nutrient levels (Table 3). Early head initiation (48.40 days) and Early harvest (83.09 days) was noticed in 45cmx30cm, as compared to 45cmx60cm (51.67 days and 86.44 days, head initiation and harvest respectively). These findings are agreed with Silata *et al.* [15] and Thirupa *et al.* [18]. Higher plant spacing led to a longer harvesting period of days. Increased photosynthesis and dry matter digestion caused by more leaves and increased food availability lengthened the vegetative phase and delayed the onset of the reproductive phase. In case of nutrient levels 75% NPK showed early head initiation and early harvest (47.25 days and 83.09 days respectively). Whereas, late head initiation (52.89 days) and late harvest (86.76 days) were recorded in 125% NPK. Similar results were found in Manasa *et al.* [10] and Yadav *et al.* [20]. It might be because there are more nutrients available, which hastened vegetative growth and suppressed the generative phase. The significant variation was recorded due to combined effect of nutrient and plant spacing too. The early head initiation and harvest was recorded in treatment  $S_1N_1$ .

**Table 2: Effect of plant spacing and nutrient levels on plant height (cm), number of leaves per plant and plant spread of red cabbage**

Treatment	Plant height (cm)			Number of leaves			Plant spread (cm)		
	30 DAT	60 DAT	At harvest	30 DAT	60 DAT	At harvest	30 DAT	60 DAT	At harvest
<b>Spacing (S)</b>									
<b>S<sub>1</sub></b>	16.78	20.28	26.66	11.98	16.79	21.99	17.66	41.07	50.99
<b>S<sub>2</sub></b>	19.83	23.95	31.05	12.96	19.35	24.04	20.84	47.38	59.45
<b>S<sub>3</sub></b>	21.18	25.90	33.47	13.51	20.76	25.43	23.02	49.71	62.64
<b>F - test</b>	*	*	*	*	*	*	*	*	*
<b>S.Em±</b>	0.117	0.155	0.208	0.173	0.136	0.235	0.154	0.141	0.166
<b>CD at 5%</b>	0.353	0.470	0.629	0.523	0.410	0.711	0.464	0.427	0.501
<b>Nutrients (N)</b>									
<b>N<sub>1</sub></b>	17.03	20.69	26.58	11.93	16.71	21.65	18.29	41.87	52.17
<b>N<sub>2</sub></b>	19.21	22.90	30.05	12.64	19.01	23.74	21.02	46.00	57.64
<b>N<sub>3</sub></b>	21.55	26.54	34.56	13.87	21.18	26.06	22.22	50.29	63.28
<b>F - test</b>	*	*	*	*	*	*	*	*	*
<b>S.Em±</b>	0.117	0.155	0.208	0.173	0.136	0.235	0.154	0.141	0.166
<b>CD at 5%</b>	0.353	0.470	0.629	0.523	0.410	0.711	0.464	0.427	0.501
<b>Interaction (SXN)</b>									
<b>S<sub>1</sub>N<sub>1</sub></b>	15.73	18.30	23.10	11.40	15.13	21.00	16.23	38.80	46.97
<b>S<sub>1</sub>N<sub>2</sub></b>	16.32	19.24	25.06	12.13	17.23	22.03	18.83	40.53	50.47
<b>S<sub>1</sub>N<sub>3</sub></b>	18.30	23.30	31.83	12.40	18.00	22.93	17.93	43.87	55.53
<b>S<sub>2</sub>N<sub>1</sub></b>	17.14	20.57	25.83	12.20	17.73	21.73	17.50	41.67	52.40
<b>S<sub>2</sub>N<sub>2</sub></b>	20.07	24.13	32.20	12.47	18.13	24.07	21.73	48.20	59.87

<b>S<sub>2</sub>N<sub>3</sub></b>	22.29	27.15	35.13	14.20	22.20	26.33	23.30	52.27	66.07
<b>S<sub>3</sub>N<sub>1</sub></b>	18.23	23.20	30.80	12.20	17.27	22.23	21.13	45.13	57.13
<b>S<sub>3</sub>N<sub>2</sub></b>	21.23	25.33	32.90	13.33	21.67	25.13	22.50	49.27	62.57
<b>S<sub>3</sub>N<sub>3</sub></b>	24.07	29.17	36.71	15.00	23.33	28.93	25.43	54.73	68.23
<b>F - test</b>	*	*	*	*	*	*	*	*	*
<b>S.Em±</b>	0.202	0.269	0.361	0.300	0.235	0.408	0.266	0.244	0.287
<b>CD at 5%</b>	0.611	0.814	1.090	0.906	0.710	1.232	0.804	0.739	0.867

S<sub>1</sub> = 45 cm x 30 cm, S<sub>2</sub> = 45 cm x 45 cm, S<sub>3</sub> = 45 cm x 60 cm, N<sub>1</sub> = 75%NPK, N<sub>2</sub> = 100%NPK, N<sub>3</sub> = 125%NPK

### 3.2Yield and quality parameters

Yield parameters were significantly influenced by plant spacing and application of nutrient levels (Table 3). Maximum head circumference, head diameter and head height (39.15cm, 12.43 cm and 12.82cm respectively) were noticed in 45cmx60cm spacing, while minimum (33.55cm, 11.13cm and 11.47cm head circumference, head diameter and head height respectively) was recorded in 45cmx30cm spacing. These results were agreed with [the findings of Agarkar et al. \[2\]](#) and Shamima *et al.*[13]. Wider spacing gives each plant more room and less competition between plants, that is conducive to the development of a head size. Maximum head circumference (39.72cm), head diameter (12.54cm) and head height (12.96cm) were recorded in 125%NPK. Whereas, 75% NPK showed minimum head circumference (33.66 cm), head diameter (11.23cm) and head height (11.42cm). Similar [results were also observed earlier by findings were found in Yebirzafet al.\[21\]](#) and Manasa *et al.*[10]. Increased nutritional availability may have expedited the production of chlorophyll and amino acids, and efficient use of carbohydrates and their organic components may have led to an increase in head size. The highest head circumference, diameter and height was found in combined effect of 45cmx60cm with 125% NPK. Nutrients and spacing influenced the head size. Similar results were reported by Joshi *et al.*[8].

**Table 3: Effect of plant spacing and nutrient levels on days taken for head initiation, days to harvest, head circumference, head diameter and head height of red cabbage**


Treatment	Days taken for head initiation	Days to harvest	Head circumference (cm)	Head diameter (cm)	Head height (cm)
<b>Spacing (S)</b>					
<b>S<sub>1</sub></b>	48.40	83.09	33.55	11.13	11.47
<b>S<sub>2</sub></b>	49.51	84.09	37.20	11.92	12.17
<b>S<sub>3</sub></b>	51.67	86.44	39.15	12.43	12.82
<b>F -test</b>	*	*	*	*	*
<b>S.Em±</b>	0.134	0.176	0.329	0.072	0.087
<b>CD at 5%</b>	0.406	0.532	0.994	0.217	0.263
<b>Nutrients (N)</b>					
<b>N<sub>1</sub></b>	47.25	83.09	33.66	11.23	11.42
<b>N<sub>2</sub></b>	49.44	83.78	36.52	11.73	12.08
<b>N<sub>3</sub></b>	52.89	86.76	39.72	12.54	12.96
<b>F -test</b>	*	*	*	*	*
<b>S.Em±</b>	0.134	0.176	0.329	0.072	0.087
<b>CD at 5%</b>	0.406	0.532	0.994	0.217	0.263
<b>Interaction (SXN)</b>					
<b>S<sub>1</sub>N<sub>1</sub></b>	45.60	82.13	31.83	10.65	11.09

<b>S<sub>1</sub>N<sub>2</sub></b>	48.27	83.07	32.28	11.07	11.21
<b>S<sub>1</sub>N<sub>3</sub></b>	51.33	84.07	36.53	11.73	12.11
<b>S<sub>2</sub>N<sub>1</sub></b>	47.27	83.07	33.58	11.45	11.39
<b>S<sub>2</sub>N<sub>2</sub></b>	49.13	83.13	37.79	11.95	12.29
<b>S<sub>2</sub>N<sub>3</sub></b>	52.13	86.07	40.23	12.35	12.84
<b>S<sub>3</sub>N<sub>1</sub></b>	48.87	84.07	35.56	11.58	11.79
<b>S<sub>3</sub>N<sub>2</sub></b>	50.93	85.13	39.48	12.18	12.75
<b>S<sub>3</sub>N<sub>3</sub></b>	55.20	90.13	42.41	13.54	13.93
<b>F -test</b>	*	*	*	*	*
<b>S.Em±</b>	0.232	0.305	0.569	0.125	0.151
<b>CD at 5%</b>	0.703	0.922	1.722	0.377	0.456


S<sub>1</sub> = 45 cm x 30 cm, S<sub>2</sub> = 45 cm x 45 cm, S<sub>3</sub> = 45 cm x 60 cm, N<sub>1</sub> = 75%NPK, N<sub>2</sub> = 100%NPK, N<sub>3</sub> =125%NPK

Significant results were observed in head compactness with respect to plant spacing and nutrient levels (Fig. 1). Highest head compactness (0.43) was recorded in wider spacing and lowest (0.48) was observed in closer spacing. Plant spacing has a good effect on head compactness. This may be the result of increased plant competition for nutrients and available space at closer plant spacing, which encouraged the growth of compact heads. Whereas, maximum head compactness (0.43) was noticed in 75% NPK and minimum (0.48) was recorded in 125% NPK. This can be due to the red cabbage heads receiving the ideal amount of nutrients, which led to the heads being more compact. The maximum head compactness (0.38) was observed in interaction effect of 45cmx30cm with 75% NPK. These results were in agreement with Riad *et al.* (2009) in cabbage.

**Fig.1: Effect of plant spacing, nutrient levels and interaction on head compactness of red cabbage**







$S_1$  = 45 cm x 30 cm,  $S_2$  = 45 cm x 45 cm,  $S_3$  = 45 cm x 60 cm,  $N_1$  = 125:37.5:37.5 kg/ha,  $N_2$  = 180:50:50 kg/ha and  $N_3$  = 225:62.5:62.5 kg/ha

Significant variation in head volume was observed in plant spacing and nutrient levels. Highest head volume (1073.17cc) and fresh head weight (1214.91g) were recorded with spacing of 45cmx60cm as compared to spacing 45cmx30cm (Table 4). These findings were agreed with the observations of Abed *et al.* [1]. This could be attributed to the fact that there are fewer plants per square foot, which creates more conducive growing circumstances like more room for shoot, root, and leaf growth than there would be with more tightly spaced plants. With respect to nutrients, maximum head volume (1099.04cc) and fresh weight of head (1256.18g) were recorded in highest nutrient level. Whereas, lowest nutrient level observed minimum head volume and fresh weight of head. These results were conformity with Verma and Nawange [19] and Manasa *et al.* [10]. Increased nutritional availability, which may have led to the production of more plant metabolites. Head volume may have increased because metabolites were more readily available to the plant. Interaction effect of 45cmx60cm spacing with 125% NPK recorded maximum head volume (1360.19) and fresh weight of head (1421.80g). Whereas, minimum was observed in 45x30cm with 75% NPK.

Higher yield per plot (61.72kg) and yield per hectare (69.26t) was recorded in closest spacing and wider spacing observed for lower yield (40.09kg/plot and 45.00t/ha) (Table 4). Similar findings were found in Kaur *et al.* [9] and Silatare *et al.* [15]. The maximum yield was found to be highest at a reduced plant spacing due to having more plants per unit area and a bigger ground cover of leaf area, which improved interception of sunlight and subsequently increased assimilate production. Maximum yield (57.43kg/plot and 64.45t/ha) was observed in 125% NPK whereas, 75% NPK observed for minimum (44.62kg/plot and 56.03t/ha) yield. experimental findings were in consonance with the findings of the Sultana *et al.* [17] and Prasad *et al.* [11]. Because applying nutrients boosted plant vigour in the form of height and leaf count, which in turn increased photosynthetic efficiency and, ultimately, increased crop yield, the importance of nutrients in increasing output was well established. combined effect of closer spacing with higher nutrient level showed highest yield (66.32kg/plot) and (74.43t/ha). These results agreed with Kaur *et al.* [9].

**Table 4: Effect of plant spacing and nutrient levels on head volume (cc) and fresh weight of head (g), yield per plot (kg) and yield per hectare (t) of red cabbage**

Treatment	Head volume (cc)	Fresh weight of head (g)	Yield per plot (kg)	Yield per hectare (ha)
Spacing (S)				
$S_1$	762.47	934.69	61.72	69.26

<b>S<sub>2</sub></b>	922.45	1140.04	50.16	56.30
<b>S<sub>3</sub></b>	1073.17	1214.91	40.09	45.00
<b>F -test</b>	*	*	*	*
<b>S.Em±</b>	15.91	14.17	0.65	0.58
<b>CD at 5%</b>	48.11	42.83	1.97	1.75
<b>Nutrients (N)</b>				
<b>N<sub>1</sub></b>	764.35	951.84	44.62	50.08
<b>N<sub>2</sub></b>	894.69	1081.62	49.92	56.03
<b>N<sub>3</sub></b>	1099.04	1256.18	57.43	64.45
<b>F -test</b>	*	*	*	*
<b>S.Em±</b>	15.91	14.17	0.65	0.58
<b>CD at 5%</b>	48.11	42.83	1.97	1.75
<b>Interaction (SXN)</b>				
<b>S<sub>1</sub>N<sub>1</sub></b>	673.73	872.20	57.63	64.68
<b>S<sub>1</sub>N<sub>2</sub></b>	724.57	927.20	61.20	68.68
<b>S<sub>1</sub>N<sub>3</sub></b>	889.12	1004.67	66.32	74.43
<b>S<sub>2</sub>N<sub>1</sub></b>	782.14	979.53	43.10	48.37
<b>S<sub>2</sub>N<sub>2</sub></b>	937.38	1098.53	48.33	54.25
<b>S<sub>2</sub>N<sub>3</sub></b>	1047.82	1342.07	59.05	66.27
<b>S<sub>3</sub>N<sub>1</sub></b>	837.18	1003.80	33.13	37.18
<b>S<sub>3</sub>N<sub>2</sub></b>	1022.14	1219.13	40.23	45.16
<b>S<sub>3</sub>N<sub>3</sub></b>	1360.19	1421.80	46.92	52.66
<b>F -test</b>	*	*	*	*
<b>S.Em±</b>	27.55	24.53	1.13	1.01
<b>CD at 5%</b>	83.10	74.19	3.41	3.04

S<sub>1</sub> = 45 cm x 30 cm, S<sub>2</sub> = 45 cm x 45 cm, S<sub>3</sub> = 45 cm x 60 cm, N<sub>1</sub> = 75%NPK, N<sub>2</sub> = 100%NPK, N<sub>3</sub> =125%NPK

### 3.4 Soil analysis

Effect of plant spacing, nutrient levels and their interactions significantly influenced on uptake of nitrogen, phosphorus and potassium (Table 5). Highest uptake of nitrogen (330.15 kg/ha), Phosphorus (29.11 kg/ha) and Potassium (166.14 kg/ha) were recorded in the treatment of with S<sub>3</sub> (45 cm x 60 cm)- whereas, N<sub>3</sub> (125% NPK) achieved higher nitrogen (332.04 kg/ha), phosphorus (31.39 kg/ha) and potassium (167.05 kg/ha) uptake by plants. In interaction effect S<sub>3</sub>N<sub>3</sub> (45 cm x 60 cm + 125% NPK) recorded higher uptake of nitrogen (375.93 kg/ha), phosphorus (35.94 kg/ha) and potassium (185.11 kg/ha). The lowest uptake of (179.87kg/ha, 76 kg/ha and 86.60 kg/ha) nitrogen, phosphorus and potassium respectively were recorded with combination of S<sub>1</sub>N<sub>1</sub> (45 cm x 30 cm + 75% NPK).

**Table 5: Effect of different plant spacing and nutrient levels on uptake of nutrients of red cabbage**

Treatment	Nitrogen uptake (kg/ha)	Phosphorus uptake (kg/ha)	Potassium uptake (kg/ha)
<b>Spacing (S)</b>			
<b>S<sub>1</sub></b>	278.55	23.36	112.29
<b>S<sub>2</sub></b>	341.78	26.81	146.99
<b>S<sub>3</sub></b>	375.93	29.11	166.14
<b>F -test</b>	*	*	*



<b>S.Em±</b>	2.44	0.29	1.29
<b>CD at 5%</b>	7.38	0.88	3.90
<b>Nutrients (N)</b>			
<b>N<sub>1</sub></b>	228.27	20.29	112.00
<b>N<sub>2</sub></b>	293.27	27.60	146.36
<b>N<sub>3</sub></b>	332.04	31.39	167.05
<b>F- test</b>	*	*	*
<b>S.Em±</b>	2.44	0.29	1.29
<b>CD at 5%</b>	7.38	0.88	3.90
<b>Interaction (SXN)</b>			
<b>S<sub>1</sub>N<sub>1</sub></b>	179.87	17.76	86.60
<b>S<sub>1</sub>N<sub>2</sub></b>	226.39	25.84	112.45
<b>S<sub>1</sub>N<sub>3</sub></b>	278.55	26.49	137.83
<b>S<sub>2</sub>N<sub>1</sub></b>	234.14	21.17	112.71
<b>S<sub>2</sub>N<sub>2</sub></b>	315.46	27.50	150.03
<b>S<sub>2</sub>N<sub>3</sub></b>	341.78	31.75	178.22
<b>S<sub>3</sub>N<sub>1</sub></b>	276.62	21.94	136.70
<b>S<sub>3</sub>N<sub>2</sub></b>	337.90	29.45	176.60
<b>S<sub>3</sub>N<sub>3</sub></b>	375.93	35.94	185.11
<b>F -test</b>	*	*	*
<b>S.Em±</b>	4.23	0.50	2.24
<b>CD at 5%</b>	12.78	1.52	6.76

S<sub>1</sub> = 45 cm x 30 cm, S<sub>2</sub> = 45 cm x 45 cm, S<sub>3</sub> = 45 cm x 60 cm, N<sub>1</sub> = 75%NPK, N<sub>2</sub> = 100%NPK, N<sub>3</sub> = 125%NPK

Different plantings spacing, nutrient levels and their interaction significantly influenced the available soil nitrogen, phosphorus and potassium (Table 6). Available soil nitrogen (245.23 kg/ha), soil phosphorus (69.49 kg/ha) and soil potassium in soil (131.81 kg/ha) was maximum in S<sub>3</sub> (45 cm x 60 cm). Lowest available soil nitrogen (225.81 kg/ha), soil phosphorus (57.77 kg/ha) and soil potassium (116.77 kg/ha) in soil was recorded with S<sub>1</sub> (45 cm x 30 cm). Significantly higher amount of available soil nitrogen (248.42 kg/ha), soil phosphorus (70.88 kg/ha) and soil potassium (130.86 kg/ha) was recorded with an application of N<sub>3</sub> (125% NPK). Whereas, lowest available soil nitrogen (226.44 kg/ha), soil phosphorus (58.47 kg/ha) and soil potassium (119.14 kg/ha) was noticed in N<sub>1</sub> (75% NPK). The interaction effect of S<sub>3</sub>N<sub>3</sub> (45 cm x 60 cm + 125% NPK) recorded highest available soil nitrogen (252.71 kg/ha), soil phosphorus (75.74 kg/ha) and soil potassium (140.81 kg/ha). Different planting spacing and nutrient levels significantly influenced available soil nitrogen, phosphorus and potassium in red cabbage. These results were in agreement with the findings of Sharma and Arya [14].

**Table 6: Effect of different plant spacing and nutrient levels on available nutrients after harvest of red cabbage**


<b>Treatment</b>	<b>Available Nitrogen (kg/ha)</b>	<b>Available Phosphorus (kg/ha)</b>	<b>Available Potassium (kg/ha)</b>
<b>Spacing (S)</b>			
<b>S<sub>1</sub></b>	225.81	57.77	116.77
<b>S<sub>2</sub></b>	242.74	67.46	124.84
<b>S<sub>3</sub></b>	245.23	69.49	131.81
<b>F -test</b>	*	*	*
<b>S.Em±</b>	1.49	0.39	0.67

CD at 5%	4.49	1.19	2.01
<b>Nutrients (N)</b>			
N <sub>1</sub>	226.44	58.47	119.14
N <sub>2</sub>	238.91	65.36	123.42
N <sub>3</sub>	248.42	70.88	130.86
F -test	*	*	*
S.Em±	1.49	0.39	0.67
CD at 5%	4.49	1.19	2.01
<b>Interaction (SXN)</b>			
S <sub>1</sub> N <sub>1</sub>	210.95	52.07	111.34
S <sub>1</sub> N <sub>2</sub>	223.99	57.05	116.41
S <sub>1</sub> N <sub>3</sub>	242.49	64.18	122.57
S <sub>2</sub> N <sub>1</sub>	232.23	59.98	120.59
S <sub>2</sub> N <sub>2</sub>	245.93	69.66	124.73
S <sub>2</sub> N <sub>3</sub>	250.07	72.73	129.19
S <sub>3</sub> N <sub>1</sub>	236.15	63.35	125.48
S <sub>3</sub> N <sub>2</sub>	246.82	69.37	129.13
S <sub>3</sub> N <sub>3</sub>	252.71	75.74	140.81
F -test	*	*	*
S.Em±	2.57	0.68	1.15
CD at 5%	7.78	2.06	3.48

S<sub>1</sub> = 45 cm x 30 cm, S<sub>2</sub> = 45 cm x 45 cm, S<sub>3</sub> = 45 cm x 60 cm, N<sub>1</sub> = 75%NPK, N<sub>2</sub> = 100%NPK, N<sub>3</sub> =125%NPK

In the present study, the maximum gross returns, net returns and B: C ratio (2.17) were realized with S<sub>1</sub> (45 cm x 30 cm) and supplied with N<sub>3</sub> (125% NPK) (Fig. 2). This was mainly due to higher head yield as compared to other plant spacing and nutrient levels.

**Fig.2: Effect of plant spacing, nutrient levels on benefit cost ratio of red cabbage**



S<sub>1</sub> = 45 cm x 30 cm, S<sub>2</sub> = 45 cm x 45 cm, S<sub>3</sub> = 45 cm x 60 cm, N<sub>1</sub> = 125:37.5:37.5 kg/ha, N<sub>2</sub> = 180:50:50 kg/ha and N<sub>3</sub> = 225:62.5:62.5 kg/ha

#### 4. CONCLUSION

According to the experimental results of the study, the wider spacing (45 cm 60 cm) and higher nutrient levels (125% NPK) resulted in improved growth and development of the plants. Closer spacing (45 cm x 30 cm) and lower levels of nutrients (75% NPK) led to an earlier harvest, but less spacing (45 cm x 30 cm) and greater levels of nutrients (125% NPK) produced a larger quantitative yield per unit area. Conversely, closer spacing (45 cm x 30 cm) and lower nutrient levels (75% NPK) led to an earlier harvest. However, less spacing (45 cm x 30 cm) and greater nutrient levels (125% NPK) produced a larger quantitative yield per unit area.

#### REFERENCES

1. Abed MY, El-said MM, Shebl EF. Effect of planting date and spacing on yield and quality of cabbage (*Brassica oleracea* var. *capitata* L.). J Plant Prod Sci. 2015;6(12):2093-2102.
2. Agarkar UR, Dadmal KD, Nikas NS, Piwlatkar GK. Effect of nitrogen levels and spacing on growth and yield of broccoli (*Brassica oleracea* var. *italica* L.). Green Farming. 2010;1(5):477-479.
3. Ghosh PK, Ramesh P, Bandyopadhyay KK, Tripathi AK, Hati KM, Misra AK, Acharya CL. Comparative effectiveness of cattle manure, poultry manure, phosphocompost and fertilizer-NPK on three cropping systems in vertisols of semi-arid tropics. I. Crop yields and system performance. Bioresour Technol. 2004;95(1):77-83.
4. Hagivara A, Yoshiho H, Ichihara T, Kawabe M, Tamano S, Aoki H, et al. Prevention by natural food anthocyanins, purple sweet potato color and red cabbage color, of 2-amino-1-methyl-6-phenylimidazo pyridine – associated colorectal carcinogenesis in rats initiated with 1,2-dimethylhydrazine. J Toxicol Sci. 2002;27(1):57-68.
5. Haque FA, Islam N, Islam MN, Ullah A, Sarkar MD. Growth, yield and profitability of cabbage (*Brassica oleracea* L.) as influenced by applied nitrogen and plant spacing. The Agric. 2015;13(1):35-45.
6. Haque KMF, Jahangir A A, Haque ME, Mondal RK, Jahan MAA, Sarker MAM. Yield and nutritional quality of cabbage as affected by nitrogen and phosphorus fertilization. Bangladesh J Sci Ind Res. 2006;41(1):41-46.
7. Jackson ML. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd. 1973; New Delhi, 498.
8. Joshi TN, Budha CB, Sharma S, Baral SR, Pandey NL, Rajbhandari R. Effect of different plant spacing on the production of hybrid cauliflower (*Brassica oleracea* var. *botrytis*) under the agro-climatic conditions of mid-hills Region of Nepal. J Plant Prot Res. 2018;1(1):105.
9. Kaur P, Singh S, Sidhu MK. Response of different levels of nitrogen and spacing response of different levels of nitrogen and spacing on yield and quality of cauliflower grown under central region of Punjab. Bioscan. 2020;15(1):123-128.
10. Manasa S, Mukunda L, Sadarunnisa S, Rajasekharam T. Studies on effect of spacing on yield and yield attributing parameters of red cabbage (*Brassica oleracea* var. *capitata* f. *rubra*). Int J Curr Microbiol Appl Sci. 2017;6(12):3143-3147.
11. Prasad PH, Bhunia P, Naik A, Thapa U. Response of nitrogen and phosphorus levels on the growth and yield of chinese cabbage (*Brassica campestris* L. var. *pekinensis*) in the gangetic plains of West Bengal. J crop weed. 2009;5(2):75-77.
12. Sarker MY, Hasan AK, Nasreen A, Naher Q, Baset MA. Effect of plant spacing and sources of nutrients on the growth and yield of cabbage. Pak J Biol Sci. 2002;5(6):636-639.
13. Shamima K, Haque MA, Ahmed F, Sultana F, Ali MY. Effect of plant spacing and potassium on growth and yield of cabbage. J Agrofor Environ. 2018;12(1-2):93-96.
14. Sharma KC, Arya PS. Effect of nitrogen and farmyard manure on cabbage (*Brassica oleracea* var. *capitata*) in dry temperate zone of Himachal Pradesh. Indian J Agric Sci. 2001;71(1):60 – 61.

15. Silatar P, Patel GS, Acharya SK, Vadodaria JR. Performance of different varieties and plant spacing on growth and yield of knolkhol (*Brassica oleracea* var. *gongylodes*). Int J Agric Sci. 2018;8(7):1476-1479.
16. Singh MK, Chand T, Kumar M, Singh KV, Lodhi SK, Singh VP, Sirohi VS. Response of different doses of NPK and boron on growth and yield of broccoli (*Brassica oleracea* L. var. *italica*). I J Bio-resource Stress Manag. 2015;6(1):108-112.
17. Sultana J, Siddique MA, Rashid MHA. Effects of cowdung and potassium on growth and yield of kohlrabi. J Bangladesh Agril Univ. 2012;10(1):27-32.
18. Thirupal D, Madhumathi C, Syamsundar RP. Effect of planting dates and plant spacing on growth, yield and quality of broccoli under Rayalaseema zone of Andhra Pradesh, India. Plant Arch. 2014;14(2):1095-1098.
19. Verma H, Nawange DD. Effect of different levels of nitrogen and sulphur on the growth, yield and quality of cabbage (*Brassica oleracea* var. *capitata* L.). Agric Sci Dig. 2015;35(2):152-154.
20. Yadav LP, Kavita A, Maurya IB. Effect of nitrogen and biofertilizers on growth of cabbage (*Brassica oleracea* var. *capitata* L.) var. Pride of India. Progressive Hortic. 2012;44(2):318-320.
21. Yebirzaf Y. Effect of different rate of nitrogen fertilizer on the growth and yield of cabbage (*Brassica oleraceae*) at Debre Markos, North West Ethiopia. S Afr J Plant Soil. 2017;11(7):276-281.