

Studies on seasonal variation of zooplankton population in Elanthakulam , Palayamkottai, Tirunelveli District

Abstract

In Elanthakulam Pond in the Tirunelveli District, a study of the zooplankton population was carried out, and plankton samples were taken from August to October of 2023. Abiotic and biotic variables have worked together to produce seasonal shifts in the zooplankton population pattern. In order to evaluate the zooplankton population in the Elanthakulam pond, the current study was conducted. The findings show that during the research period, 10 species from four genera were identified. Of the ten species, four belonged to the rotifer genus, three to the Cladoceran genus, two to the copepod genus, and one to the Ostracoda genus. The presnet study suggests that due to unfavourable environmental conditions, zooplankton did not reach its peak during the monsoon season. It also shows a minor fall in August and September, followed by an increase in October.

Comment [W1P1]: biotic

Comment [W1P2]: write any methods before the results

Keywords: Elanthakulam, Cladiceran, Moina, Daphnia, Ostracoda

Introduction:

According to Ali (1999), biodiversity can also be defined as the variation found in all living things, including those found in terrestrial, marine, and other aquatic ecosystems, as well as the ecological complexes of which they are a part. Details regarding the evenness, dominance, richness, and variety of species. Analysing the ecosystem's biological components is crucial to comprehending harmful environmental changes (Krishnamoorthy and Subramanian, 1999). Major taxonomic groups are represented in the rich array of zooplankton found in Indian water bodies. Numerous of these types have distinct physiological and environmental assemblages. Any aquatic habitat's population size, composition, and distribution can reveal information about the environmental conditions that are present there.

It is evident that a variety of environmental elements combine to create the right conditions for zooplankton growth in both seasonal and geographical contexts (Shah and Pandit 2013). An essential component of an aquatic ecosystem's relevance and a major player in energy transmission is zooplankton. Because they are highly sensitive to their surroundings, plankton populations' tolerance, abundance, variety, and dominance in the habitat will all shift in response to environmental changes. As a result, plankton population observation could be a trustworthy method for biomonitoring research that evaluates the level of pollution in aquatic environments. The ecosystem and food chain of ponds, lakes, and reservoirs depend heavily on freshwater zooplankton (Manicam et al., 2014). Zooplankton consume phytoplankton as food. They are in control of consuming millions of tiny algae that would otherwise spread uncontrollably. Different zooplankton species have distinct life histories that are impacted by predation pressure, feeding ecology, and seasonal changes in biotic variables. Primary consumers, which consume phytoplankton, and secondary consumers, which eat other zooplankton, make up the zooplankton community. They offer a direct conduit between upper trophic levels, such fish, and primary producers. During their larval stages, almost all fish rely on zooplankton as their primary food source, and some fish consume it for the entirety of their lives (Madin et al., 2001). The freshwater zooplankton population is essential to the food chain's food web because it recycles nutrients and moves organic matter from primary producers like diatoms to secondary consumers like fish. The amount of fish stock is determined in part by zooplankton, and the decline in the Copepod population is thought to be the cause of the fishing resources' failure (Stottrup 2000 and Tiwari et al., 1991). The water quality is

assessed in terms of domestic, municipal, and industrial contamination using species diversity indices of zooplankton communities (Acharjee et al., 1995). Thus, zooplankton can serve as a sorority indicator. For the purpose of managing zooplankton populations and maximising system production, research on their variety, density, commonness, and energy levels is currently necessary. When taking into account an aquatic system, planktons show to be quite significant since they can respond instantly to changes in the surrounding environment (Thakur et al., 2013 and Malik et al., 2013). Many biotic and abiotic elements, including as light, temperature, available nutrients, hydrodynamics, predation, oxygen content, pH, and so forth, affect the growth and development rate of plankton (Dhar et al., 2012). The trophic level that comes after the phytoplanktons is called the zooplanktons (Malik et al., 2013 and Shanthala et al., 2008). The physicochemical characteristics of the water body have an impact on the zooplankton population, which also varies with the seasons (Hulyal and Kaliwal 2007 and Kudari and Kanamadi 2007). Unlike physicochemical methods that have led to the detection of one pollutant at a time, the analysis of such indicator organisms, both qualitative and quantitative, has resulted in an assisting option to combine the effects of a number of contaminants. Furthermore, the current state of many water bodies has been ascertained by using indices and other technologies. According to Mahadev et al. (2007), biomonitoring has emerged as a crucial component of studies on water pollution and makes a significant contribution to the field's understanding of water quality assessment. The physicochemical parameters of the environment cause fluctuations in the zooplankton population; in particular, rotifer species are affected by biotic variables (Karuthapandi et al., 2012). The dominance of fish, macroinvertebrates, and water fowl, as well as their feeding preferences, determine the amount and composition of zooplanktons (Russell et al., 2006). In a similar vein, Jafari et al. (2011) investigated the relationship between the physicochemical conditions of the Haraz River and the variety and compositions of zooplankton. Due to their heterotrophic nature, zooplanktons connect primary producers to higher trophic levels and play a crucial role in the food web. Because there are less opportunities for an individual in a water body to remain in the eutrophic zone, where photosynthesis takes place, the abundance of zooplankton is declining (Dhembare, 2011). When assessing the temperature, pollution, and nutrient levels of an ecosystem, zooplanktons play a significant role (Purushothama et al.,

2011). Evaluating the zooplankton population in the Eanthakulam pond, Tirunelveli District is the aim of the present study.

Material and Methods:

The plankton sample used in this investigation was collected at Elanthakulam Pond during the winter months of August, September, and October of 2016 (Fig 1a-b). A 25µm-mesh plankton net was swept over surface water, and the plankton it caught was then placed in a different plastic container. To get plankton, around 1 litre of surface water was sieved via a net. Without delay, the zooplankton was preserved in 4% formalin for subsequent microscopic examinations. Post-Clegg (1956), Edmondson (1959), Hutchinson (1967), Michael (1973), Ward and Whipple (1963), Pennak (1978), APHA (1989), and Sridharan (1989) plankters were identified. Useful indices of species organisation in communities, as described by Odum (1971), were also computed when plankters were identified.

Result:

Tables 1 to 5 show the zooplankton that was seen in the Elanthakulam. The tables display that the zooplankton present in the pond belonged to four distinct groups: Ostracoda, Cladocera, Rotifera, and Copepoda. During the study period, a total of 10 species were reported in the Elanthakulam pond. Four rotifer species in total were identified: *Keretella cochlearis*, *Brachionus rubens*, *Brachionus caudatus*, and *Brachionus calyciflorus*. Table 1 lists the different zooplankters that constituted the Rotifera group. The table makes it clear that a total of 4 species from 4 genera were found. Of them, three species belong to the genus *Brachionus*, whereas just one species represents genera like *Keretella*. For *Brachionus calyciflorus* and *B. rubens*, the best counts were recorded in February; for *B. caudatus*, the best counts were recorded in January; and for *Keretella*, the best counts were recorded in February.

Comment [W1P3]: is it pond?
From the picture, it seems like a lake or reservoir

Comment [W1P4]: This study actually mostly related to water quality, especially when conducted during winter season. But no any data presented regarding water quality

Comment [W1P5]: Write the scientific name in italic.
Check all the scientific names in this manuscript

Table 2 lists the different zooplankters that represented the Cladoceran and Anostracan groups. The table makes it clear that a total of 3 species from 2 genera were found. Of these, two species belong to the genus *Moina*, while one species represents genera such as *Daphnia*. *Moina micrura* chose to record her maximum counts in January, while *Daphnia pules* and *Moina brachiata* preferred to record their best counts in February. Table 3 lists the different zooplankton that constitute the copepod group. The table makes it clear that a total of 2 species from 2 genera were found. Of them, one species represented the genus *Diaptamus*, while another species represented genera such as *Mesocyclops hyalinus*. *Mesocyclops hyalinus* favoured January to record their highest counts, while *Diaptamus castor* preferred February. Table 4 showed the zooplankters that constituted the Ostracode group. A total of solitary species from solitary genera were noted. One species from the genus *Cypris* was present among them. The best counts of *Cypris* Ostracodan were recorded in January, and there were less of them available during this season.

Disussion:

While being present all year round, rotifers preferred the months of August through October for recording their counts, according to an overall comparison of the population. A detailed examination indicates that, out of this time frame, December to February seems to be the most preferred month because during these months, four of the species that were noted reached their peak counts. According to research, one of the most prevalent rotifer genera found in the majority of India's water bodies is *Brachionus* (Sreenivasan, 1974; Rajasekhar et al., 2010; Singh et al., 2012; Srivastava, 2013). Therefore, it makes sense that *Brachionus* was the most prevalent genera in this pond during the current investigation as well. It has also been found via numerous researchers' studies that rotifers seem to favour particular months of the year over others in distinct bodies of water. When it came to their preferences, Michael (1969), Chourasia and Adoni (1985), Singh et al. (2012), Tidame and Shinde (2012), Kastoorigbai (1991), and Sivakami (1996) stated that they liked June through August, Jayanthi (1994), and Rajasekhar et al. (2010) stated that they preferred September and October. The present observation is consistent with these reports. It has been suggested by Tidame and Shinde (2012) that rotifers are used as an important aquatic faunal component for biomonitoring, while Bogdan and Gilbert (1984) claim that rotifers are the dominant members of the

zooplankton in most aquatic systems and that almost all fish feed on tiny rotifers during their early development. Sharma (1991) also notes that of the different rotifers that have been identified so far, rotifers belonging to the genus *Brachionus* are more suitable for feeding fish larvae. The most significant soft-bodied metazoans, or invertebrates, with a brief life cycle among plankton are called rotifers. Table 4 displays the different zooplankters that represented the Anostracan and Cladoceran groups. A single species represented the genus *Cypris*. The best counts of *Cypris* ostracodan were recorded in January, and there were less of them available during this season. Numerous studies indicate that Chourasia and Adoni (1985) said they liked October and April, Khan et al. (1986) said they liked summer, and Haque and Khan (1997) said they liked December, May, and August. According to Rajasekhar et al. (2010), they both liked the winter and the rainy season (Tidame and Shinde, 2012). A vital class of zooplankton, cladocerans are the most beneficial and nutrient-dense group of crustaceans for fish further up the food chain. Cladocera constituted the major group within the Zooplankton. *Daphnia* sp., *Moina* sp., *Ceriodaphnia* sp., and *Bosmina* sp. are the representatives of this group. According to Murugan (1998), this group consumes tiny zooplankton, bacterioplankton, and algae. They are also quite sensitive to pollutants; in fact, they might react even when the concentration of the contaminants is very low. Numerous researchers have examined the physicochemical characteristics, biotic components, and seasonal variations in zooplankton population density, composition, and abundance in freshwater bodies (Thirupathaiah et al., 2011; Patel et al., 2013).

Conclusion

The present study concludes that zooplankton were not maximum in winter season because unfavorable environmental condition.

Reference:

Ali SS. Fresh water Fisheries Biology, 1st Ed. Naseem Book Depot, Hyderabad, 1999

Acharjee, B., A. Dutta, M. Chaudhury and B.Pathak. 1995. Phytoplankton species diversity indices in dighali beel, Assam. *India Environ. Ecol.*, 13(3): 660-662.

APHA (1989). Standard methods for the examination of water and wastewater. American Public Health Association, Washington, USA.

Bogdan, K. G. and Gilbert, J. J. (1984). Body size and food size in freshwater zooplankton. *Proc. Nail.Acad. Sci. USA*, 81: 6427-6431.

Clegg, J. (1956). *The Observers Book of Pond Life*. Frederick Warne and Co. Ltd., London. pp. 1-112.

Chakrapani, B.K., M.B. Krishna and T.S. Srinivasa, 1996. A Report on the water quality, plankton and bird populations of the lakes in and around Bangalore and Maddur, Karnataka, India. Department of Ecology and Environment, Government of Karnataka.

Chourasia, S. K. and Adoni, A. D. (1985). Zooplankton dynamics of a shallow eutrophic lake. *Bull. Bot. Soc. Sugar*, 32: 30-39.

Dhanapathi M.V.S.S.S. 2000. Taxonomic notes on rotifers from India. IAAB Publ. Hyderabad. Pp.12-168.

Dhembare, A.J. 2011. Diversity and its indices in zooplankton with physicochemical properties of Mula Dam Water Ahmednagar, Maharashtra India. *Eur. J. Exp. Biol.*, 1(4): 98 103.

Edmondson, W. T. (1959). *Freshwater Biology*. 2nd ed. John Wiley and Sons Inc., New York. p. 1248.

Goldman, C. R. and Horne, A. J. (1983). *Limnology*. McGraw-Hill International Book Co., New Delhi, p. 464

Comment [W1P6]: Most of the cited literatures are not up to date

Haque, N. and Khan, A. A. (1997).Temporal and spatial distribution of cladoceran population in a freshwater lake at Aligarh. *J.Freshwater Biol.*, 6: 225-229.

Hulyal S.B., , B. B. Kaliwal (2007) Water quality assessment of Alamatti Reservoir of Bijapur (Karnataka State, India) with special reference to zooplankton Environmental monitoring Assessment (2008) 139:299-306

Jafari, N., Nabavi, S., Akhavan, M. 2011.Ecological investigation of zooplankton abundance in the river Haraz, Northeast\Iran: impact of environmental variables. *Arch. Biol. Sci. Belgrade*, 63(3): 785-798.

Jayanthi, M. (1994). A comprehensive study of three contrasting lentic systems in the context of aquaculture. Ph.D. Thesis, Bharathidasan University, Tiruchirappalli, India.

Joydip Dhar, Randhir Singh Baghel, Anuj Kumar Sharma (2012). Role of instant nutrient replenishment on plankton dynamics with diffusion in a closed system: A pattern formation Applied Mathematics and Computation 218 (2012)8925-8936.

Karuthapandi M, Rao DV, Xavier Innocent B, Deepa J. Zooplankton diversity and trophic status of Safilguda tank, Hyderabad. *Int. J Adv Lif Sci.* 2013; 6(1):44-50.

Karuthapandi, M., Rao, D.V., Xavier Innocent, B. 2013. Zooplankton composition and diversity of Umdasager, Hyderabad. *Int. J. Life Sci. Edu. Res.*, 1(1): 21 26.

Kastooribai, R. S. (1991). A comparative study of two tropical lentic systems in the context of aquaculture. Ph. D.Thesis, University of Madras, India.

Khan, A. A., Ali, M. and Haque, N. (1986). Population ecology of zooplankton in a polluted lake at Aligarh. *Proc. Nat. Symp. Environ. Biol. Coastal Ecosystem*, 475

Krishnamoorthy K, Subramanian P. Organisation of commercially supporting metroplankton in Palk Bay and Gulf of Mannar biosphere reserve areas, Southeast of India, *Indian J Mar Sci.* 1999; 28:211-215.

Madin LP, Horgan EF and Steinberg DK (2001). Zooplankton at the Bermuda Atlantic Time-series Study (BATS) station: diel, seasonal and interannual variation in biomass, 1994-1998. *Deep Sea Research*;48: 2063-2082.

Manickam N, Saravana Bhavan P, Santhanam P, Muralisankar T, Srinivasan V, Radhakrishnan S, Vijayadevan K, Chitrarasu P and Jawahar Ali A (2014). Seasonal Variations of Zooplankton Diversity in a Perennial Reservoir at Thoppaiyar, Dharmapuri District, South India. *Austin Journal of Aquaculture and Marine Biology*; 1(1): 1-7.

Maruthanayagam, C., M. Sasikumar and C. Senthilkumar, 2003. Studies on zooplankton population in Thirukkulam pond during summer and rainy seasons. *Nature Environ. Pollut. Technol.*, 2: 13-19.

Michael, R. G. (1973). A guide to the study of freshwater organisms. *J. Madurai Kamaraj University, India (Suppl.)*, pp. 1-186.

Michael, R. G. (1969). Seasonal trends in Physico-chemical factors and plankton of a freshwater fish lake and their role in fish culture. *Hydrobiologia.*, 33: 144-161.

Neetu Malik, A. K. Biswas, C. B. Raju (2013) *Plankton as an Indicator of Heavy Metal Pollution in a Freshwater Reservoir of Madhya Pradesh, India* *Bull Environ Contam Toxicol* (2013)90:725-729

Odum, E. P. (1971). *Fundamentals of ecology*. W.B. Saunders Co. Philadelphia, London, Toronto(1971).

Patel, V., Shukla, S.N., Patel, V.K. 2013. Studies on the diversity of zooplankton and their seasonal variation in Govindgarh lake at Rewa (M.P), India. *Indian J. Appl. Res.*, 3(11): 544-546.

Pennak, R. W. (1978). *Freshwater invertebrates of the United States*. 2nd ed. John Wiley and Sons, New York.p. 803.

Purushothama, R., Sayeswara, H.A., Goudar, M.A., Harishkumar, K. 2011. Physicochemical profile and zooplankton community composition in Brahmana Kalasi Tank, Sagar, Karnataka, India. *Ecoscan.*, 5(3): 99-103.

Rajasekhar, M., Vijayakumar, K. and Paerveen, Z. (2010). Seasonal variations in Zooplankton Community in Freshwater reservoir, Gulbarga District, Karnataka, South India. *Int. J. Systems Biol.*, 2: 6-11.

Russell, J., Shiel, A., Justin, F., Costelloe, B.T., Julian, R.W., Reid, A.C., Peter Hudson, A., Powling, J. 2006. Zooplankton diversity and assemblage in Aride zone river of lake Eyre Basin, Australia. *J.Mar. Freshwater Res.*, 57: 49-60.

- Shah JA, Pandit AK. Relation between physico-chemical limnology and crustacean community in Wular Lake of Kashmir Himalaya, Pak J Biol Sci. 2013; 16(19):976-983.
- Sharma, S. S. S. (1991). Rotifers and aquaculture. *Environ. Ecol.*, 9: 414-428.
- Singh, R. K., Pandey, M. K., Kumari, R. and Ranjam, P. (2012). Study on the diversity and seasonal variation of zooplankton in Mahendra Nath Pond, Siwan, Bihar. *Int. J. Pharm.Biol. Arch.*, 3: 867-890.
- Srivastava, S. K. (2013). Monthly variations in the occurrence of zooplankton in afreshwater body, Ramgarh Lake, Gorakhpur, UP. *Int. J. Appl. Biosci.*, 1: 23-27.
- Sridharan, V. T. (1989). Phytoplankton and algae studies. Techniques of plankton methodology. Prepared for Training workshop on Integrated Environmental Research programme on Kaveri River. pp. 1-15.
- Sreenivasan, A. (1974). Limnological studies and primary production in temple lake ecosystem. *Hydrobiol.*, 48: 117-125.
- Stottrup, J.G. 2000. The elusive copepods: Their production and suitability in marine aquaculture. *Aquaculture Res.*, 31: 703-711.
- Sukumaran, P.K. and A.K. Das, 2002. Plankton abundance in relation to physicochemical features in a peninsular man-made lake. *Environ. Ecol.*, 20: 873-879.
- Thakur, R. Jindal, UdayBhan Singh, A. S. Ahuwalia (2013) Plankton diversity and water quality assessment of three freshwater lakes of Mandi (Himachal Pradesh, India) with special reference to planktonic indicators Environmental monitoring Assessment (2013) 185:8355-8373
- Thirupathaiah, M., Sammatha, C.H., Sammaiah, C.H. 2011. Diversity of zooplankton in freshwater lake of Kamalapur, Karimnagar District (A.P) India. *Ecoscan*, 5(1 2): 85 87.
- Tiwari, L.R. and V.R. Nair. 1991. Contribution of zooplankton to the fishery of Dharamtar creek, adjoining Bombay harbour. *J. Indian. Fish. Ass.*, 21: 15-19.

Tidame, S. K. and Shinde, S. S. (2012). Studies on seasonal variations in physico-chemical parameters of the temple pond, Nashik District (MS), India. *Int. Multi. Res. J.*, 2: 29-32.

Vijaykumar A. Kudari, Ravishankar D. Kanamadi (2007) Impact of changed trophic status on the zooplankton composition in six water bodies of Dharwad district, Karnataka state (South India) Environmental monitoring Assessment (2008) 144:301-313

Ward, H. B., and Whipple, G. C. (1963). *Fresh-Water Biology*. John Wiley & Sons; New York

Table 1. Rotifer population of Elanthakulam pond

S.No	Rotifer	August	September	October
1	<i>Brachionus calyciflorus</i>	0	10±2	15±2
2	<i>Brachionus caudatus</i>	3±2.3	11.6±3.5	8.3±3.5
3	<i>Brachionus rubens</i>	0	4.3±0.5	3±2.5
4	<i>Keretella cochlearis</i>	6±2	2.6±1.1	9.6±2

Table 2. Cladoceran and Anostracan population of Elanthakulam pond

S.No	Cladoceran and Anostracan	August	September	October
1	<i>Daphnia pulex</i>	0.5±1.6	5±1.7	9.3±3
2	<i>Moina brachiata</i>	8.3±4.7	1±1	10±2
3	<i>Moina micrura</i>	2.6±1.1	10.6±5	6±1.7

Table 3. Copepod population of Elanthakulam pond

S.No	Copepod	August	September	October
1	<i>Diaptomus castor</i>	9.3±3	4±1	12.6±7
2	<i>Mesocyclops hyalinus</i>	3.3±1.1	13.3±5.7	8±2

Table 4. Ostracodan population of Elanthakulam pond

S.No	Ostracodan	August	September	October
1	<i>Cypris Ostracodon</i>	0	2.6±1.5	1.3±0.5


Table.5 Zooplankton Total Count of Elanthakulam pond

S.No	Zooplankton	August	September	October
1	Rotifera	14	38	56
2	Copepoda	19	26	31
3	Cladocera	19	25	38
4	Ostracoda	0	4	2
	Total	52	93	127

Figure 1



a. Collection site




b. Collection of Zooplankton


Figure 2: Rotifer




a. *Brachionus calyciflorus*



b. *Brachionus caudatus*



c. *Brachionus rubens*



d. *Keretella cochlearis*

Figure 3 - Cladoceran and Anostracan



a. *Daphnia pulex*



b. *Moina brachiata*



c. *Moina micrura*

Figure 4 - Ostracoda



a. *Cypris ostracodan*

Figure 5 -Copepod



a. *Cyclops hyalinus*



b. *Diaptamus castor*

Fig 6: Abundance of different groups of zooplankton during August,September and October in the surface water of Elanthakulam pond

