

Original Research Article

Prevalence of Malnutrition and Associated Factors Among HIV Seropositive Adults on Antiretroviral Therapy at the HIV Care and Treatment Centre of the Regional Hospital, Buea, Cameroon

ABSTRACT

Aims: The study aimed to determine the prevalence of malnutrition and associated factors among human immunodeficiency virus (HIV) seropositive adults on antiretroviral therapy at the HIV care and treatment center of the Buea Regional hospital, Cameroon.

Study design: A Hospital-based Cross sectional study design

Place and duration of study: Buea regional hospital at the HIV care and treatment centre, South West Region of Cameroon between April 2022 to June 2022.

Materials and method: We included 139 HIV seropositive individuals (17 men, 122 women age range 26 to 59 years). Nutritional status was assessed using anthropometric and clinical methods. A 24-hour dietary diversity was assessed using a questionnaire. Data was analyzed using SPSS version 25.0.

Results: The prevalence of underweight, overweight and obesity among the participants were 1.4 %, 38.8%, and 24.4% respectively. Sex of the participants ($\chi^2 = 4.715$, $p = 0.030$) and attendance to HIV-related counseling sessions ($\chi^2 = 4.512$, $p = 0.034$) were significantly associated with malnutrition. The mean dietary diversity score of the study respondents was 4.97 ± 1.6 with only 37% of the respondents achieving a minimum dietary diversity. In the multivariate logistic regression analysis, the only significant predictor of malnutrition identified was HIV counselling session. The odds of those who were counseled being malnourished was 3.29 times more than those who were not counseled (AOR=3.29, $p=0.036$).

Conclusions: There is a high burden of overweight and obesity in the HIV population of the Regional Hospital Buea. Factors significantly associated with malnutrition were HIV related counseling session and sex. Majority of the HIV seropositive patients in the Regional Hospital Buea had a poor dietary diversity. Nutritional counseling should be an important part of persons living with HIV/AIDS (PLWHA) routine follow-up.

Keywords: *Prevalence, Malnutrition, HIV, Adults, Antiretroviral therapy, Regional Hospital Buea*

ABBREVIATIONS

AIDS	: ACQUIRED IMMUNE DEFICIENCY SYNDROME
HIV	: HUMAN IMMUNODEFICIENCY VIRUS
ART	: ANTIRETROVIRAL THERAPY
BMI	: BODY MASS INDEX
CD4+	: CLUSTER OF DIFFERENTIATION 4
CDC	: CENTRE FOR DISEASE CONTROL AND PREVENTION
PLWHA	: PERSONS LIVING WITH HIV/AIDS
FAO	: FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS
FFQ	: FOOD FREQUENCY QUESTIONNAIRE
IBM	: INTERNATIONAL BUSINESS MACHINES CORPORATION
IDDS	: INDIVIDUAL DIETARY DIVERSITY SCORE
MUAC	: MID UPPER ARM CIRCUMFERENCE
WHR	: WAIST TO HIP RATIO
MDD	: MINIMUM DIETARY DIVERSITY

DDS : DIETARY DIVERSITY SCORE

1. INTRODUCTION

More than 37.7 million people are living with HIV in the world out of which about 36 million are adults. The greatest burden of the disease is concentrated in sub-Saharan Africa [1]. In 2021, the Prevalence of HIV among adults' ages 15-49 years in Cameroon was 2.9% [2]. In the Fako Division of the South West Region of Cameroon, the adult prevalence of HIV/AIDS was estimated at 3.2% in 2018 [3]. The availability of Antiretroviral Therapy (ART) has extended the lives of many people living with HIV/AIDS and has greatly reduced morbidity and death due to AIDS and its related complications [4]. Adherence to ART is essential for long-term therapeutic success because it is necessary not only in reducing the risk of emergence of HIV resistance strains, but also in improving the health status of the PLWHA [5]. ART has been shown to improve the quality of life of PLWHA [6].

Malnutrition refers to deficiencies, excesses, or imbalances in a person's intake of energy and/or nutrients [7]. Poor nutrition crosses economic lines and leads to health problems caused by eating too little (under-nourishment), too much (over nourishment) or an unbalanced diet that lacks essential nutrients for a healthy life (micronutrient deficiencies) [8]. HIV infects CD4 cells causing immune suppression which can further be worsened by poor nutrition [9, 10]. Among adults with advanced HIV infection in sub-Saharan Africa, the magnitude of malnutrition (BMI-below 18.5) was shown to be between 10 and 30%, where social factors like income, employment, inadequate food intake, and sanitation were identified as risk factors [11].

There are multiple risk factors associated to malnutrition secondary to HIV. These include food insecurity, lack of access to medical treatment, increased energy and nutrient requirements, drug side effects, gastrointestinal symptoms, reduced food intake, malabsorption and comorbidities among others [12]. In the elderly people, age-associated changes in body physiological functions, hormonal or vitamin imbalances, and sensory loss may lead to high possibility of malnutrition [13].

Dietary diversity is the number of different food groups consumed over a given period that provides a balance of nutrients that promote healthy, growth and development. Low dietary diversity is associated with a higher probability of metabolic syndrome and cardiovascular disease risk factors in adults [14]. Eating a diversity of foods (varieties of food groups) is an internationally accepted recommendation for a healthy diet, and is associated with positive health outcomes such as reduced incidence of mortality. Dietary diversity is therefore a key concept that should be promoted in managing the nutritional situation of PLWHA [15].

The staple foods of Cameroon vary from region to region and depend on climate, urbanization, household income, nutritional education, health, gender, and age group. Generally, starchy foods (e.g. corns; cassava, yam, cocoyam, potatoes; plantains; unripe bananas; and rice) dominate in Cameroonian dishes and are served with sauces, which usually contain leafy vegetables, legumes and seeds with meat or fish added, depending on affordability, availability and cultural habits [16, 17]. Although the

prevalence of malnutrition and associated factors have been reported in many countries, there is little data on the burden of malnutrition and associated factors among PLWHA in Cameroon in general and particularly in the South West Region of Cameroon. This study was therefore conducted to assess the prevalence of malnutrition and its associated factors among adult HIV/AIDS persons on ART at the HIV care and treatment centre of the Buea Regional Hospital to generate evidence-based data to inform public health interventions to improve the nutrition status of HIV-infected patients.

2. MATERIAL AND METHODS

2.1 Study design and settings

The study was a hospital-based cross-sectional study carried out at the HIV Care and Treatment Centre of the Regional Hospital Buea, a secondary health care facility. About 2000 adult HIV/AIDS persons are enrolled in the HIV/AIDS Unit (Care and Treatment Centre) in this Hospital. The Centre provides various services such as reception of HIV/AIDS patients, intake service, referral drug refill, screening, medical consultation and counselling.

2.3 Study population and sampling

The study population consisted of adult persons living with HIV/AIDS who had been on uninterrupted ART for at least six months. Adult HIV/AIDS persons who had kyphoscoliosis were excluded from the study. A total of 139 HIV Seropositive adults on antiretroviral therapy who consented were included in to the study. Recruitment of participants into the study was consecutive.

2.5 Data collection

Data on demographic factors, socioeconomic factors, and health-related factors were collected using a structured questionnaire from April 2022 to June, 2022. The clinical characteristics of the patients were extracted from their treatment registers using a data extraction form. Measurement of body weight was conducted using a manual weighing balance and recorded to the nearest 0.1kg. The body weight was taken with participants in light clothing and shoes taken off. Measurement of height was conducted using a measuring gauge that was regulated to the nearest 0.1cm. BMI (kg/m^2) was calculated as a proxy nutritional status by dividing body weight (kg) by the square of the height (m^2). It was classified according to the conventional classification of the World Health Organization. BMI was stratified according to the WHO criteria: $<18.5 \text{ kg/m}^2$ (underweight), >18.5 to 25 kg/m^2 (normal weight), and $>25 \text{ kg/m}^2$ (overweight and obese) [20]. The measurement of waist and hip circumferences was conducted using a stretched resistant tape that was regulated to the nearest 0.1cm. Waist circumference was measured at the midpoint between the lower margin of the least palpable rib and the top of the iliac crest, using a stretch-resistant tape. Hip circumference was measured around the widest portion of the buttocks. For both measurements, the subject stood with feet close together, arms at the side that guarantee evenly distribution of the body weight. The subjects were relaxed, and the measurements were

taken at the end of a normal expiration. The measurements were collected twice and the average of each taken [21]. The waist to hip ratio (WHR) was calculated as waist circumference (cm) divided by the hip circumference (cm) to get the abdominal obesity [22]. Guideline and cut-offs for underweight, normal weight, overweight, obesity for BMI were, <18.5, < 18.5 to 24.9, 25 to 29.9, > 30 [20]. respectively and WHR for men low, moderate and high is ≤ 0.95 , 0.96 to 1.0, and ≥ 1.0 respectively and for women is ≤ 0.80 , 0.81 to 0.85 and ≥ 0.86 [22] respectively.

Minimum dietary diversity was assessed using a minimum dietary diversity questionnaire, as indicator of the nutrient adequacy of the diet of the persons based on guidelines from the Food and Agricultural Organization (FAO) [23]. Dietary diversity was recognized by using the approved 24- hour recall method adopted from the food and agriculture organization (FAO). The MDD was made up of 10 main food groups. A PLWHA was classified as having food insecurity or inadequate dietary diversity if he or she consumed less than five food groups. A PLWHA who consumed five or more food groups was classified as having reached MDD with good diet variety and food security [24].

2.6 Data analysis

Data was analyzed using the Statistical Package for Social Sciences (SPSS) Version 25.0. Descriptive statistics were performed for categorical data, mean and standard deviation and median were calculated for continuous variables. Logistic regression was used to identify factors associated to malnutrition. Simple logistic regression was used to screen predictors at p-value < 0.2 and a multiple logistic regression was used to select the factors associated with nutritional status of children at p-value ≤ 0.05 .

3. RESULTS

3.1 Socio-demographic Characteristics of study participants

A total of 139 participants were enrolled into the study of which 122 (87.8%) were females. The mean age of participants was 43 ± 7 years with 46.6% participants in the age group of 36 - 45 years. Majority of the participants 134(96.4. %) were Christians with, 58 (41.7%) were married, 69 (49.6%), were self-employed and 77 (55.4%) had an estimated monthly income (FCFA) between 50,000 to 100,000 (Table 1).

Table 1: Socio-demographic characteristics of study participants

Variable	Categories	Frequency	Percentage
Sex	Female	122	87.8
	Male	17	12.2

	Total	139	100
Age groups (years)	26- 35	18	12.9
	36 – 45	62	44.6
	46 – 55	50	36
	> 55	9	6.5
	Total	139	100
Marital Status	Cohabiting	5	3.6
	Divorced	11	7.9
	Married/Concubinage	58	41.7
	Single	46	33.1
	Widow(er)	19	13.7
	Total	139	100
Religion	Christian	134	96.4
	Muslim	5	3.6
	Total	139	100
Profession	Private employee	37	26.6
	Public employee	19	13.7
	Self-employee	69	49.6
	Unemployed	14	10.1
	Total	139	100
Education Level	Not a student	4	2.9
	High school	14	10.1
	Primary school	61	43.9
	Secondary school	47	33.8
	University	13	9.4
	Total	139	100
Residence	Rural	102	73.4
	Urban	37	26.6
	Total	139	100
Household size	1 _ 3	54	38.8
	4 _ 6	62	44.6
	7 +	23	16.5
	Total	139	100
Food money per day	1000 – 2000	94	67.6
	3000 – 5000	40	28.8
	> 5000	5	3.6
	Total	139	100
Income (FCFA)	<50000	55	39.6
	>100000	7	5
	50000-100000	77	55.4
	>100000	7	5

	Total	139	100
--	--------------	------------	------------

3.2 Clinical characteristics of the study participants

Regarding the HIV/AIDS-related characteristics of participants; more than half 78 (56.1%) were at clinical stage I, 115 (83.3%) of the participants had an undetectable viral load, 126 (90.6%) had disclosed their HIV status, 117(84.2%) were on first line treatment and 15 (10.8%) did not attend HIV-related counseling (table 2).

Table 2: Clinical characteristics of the study participants

Variable	Categories	Frequency	Percent
Duration of ART	≥3 years	119	85.6
	6 months to 3 years	20	14.4
	Total	139	100
Clinical stage	Stage I	78	56.1
	Stage II	18	12.9
	Stage III	39	28.1
	Stage IV	4	2.9
	Total	139	100
HIV advance disease	No	136	97.8
	Yes	3	2.2
	Total	139	100
Condition	Cerebral Toxoplasmosis	1	33.3
	Pulmonary TB	2	66.7
	Total	3	100
Most recent viral load (copies/ml)	40-1000	17	12.3
	>1000	6	4.3
	Undetectable	115	83.3
	Total	139	100
HIV/AIDS-related symptoms	No	133	95.7
	Yes	6	4.3
	Total	139	100
Treatment Regimen	1st line	117	84.2
	2nd line	22	15.8
	Total	139	100
Depressed or anxious	No	128	92.1
	Yes	11	7.9
	Total	139	100

Disclosure of HIV/AIDS status	No	13	9.4
	Yes	126	90.6
	Total	139	100
HIV related counselling session	No	15	10.8
	Yes	124	89.2
	Total	139	100
Diarrhoea	No	135	97.1
	Yes	4	2.9
	Total	139	100

3.3 Prevalence of malnutrition among HIV/AIDS patients based on Body Mass Index (BMI)

The average BMI of the study population was $24.56 \pm 4.97 \text{ kg/m}^2$. Based on BMI classification, 2(1.4%) of the participants were underweight, 49(35.3%) had normal weight, 54(38.8%) were overweight and 34(24.4%) were obese (Figure 1).

Figure 1: Prevalence of malnutrition among PLWHA based on BMI

Malnutrition in HIV persons classified based on the waist-hip ratio

Table 3 shows the waist to hip ratio categories whereas figure 3 shows the variation of WHR in the study population. Abdominal obesity was more present in women 67(48.2%) than in men 1(0.7%) as shown in figure 2. The mean WHR was 2.5 ± 13.3 . The overall prevalence of malnutrition based on WHR was 68(48.9%).

Table 3: Categories of Waist-hip ratio of persons living with HIV/AIDS
Categories of Waist-hip ratio of persons living with HIV/AIDS

Variable	Categories	Frequency(n)	Percent(%)
Waist/Hip ratio categories	High	68	48.9
	Low	38	27.3
	Moderate	33	23.7
	Total	139	100
Malnutrition(overnutrition)	Normal	71	51.1
	Malnourished	68	48.9
	Total	139	100

Figure 2: WHR distribution by Sex among PLWHA at the Buea Regional Hospital

3.4 Demographic factors associated with malnutrition among study participants

Of the demographic factors studied, there was a significant association between sex and malnutrition ($\chi^2=4.72$, $p=0.03$) with females being more malnourished than males (table 3).

Table 4: Association between malnutrition and socio-demographic characteristics

Variables	Categories	n	Nutritional status				2	p-value
			Malnourished	(%)	Normal	(%)		
Residence	Rural	102	67	48.2	35	25.18	0.15	0.701
	Urban	37	23	16.55	14	10.07		
	Total	139	90	64.75	49	35.25		
Sex	Female	122	83	59.71	39	28.06	4.72	0.030
	Male	17	7	5.04	10	7.19		
	Total	139	90	64.75	49	35.25		
Age groups (years)	> 55	9	6	4.32	3	2.16	3.74	0.291
	26- 35	18	8	5.76	10	7.19		
	36 – 45	62	42	30.22	20	14.39		
	46 – 55	50	34	24.46	16	11.51		
	Total	139	90	64.75	49	35.25		
Marital Status	Cohabiting	5	4	2.88	1	0.72	5.79	0.204
	Divorced	11	6	4.32	5	3.6		
	Married/Concubine	58	32	23.02	26	18.71		
	Single	46	33	23.74	13	9.35		
	Widow(er)	19	15	10.79	4	2.88		
	Total	139	90	64.75	49	35.25		
Religion	Christian	134	86	61.87	48	34.53	0.53	0.421
	Muslim	5	4	2.88	1	0.72		
	Total	139	90	64.75	49	35.25		
Profession	Private employee	37	24	17.27	13	9.35	1.57	0.667
	Public employee	19	10	7.19	9	6.47		
	Self-employee	69	47	33.81	22	15.83		
	Unemployed	14	9	6.47	5	3.6		
	Total	139	90	64.75	49	35.25		
Education Level	High school	14	10	7.19	4	2.88	1.75	0.808
	Not a student	4	2	1.44	2	1.44		

	Primary school	61	41	29.5	20	14.39		
	Secondary school	47	30	21.58	17	12.23		
	University	13	7	5.04	6	4.32		
	Total	139	90	64.75	49	35.25		
Household size	1 _ 3	54	38	27.34	16	11.51	1.22	0.542
	4 _ 6	62	38	27.34	24	17.27		
	7 +	23	14	10.07	9	6.47		
	Total	139	90	64.75	49	35.25		
Income (FCFA)	<50000	55	32	23.02	23	16.55	2.32	0.309
	50000-10000	77	54	38.85	23	16.55		
	>100000	7	4	2.88	3	2.16		
	Total	139	90	64.75	49	35.25		
Food money(FCFA)	1000 – 2000	94	61	43.88	33	23.74	1.46	0.478
	3000 – 5000	40	27	19.42	13	9.35		
	> 5000	5	2	1.44	3	2.16		
	Total	139	90	64.75	49	35.25		

3.5 Association between malnutrition and clinical characteristics

Table 4 presents the association bet malnutrition clinical variables. Attendance at HIV-related counseling sessions was significantly associated to malnutrition ($\chi^2=4.512$, $p=0.03$). Those who attended HIV-related counselling were more malnourished (60.43) than those who did not attend the counselling.

Table 5: Association between malnutrition and clinical characteristics of participants

Variables	Categories	n	Malnutrition				χ^2	P-value
			Malnourished	%	Normal	%		
Duration on ART	3 years and above	119	77	55.4	42	30.22	0.01 0.980	
	6 months to 3 years	20	13	9.35	7	5.04		
	Total	139	90	64.75	49	35.25		
Clinical stage	Stage I	78	53	38.13	25	17.99	2.27 0.530	
	Stage II	18	9	6.47	9	6.47		
	Stage III	39	25	17.99	14	10.07		
	Stage IV	4	3	2.16	1	0.72		
	Total	139	90	64.75	49	35.25		
Recently diagnosed with HIV advanced disease	No	136	87	62.59	49	35.25	1.67	0.196
	Yes	3	3	2.16	0	0		
	Total	139	90	64.75	49	35.25		

Most recent viral load (copies/ml)	40 – 1000	17	9	6.47	8	5.76	1.80	0.411
	>1000	6	5	3.6	1	0.72		
	Undetectable	115	75	53.96	40	28.78		
	Total	138	89	64.03	49	35.25		
HIV/AIDS related symptoms	No	133	86	64.66	47	35.33	1.81	0.473
	Yes	6	4	2.88	2	1.44		
	Total	139	90	64.75	49	35.25		
Treatment regimen	1st line	117	79	56.83	38	27.34	2.49	0.115
	2nd line	22	11	7.91	11	7.91		
	Total	139	90	64.75	49	35.25		
Depression and anxiety	No	128	82	58.99	46	33.09	0.33	0.564
	Yes	11	8	5.76	3	2.16		
	Total	139	90	64.75	49	35.25		
Disclosure of HIV/AIDS status	No	13	9	6.47	4	2.88	0.13	0.722
	Yes	126	81	58.27	45	32.37		
	Total	139	90	64.75	49	35.25		
HIV related counseling session	No	15	6	4.32	9	6.47	4.51	0.034
	Yes	124	84	60.43	40	28.78		
	Total	139	90	64.75	49	35.25		
Diarrhoea	No	135	87	62.59	48	34.53	0.19	0.663
	Yes	4	3	2.16	1	0.72		
	Total	139	90	64.75	49	35.25		

In the multivariate logistic regression analysis, only HIV counseling session was associated with nutritional status. The odds of those who attended HIV counseling sessions being malnourished was 3.29 times higher than those who did not attend any HIV related counseling session (AOR=3.29, p=0.036) (table 6).

Table 6: Factors independently associated with malnutrition among HIV/AIDS persons on follow-up care at the Buea Regional Hospital

Variable	Levels	AOR	95% CI		P-value
			Lower	Upper	
Treatment Regimen	2nd line	0.46	0.18	1.17	0.102
	1st line	1			
HIV related counselling session	Yes	3.29	1.08	9.99	0.036
	No	1			

3.6 Dietary diversity of PLWHA

Among the ten main food groups, participants consumed more of grains, white roots and tubers, and plantains 139 (100%), meat, poultry and fish 119 (85.6%), and dark

green leafy vegetables 67 (48.2%), while eggs 15 (10.8%) was the lowest consumed food (Figure 3).

Figure 3: Distribution of 10 main groups consumed in the last 24-hour by HIV/AIDS persons aged 21-59 years old

For the dietary diversity, only 51 (37%) had a good dietary diversity while 63%% had poor dietary diversity (Figure 4).

Figure 4: Distribution of participants' dietary diversity score

4. Discussion

Prevalence of malnutrition:

The majority of the study participants were overweight and obese. The prevalence of undernutrition in our study was 1.4%. This was lower compared to that reported (8.5%) in a study conducted in the Centre Region of Cameroon [19], 19.5% reported in Tanzania [25], 10% in Zimbabwe[26] and 19.2% in Senegal [27]. The difference in the prevalence of undernutrition might be due to differences in socioeconomic and other factors such as food habit and cultures. Also, these differences could be explained by the discrepancy in health care awareness of the community or feeding practices of different ethnic groups. Improvement in healthcare services could also account for these differences.

Overweight and obesity were more dominant in our study than undernutrition. This result raises a concern with the quality of food consumed by participants. Many food types consumed by our study participants are rich in carbohydrates and this could explain the high prevalence of overweight and obesity in this study population. Also, Dolutegravir, a recently added integrase inhibitor antiretroviral drug has been associate to short-term weight gain in HIV/AIDS persons who begin ART[28]. Mostly women were obese in our study cohort with almost half of the study population with abdominal obesity. Abdominal obesity might reflect a metabolic syndrome which has been shown to affect 32.8% of PLWHA in Cameroon [29].

Factors associated with malnutrition

Females were more malnourished than males. This finding is similar to that of a study conducted in Ethiopia [30]. This may be due to the low number of males that were included in the study. In order to sustain their children and husband's life, women live a more sedentary life style to pay attention to their family. Those who attended counseling sessions were more prone to malnutrition than those who did not. This might be because not all persons receive nutritional advice during their follow-up, although all of

them received health talks. This suggests that the reinforcement of the caregiver's implication on nutritional education of persons should be improved. This finding is comparable to that reported by Perpetue *et al.*, (56.1%) in 2021 in Cameroon [19].

The only predictor of malnutrition identified was the treatment regimen, although there was no significant association. The odds of those on 2nd line treatment regimen being malnourished was 0.46 less than those who were on 1st line treatment. Those who were in clinical stage I were more malnourished than those in clinical stage II, III and IV, but the difference was not significant. This is similar to a study reported in Ethiopia where the stage of HIV didn't show any significant effect on respondent's nutritional status [30]. Our findings are contrary to those of a studies conducted at Sella Hospital in Ethiopia [31], where WHO clinical stage II was significantly associated to undernutrition. A study also revealed that individuals at all stages of HIV disease are at risk of nutritional deficiency, but clinical stages show the severity of the disease from primary HIV infection to advanced stages of HIV or AIDS [32]. This discrepancy might be due to the clinical stage of the study participants, where majority of them are found at the clinical stage one in the current study.

With regards to HIV/AIDS related symptoms, there was no significant association between malnutrition and HIV/AIDS related symptoms. This is in contrast to the findings of a study carried out at the Elege Hiwot referral Hospital in Ethiopia where those who developed disease symptoms two weeks prior to the survey were almost two times more malnourished than those who were free of symptoms [33].

In this study, the duration on ART was not significantly associated to malnutrition. This is in contrast to reports by Daniel and collaborators in a study carried out in Ethiopia where patients who were on ARV drugs for less than 12 months were 1.7 times more malnourished than those who took the drug more than a year [33]. ART contributes to the wellbeing of people living with HIV (PLHIV), and reduces the threat of the ongoing transmission of HIV.

Minimum dietary diversity:

Dietary diversity which is the number of different food groups consumed over a given time period is of great importance as it can be used as a simple indicator of nutritional adequacy in individuals [34]. The present study revealed that only 37% of the participants had achieved a minimum dietary diversity. This is similar to findings from a study carried out in Ethiopia where 29.5 % of participants consumed a diversified diet [35]. The poor dietary diversity observed in our study could be due to the ongoing socio-political crisis in the South West Region of Cameroon which might prevent PLWHA to have access to diversified foods. However, this percentage is lower than that reported in a study carried out in Ethiopia at the Hiwot Fana and Dilchora Hospitals that reported an MDD of 71.3%. [36]. This discrepancy might be due to variation in the data collection periods, i.e. data collection period for the Hiwot Fana & Dilchora Hospitals was November to February, which was harvesting time when there was better food access.

CONCLUSIONS

The prevalence of overweight and obesity among PLHIV at the Buea regional hospital treatment site (was very high with female more affected than males. Factors significantly associated with malnutrition were HIV-related counseling and sex of the participants. Majority of HIV/AIDS patients in Buea had a poor dietary diversity. This study highlight the need to develop and implement a nutritional based intervention in HIV treatment centre for the improvement of the nutritional status of PLHIV.

ETHICAL APPROVAL AND CONSENT

This study was approved by the Institutional Review Board of the Faculty of Health Sciences, University of Buea (Ref. No. 2022/177104/UB/SG/IRB/FHS). Administrative clearance was obtained from the Regional Delegation of Public Health for the South West region, (Ref. No. RII/MINSANTE/SWR/RDPH/PS/419/424) and written informed consent was obtained from the patients recruited into this study.

REFERENCES

1. UNAIDS (2021). Global HIV & AIDS statistics — Fact sheet. <https://www.unaids.org/en/resources/fact-sheet>. Accessed 19 Nov 2021
2. Knoema (2021). Cameroon HIV prevalence, 1960-2022. In: Knoema. <https://knoema.com//atlas/Cameroon/HIV-prevalence>. Accessed 21 May 2023
3. Bonono L (2021). Cameroon National Strategic Plan For fight against HIV/AIDS and STIs 2021-2023.
4. Mekuria A, Chernet H, Habtamu J (2015). Malnutrition and Associated Factors among Adult Individuals Receiving Highly Active Antiretroviral Therapy in Health Facilities of Hosanna Town, Southern Ethiopia. *Open Access Library Journal* 02:1
5. Yaya I, Landoh DE, Saka B, Patchali PM, Wasswa P, Aboubakari A, N'Dri MK, Patassi AA, Kombaté K, Pitche P (2014). Predictors of adherence to antiretroviral therapy among people living with HIV and AIDS at the regional hospital of Sokodé, Togo. *BMC Public Health* 14:1308
6. Yaya I, Djalogue L, Patassi AA, (2019). Health-related quality of life among people living with HIV/AIDS in Togo: individuals and contextual effects. *BMC Res Notes* 12:1–6
7. WHO (2021) Fact sheets - Malnutrition. In: World Health Organization. <https://www.who.int/news-room/fact-sheets/detail/malnutrition>. Accessed 30 Jan 2022
8. FAO (2010). The spectrum of malnutrition. <https://www.fao.org/3/i2490e/i2490e02b.pdf>. Accessed 21 May 2023

9. Gebru TH, Mekonen HH, Kiros KG (2020). Undernutrition and associated factors among adult HIV/AIDS patients receiving antiretroviral therapy in eastern zone of Tigray, Northern Ethiopia: a cross-sectional study. *Arch Public Health* 78:100
10. Mulu H, Hamza L, Alemseged F (2016). Prevalence of Malnutrition and Associated Factors among Hospitalized Patients with Acquired Immunodeficiency Syndrome in Jimma University Specialized Hospital, Ethiopia. *Ethiop J Health Sci* 26:217–226
11. Kabalimu TK, Sungwa E, Lwabukuna WC (2018). Malnutrition and associated factors among adults starting on antiretroviral therapy at PASADA Hospital in Temeke District, Tanzania. *Tanzania Journal of Health Research* 20:
12. Osuna-Padilla IA, Salazar Arenas M de los A, Rodríguez- Moguel NC, Aguilar-Vargas A, Montano Rivas JA, Ávila-Ríos S (2022). Phase angle as predictor of malnutrition in people living with HIV/AIDS. *Nutrition in Clinical Practice* 37:146–152
13. Thet D, Siritientong T, Sangarlangkarn A, Lwin HMS, Phonphithak S, Apornpong T, Avihingsanon A (2022). Deterioration of Nutritional Status and Its Negative Association with Depression Among Older HIV-Infected Asian Population: A Four-Year Longitudinal Study. *AIDS Research and Human Retroviruses* 38:592–600
14. Drimie S, Faber M, Vearey J, Nunez L (2013) Dietary diversity of formal and informal residents in Johannesburg, South Africa.
15. Weldegebreal F, Digaffe T, Mesfin F, Mitiku H (2018) Dietary diversity and associated factors among HIV positive adults attending antiretroviral therapy clinics at Hiwot Fana and Dilchora Hospitals, eastern Ethiopia. *HIV AIDS (Auckl)* 10:63–72
16. Sharma S, Mbanya JC, Cruickshank K, Cade J, Tanya AKN, Cao X, Hurbos M, Wong MR (2007) Nutritional composition of commonly consumed composite dishes from the Central Province of Cameroon. *International journal of food sciences and nutrition*. <https://doi.org/10.1080/09637480701288454>
17. Mennen LI, Mbanya JC, Cade J, Balkau B, Sharma S, Chungong S, Cruickshank JK (2000). The habitual diet in rural and urban Cameroon. *Eur J Clin Nutr* 54:150–154
18. Moenikia M, Zahed-Babelan A (2010). A study of simple and multiple relations between mathematics attitude, academic motivation and intelligence quotient with mathematics achievement. *Procedia - Social and Behavioral Sciences* 2:1537–1542
19. Perpetue FFD, Charles KT, Anatole PC (2021). Nutritional Assessment of HIV/AIDS patients in Centre Region of Cameroon: A pilot study. *Journal of Food Science and Nutrition Research* 4:77–93
20. World Health Organization (2011). Waist circumference and waist-hip ratio : report of a WHO expert consultation, Geneva, 8-11 December 2008.

21. World Health Organization (2011). Waist circumference and waist-hip ratio : report of a WHO expert consultation, Geneva, 8-11 December 2008.
22. Lana B (2017). Waist-to-hip ratio: How does it affect your health? Medical News Today
23. Kennedy G, Ballard T, Dop MC, European Union (2011). Guidelines for measuring household and individual dietary diversity. Food and Agriculture Organization of the United Nations, Rome
24. Weerasekara PC, Withanachchi CR, Ginigaddara GAS, Ploeger A (2020). Understanding Dietary Diversity, Dietary Practices and Changes in Food Patterns in Marginalised Societies in Sri Lanka. *Foods* 9:1659
25. Kabalimu TK, Sungwa E, Lwabukuna WC (2018). Malnutrition and associated factors among adults starting on antiretroviral therapy at PASADA Hospital in Temeke District, Tanzania. *Tanzania Journal of Health Research*. <https://doi.org/10.4314/thrb.v20i2.5>
26. Takarinda KC, Mutasa-Apollo T, Madzima B, Nkomo B, Chigumira A, Banda M, Muti M, Harries AD, Mugurungi O (2017). Malnutrition status and associated factors among HIV-positive patients enrolled in ART clinics in Zimbabwe. *BMC Nutrition* 3:15
27. Benzekri NA, Sambou J, Diaw B, Sall F, Niang S, Ba S, Guèye NFN, Diallo BM, Hawes SM, Gottlieb GS (2015). High Prevalence of Severe Food Insecurity and Malnutrition among HIV-Infected Adults in Senegal, West Africa. *PLoS One* 10:e0141819
28. Eifa BA, Ketema W (2022). Could a Dolutegravir-Based Antiretroviral Therapy Lead to Clinical Obesity? A Retrospective Cohort Study Conducted at Hawassa University Comprehensive Specialized Hospital in Hawassa, Sidama, Ethiopia. *AIDS Research and Treatment* 2022:e2965325
29. Mukabana B, Masika F (2018). Factors Affecting Dietary Intake and Dietary Diversity Score Among Adults Living With HIV/AIDS in Uasin Gishu District Hospital, Kenya - A Cross Sectional Study. <https://www.iosrjournals.org/iosr-jnhs/papers/vol7-issue6/Version-5/C0706051018.pdf>. Accessed 27 Jun 2022
30. Kenea MA, Garoma S, Gemedie HF (2015). Assessment of adult nutritional status and associated factors among ART users in Nekemte referral hospital and health center, east Wollega zone, Ethiopia. *J Food Nutr Sci* 3:56–63
31. Teklu T, Chauhan NM, Lemessa F, Teshome G (2020). Assessment of Prevalence of Malnutrition and Its Associated Factors among AIDS Patients from Asella, Oromia, Ethiopia. *BioMed Research International* 2020:e7360190
32. Munthali C, Taegtmeyer M, Garner PG, Lalloo DG, Squire SB, Corbett EL, Ford N, MacPherson P (2014). Diagnostic accuracy of the WHO clinical staging system for

defining eligibility for ART in sub-Saharan Africa: a systematic review and meta-analysis. *Journal of the International AIDS Society* 17:18932

33. Daniel M, Mazengia F, Birhanu D (2013). Nutritional status and associated factors among adult HIV/AIDS clients in Felege Hiwot Referral Hospital, Bahir Dar, Ethiopia. *Science Journal of Public Health* 1:24–31

34. Puwanant M, Boonrusmee S, Jaruratanasirikul S, Chimrung K, Sriplung H (2022). Dietary diversity and micronutrient adequacy among women of reproductive age: a cross-sectional study in Southern Thailand. *BMC Nutrition* 8:127

35. Tesfaw A, Jara D, Temesgen H (2018) Dietary Diversity and Associated Factors among HIV Positive Adult Patients Attending Public Health Facilities in Motta Town, East Gojjam Zone, Northwest Ethiopia, 2017. *Advances in Public Health* 2018:e6135482

36. Weldegebreal F, Digaffe T, Mesfin F, Mitiku H (2018). Dietary diversity and associated factors among HIV positive adults attending antiretroviral therapy clinics at Hiwot Fana and Dilchora Hospitals, eastern Ethiopia. *HIV* 10:63–72