

Original Research Article

Comparative performance of different fertilizer Recommendation methods on growth and yield of rice in Old Brahmaputra Floodplain soils

ABSTRACT

The experiment was conducted at the Soil Science Field Laboratory of Bangladesh Agricultural University, Mymensingh, Bangladesh from February 2020 to May 2020 to study the comparative performance of different fertilizer recommendation methods in rice (BRRI dhan29) in Old Brahmaputra Floodplain Soils (AEZ 9). The experiment was laid out in a randomized complete block design with three replications. There were six treatments namely, T1 (Control-no fertilizer), T2 (Farmers' Practice), T3 (Fertilizer Recommendation Guide-2018), T4 (BAU Soil Testing Kit), T5 (Soil Test Basis) and T6 (Rice Crop Manager). The results revealed that all the treatments showed better performances over control (T1). Treatment T4 (BAU Soil Testing Kit) produced the highest value of plant height (90.18 cm), panicle length (24.90), filled grains panicle⁻¹ (122.60) and 1000-grain weight (26.67g) while T5 (Soil Test Basis) produced maximum effective tillers hill⁻¹ (12.80). The highest grain yield (6.29 t ha⁻¹) and straw yield (7.73 t ha⁻¹) were recorded in treatment T4 (BAU Soil Testing Kit). Treatment T4 (BAU Soil Testing Kit) also showed the highest grain yield increase (61.49%) and straw yield increase (90.49%) over control. The highest total N, P and K uptake (108.84 kg ha⁻¹, 24.54 kg ha⁻¹ and 128.17 kg ha⁻¹) were obtained in the treatment T4 (BAU Soil Testing Kit). Control treatment T1 (No fertilizers) showed minimum performances in all the cases compared to other treatments. The overall result demonstrate that the application of NPK fertilizers following BAU Soil Testing Kit method of fertilizer application system is a better option for maximum and eco-friendly rice production.

Keywords: Fertilizer recommendation, methods, rice, Bangladesh

1. INTRODUCTION

Bangladesh is heavily reliant on intensive rice cultivation, ranking third globally in both area and production of rice. Rice serves as a staple for over three billion people worldwide. In 2019, Bangladesh produced 37.4 million tons of rice, contributing significantly to the national economy and food security. However, the country faces challenges such as a low average rice yield of 3.92 t ha⁻¹ compared to other nations like China and Japan (FAO, 2014).

With an annual per capita consumption of 198 kg, rice is a vital commodity in Bangladesh, utilizing 84.67% of total cropped land. Despite the country's efforts to increase production to meet the growing population's demands, challenges such as diminishing cultivable land persist. Experts anticipate a potential increase in rice production to 46.7 million tons by 2050 (Kabir et al., 2021).

Fertilizer plays a crucial role in sustaining high crop production. But the heavy dependence on NPKS fertilizers pose environmental and food safety concerns. Continuous use accelerates soil degradation, affecting physical and chemical properties. Constituting 20% of rice production costs, fertilizer expenses prompt substantial government subsidies, reaching 119 billion Taka in 2012–2013 (MoF, 2015). Crop scientists around the world are searching for sustainable farming techniques to keep the soil healthy and decrease the amount of fertilizer needed for per kg of rice produced (Betteridge et al., 2008).

This study aims to compare different fertilizer recommendation methods to enhance fertilizer use efficiency, preserve soil health, protect the environment, and achieve maximum production in the current agricultural landscape of Bangladesh. The objectives include:

1. Compare fertilizer recommendation methods and identify the best.
2. Assess the impact of methods on rice growth and yield.
3. Examine the effect of different fertilizer treatments on nutrient uptake by the rice crop.

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The experiment was conducted at Bangladesh Agricultural University's Soil Science Field Laboratory in Mymensingh (24.75° N latitude, 90.45° E longitude, 18.978m above the sea level) during the boro season (February-May, 2020) on BRRi dhan29, a widely used HYV boro rice variety in Bangladesh. The climate is sub-tropical with distinct seasonal variations. The soil, categorized as non-calcareous Dark Grey Floodplain under AEZ9, is moderately well-drained silt loam.

Land preparation involved ploughing, cross ploughing with a power tiller, and laddering. After levelling and puddling, experimental plots were arranged per treatments. Soil sampling, conducted at 0-15 cm depth before ploughing, produced composite samples from 10 spots in each plot. These samples underwent meticulous cleaning and were stored in plastic containers for subsequent physical and chemical analyses.

The experiment was comprised of six treatments including control.

T1: Control (No fertilizer)

T2: Farmers practice

T3: Fertilizer Recommendation Guide (FRG)-2018

T4: BAU soil testing kit

T5: Soil Test Basis (STB)

T6: Rice crop manager

Fertilizer rates of NPKS fertilizers for 10 m² area following different recommendation methods are presented in table .1

Table .1 Fertilizer rates of the treatments following different recommendation methods used for the experiment

Treatments	Name of the fertilizers				
	Urea (g 10 m ⁻² split ⁻¹)	TSP (g 10 m ⁻²)	MoP (g 10 m ⁻²)	Gypsum (g 10 m ⁻²)	ZnO (g 10 m ⁻²)
T ₁	0.00	0	0	0.00	0.00
T ₂	82.00	100	86	61.00	0.00
T ₃	103.68	105	120	44.40	1.92
T ₄	130.00	20	226	44.40	1.92
T ₅	135.00	35	180	94.00	1.92
T ₆	90.00	80	36	31.00	1.30

Table .2 Nutrients and their sources used for the experiment

Nutrient	Sources
Nitrogen (N)	Urea
Phosphorus (P)	Triple super phosphate
Potassium (K)	Muriate of potash
Sulphur (S)	Gypsum

Seedlings of BRR1 dhan29 rice were carefully uprooted from the seedbed before transplanting, with the transplantation occurring on February 13, 2020. Thirty-five-day-old seedlings were transplanted into the plots. The experiment employed a randomized complete block design with three replications, incorporating six treatments, including a control. Each block was subdivided into unit plots, resulting in a total of 18 plots. The unit plot size was 4×2.5 m, with 1 m spacing between blocks and 0.5 m between plots. To ensure proper crop growth, various intercultural operations were undertaken such as weeding, pest control, irrigation, drainage etc.

Harvesting occurred at full maturity, with crops bundled, threshed, cleaned, and processed. Grain and straw yields were recorded based on 14% moisture content. Sample analysis involved collecting 100 grams from each plot, oven drying, and grinding for chemical analysis. Various plant characteristics, including plant height, panicle length, number of effective tillers, number of filled grains per panicle, weight of 1000 grains, and grain and straw yields, were measured and recorded. After sun drying, straw and grain samples underwent digestion and analysis for nitrogen, phosphorus, potassium, and sulfur content using specified methods. To calculate nutrient uptake with grain and straw the following equation was used.

$$\text{Nutrient uptake (kg ha}^{-1}\text{)} = (\text{Gy} \times \text{Ngr}) / 100 + (\text{Sty} \times \text{Nst}) / 100$$

Where,

Gy= Grain yield (kg ha^{-1}), Sty= Straw yield (kg ha^{-1}), Ngr = Nutrient content in grain (%), Nst = Nutrient content in Straw (%)

Data were analysed statistically by ANOVA to examine whether treatment effects were significant (Gomez and Gomez, 1984). Mean values were compared by Duncan's Multiple Range Test (DMRT). Software package, Statistix10 was followed for statistical analysis.

3. RESULTS AND DISCUSSION

Yield contributing characteristics of rice consists of plant height, effective tillers hill^{-1} , panicle length, filled grains panicle^{-1} , and 1000-grain weight. The effects of NPK fertilizers on the plant height of BRRI dhan29 has been ranked in order of $T_2 > T_6 > T_3 > T_4 > T_5 > T_1$, the number of effective tillers hill^{-1} has been ranked in order of $T_5 > T_4 > T_3 > T_6 > T_2 > T_1$, panicle length has been ranked in order of $T_4 > T_6 > T_5 > T_3 > T_2 > T_1$, filled grains panicle^{-1} has been ranked in the order of $T_4 > T_5 > T_3 > T_2 > T_6 > T_1$ and the values of 1000-grains weight for treatments T_1, T_2, T_3, T_4, T_5 and T_6 has been recorded as 25.00g, 26.00g, 26.50g, 26.67g, 26.67g and 26.13g respectively.

List 1 :Effect of different fertilizer application methods on several plant characteristics of BRRI dhan29

Treatments	Plant height(cm)	No. of effective tillers hill^{-1} (No.)	Panicle length(cm)	Filled grains panicle^{-1} (No.)	1000- grain weight (g)
T_1	78.47b	9.27c	20.20b	83.07c	25.00b
T_2	90.97a	10.87bc	22.84a	122.53ab	26.00ab
T_3	90.63a	12.13ab	23.43a	114.77ab	26.50a
T_4	90.18a	12.14ab	24.90a	122.60a	26.67a
T_5	90.17a	12.80a	23.50a	116.5ab	26.67a
T_6	90.64a	11.53ab	23.60a	111.53b	26.13a
SE \pm	2.03	0.86	1.04	3.96	0.497
CV (%)	2.81	9.2	5.54	4.41	2.33

The figure (s) having common letter (s) in a column do not differ significantly at 5% level of significance. SE (\pm) = Standard Error of Means, CV (%) = Co-efficient of variance

T₁: Control (No fertilizer), T₂: Farmers practice, T₃: Fertilizer Recommendation Guide (FRG)-2018, T₄: BAU soil testing kit, T₅: Soil Test Basis (STB) and T₆: Rice crop manager (RCP)

The grain yield has been recorded in the rank of T₄ > T₅ > T₃ > T₂ > T₆ > T₁. The results indicated that all the treatments with fertilizer recommendation methods gave significantly higher value over the control (T₁) and in terms of straw yield the treatment has been ranked in order of T₄ > T₅ > T₃ > T₆ > T₂ > T₁.

List 2 :Influence of different fertilizer recommendation methods on grain and straw

yields of BRRI dhan29

Treatments	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)
T ₁	4.01c	4.07d
T ₂	5.18b	6.15c
T ₃	5.50ab	6.96b
T ₄	6.29a	7.73a
T ₅	5.81ab	7.08ab
T ₆	5.17b	6.43bc
SE ±	0.4412	0.3373
CV (%)	10.15	6.45

The figure (s) having common letter (s) in a column do not differ significantly at 5% level of significance.




Figure 1 Effect of applying fertilizers through different fertilizer recommendation methods on %grain and %straw yield increase over control of BRRI dhan29

In rice grain, the highest content (1.288%) was observed with BAU Soil Testing Kit (T₄), while all treatments showed significantly higher nitrogen content compared to the control. In case of rice straw, the highest N content of 0.364% was recorded in the treatment T₅ which was statistically similar with T₄. The highest nitrogen uptake in grain and straw was in T₄ (80.86 kg ha⁻¹ and 27.98 kg ha⁻¹). The highest total nitrogen uptake by rice has also found in treatment T₄ (108.84 kg ha⁻¹).

List 3 :Effect urea fertilizer through different fertilizer recommendation methods on nitrogen concentration and uptake by BRRI dhan29

Treatments	Content		Uptake (kg ha ⁻¹)		
	% N (Grain)	% N (Straw)	N (Grain)	N (Straw)	Total
T ₁	1.101d	0.328c	44.20c	13.35e	57.54d
T ₂	1.128c	0.352b	58.37b	21.70d	80.07c
T ₃	1.176b	0.352b	64.82b	24.52bc	89.34bc
T ₄	1.288a	0.362a	80.86a	27.98a	108.84a
T ₅	1.178b	0.364a	68.50b	25.74ab	94.23b
T ₆	1.176b	0.353b	60.76b	22.69cd	83.45bc
SE ±	0.0026	0.0013	4.97	1.22	5.51
CV (%)	0.27	0.44	9.69	6.59	7.89

The figure (s) having common letter (s) in a column do not differ significantly at 5% level of significance.

Phosphorus content varied at different ranges with the application of P fertilizer through different fertilizer recommendation methods in rice grain of BRRI dhan29. The highest phosphorus content in grain and straw of was recorded in the treatment T₃ (0.258% and 0.123%). Phosphorus (P) uptake by rice grain, straw and total uptake has been recorded in the treatment T₄ (15.57 kgha⁻¹, 8.965 kgha⁻¹ and 24.54 kgha⁻¹). All the treatments with fertilizer recommendations gave significantly higher value over the control (T₁).

List 4 :Effect of P fertilizer through different fertilizer recommendation methods on phosphorus concentration and uptake by BRRI dhan29

Treatments	Content		Uptake (kg ha ⁻¹)		
	% P (Grain)	% P (Straw)	P (Grain)	P (Straw)	Total
T ₁	0.242f	0.115d	9.714c	4.679c	14.39c
T ₂	0.255b	0.121b	13.196ab	7.458b	20.65b
T ₃	0.358a	0.123a	14.220ab	8.566a	22.79ab
T ₄	0.248e	0.116d	15.570a	8.965a	24.54a
T ₅	0.249d	0.116d	14.477ab	8.201ab	22.68ab
T ₆	0.254c	0.119c	13.123b	7.650b	20.77b
SE ±	0.00045	0.00052	1.082	0.405	1.273
CV (%)	0.22	0.53	9.9	6.54	7.44

The figure (s) having common letter (s) in a column do not differ significantly at 5% level of significance.

SE (±) = Standard Error of Means, CV (%) = Co-efficient of variance

T₁: Control (No fertilizer), T₂: Farmers practice, T₃: Fertilizer Recommendation Guide (FRG)-2018, T₄: BAU soil testing kit, T₅: Soil Test Basis (STB) and T₆: Rice crop manager (RCP)

Potassium content varied at different ranges with the application of K fertilizer through different fertilizer recommendation methods in rice grain of BRR1 dhan29. The highest potassium content in grain and straw of was recorded in the treatment T₄ (0.339% and 1.383%). Potassium (k)uptake by rice grain, straw and total uptake has been recorded in the treatment T₄ (21.282 kgha⁻¹, 106.89 kgha⁻¹ and 128.17 kgha⁻¹). All the treatments with fertilizer recommendations gave significantly higher value over the control (T₁).

List 5 :Effect of K fertilizer through different fertilizer recommendation methods on potassium concentration and uptake by BRR1 dhan29

Treatments	Content		Uptake (kg ha ⁻¹)		
	% K (Grain)	% K (Straw)	K (Grain)	K (Straw)	Total
T ₁	0.308f	1.227d	12.363d	49.96e	62.32e

T ₂	0.329d	1.290c	17.026bc	79.51d	96.54d
T ₃	0.333c	1.295c	18.354abc	90.19bc	108.54bc
T ₄	0.339a	1.838a	21.282a	106.89a	128.17a
T ₅	0.337b	1.352b	19.594ab	95.60b	115.20b
T ₆	0.315e	1.274c	16.274c	81.90cd	98.18cd
SE ±	0.00067	0.0125	1.405	4.39	5.11
CV (%)	0.25	1.17	9.85	6.40	6.17

The figure (s) having common letter (s) in a column do not differ significantly at 5% level of significance.

SE (±) = Standard Error of Means, CV (%) = Co-efficient of variance

T₁: Control (No fertilizer), T₂: Farmers practice, T₃: Fertilizer Recommendation Guide (FRG)-2018, T₄: BAU soil testing kit, T₅: Soil Test Basis (STB) and T₆: Rice crop manager (RCP)

Sulphur content and uptake varied at different ranges with the application of S fertilizer through different fertilizer recommendation methods in rice grain of BRRI dhan29. The highest Sulphur (S) uptake by rice grain, straw and total uptake has been recorded in the treatment T₄ (14.124 kg ha⁻¹, 17.466 kg ha⁻¹ and 31.590 kg ha⁻¹). All the treatments with fertilizer recommendations gave significantly higher value over the control (T₁).

List 6 :Effect of S fertilizer through different fertilizer recommendation methods on sulphur concentration and uptake by BRRI dhan29

Treatments	Content		Uptake (kg ha ⁻¹)		
	% S (Grain)	% S (Straw)	S (Grain)	S (Straw)	Total
T ₁	0.201e	0.218d	8.068c	8.179e	16.25d
T ₂	0.223c	0.225b	11.540b	13.867d	25.41c
T ₃	0.225b	0.226b	12.401ab	15.739bc	28.14bc
T ₄	0.225b	0.226b	14.124a	17.466a	31.59a

T ₅	0.231a	0.230a	13.372ab	16.264ab	29.64ab
T ₆	0.219d	0.223c	11.317b	14.336cd	25.65c
SE \pm	0.00076	0.00051	0.9296	0.7641	1.377
CV (%)	0.42	0.29	9.65	6.54	6.46

The figure (s) having common letter (s) in a column do not differ significantly at 5% level of significance.

SE (\pm) = Standard Error of Means, CV (%) = Co-efficient of variance

T₁: Control (No fertilizer), T₂: Farmers practice, T₃: Fertilizer Recommendation Guide (FRG)-2018, T₄: BAU soil testing kit, T₅: Soil Test Basis (STB) and T₆: Rice crop manager (RCP)

4. CONCLUSION

The experiment, conducted from February to May 2020, assessed the impact of different NPK fertilizer recommendation methods on rice (BRRI dhan29) in Old Brahmaputra floodplain soils. Employing a randomized complete block design with six treatments, BAU Soil Testing Kit (T₄) demonstrated superior performance, significantly increasing plant height, effective tillers, panicle length, filled grains, 1000-grain weight, grain yield (6.29 t ha⁻¹), straw yield (7.73 t ha⁻¹), and nutrient uptake. The control (T₁) exhibited inferior results in all aspects. Using BAU Soil Testing Kit for fertilizer application emerged as the most efficient and eco-friendly approach, contributing to higher rice yield and improved nutrient uptake compared to other methods, emphasizing the importance of adopting proper fertilizer recommendation methods in sustainable crop production in Bangladesh. Further studies are encouraged to validate these findings. [This should briefly state the major findings of the study. If you are using copy-paste option then select 'match destination formatting' in paste option OR use 'paste special' option and select 'unformatted Unicode text' option]

REFERENCES

- Abedin MJ, Saleque MA, Panullah GM, Majid MA 1998: Evaluation of two concepts of fertilization for wheat in a calcareous soil of Bangladesh. *Journal of Plant Nutrition* **21**(9) 1843-1854.
- AIS (Agricultural Information Service) 2012: *Krishi Dairy*. Agric. Info. Service. Khamarbari, Farmgate, Dhaka. pp. 10-15.
- Akhter Afrin 2002: Response of rice to soil test based fertilizer recommendation. MS Thesis, Department of Soil Science, Bangladesh Agricultural University, Mymensingh.

- Ali MR, Costa DJ, Sayed, MA, Khan MAH, Abedin JA 2009: Development of fertilizer recommendation for the cropping pattern potato-boro-t.aman in irrigated medium highland condition under AEZ -9. *Bangladesh Journal of Agricultural Research* **34(1)** 41-49.
- BARC 2018: Fertilizer Recommendation Guide. Bangladesh Agricultural Research Council (BARC), Farmgate, Dhaka 1215, 274 p.
- BBS 2014: Statistical Yearbook of Bangladesh. Bangladesh Bureau of Statistics. Statistics Division, Ministry of Planning, Govt. of Peoples Republic of Bangladesh. Dhaka. Bangladesh p. 64.
- Beneduzzi HM, de Souza EG, Moreira WKO, Ricardo S, Bazzi CL, Rodrigues M 2022: Fertilizer recommendation methods for precision agriculture – A systematic literature study. *Agricultural Engineering* **42**.
- Betteridge K, Schnug E, Haneklaus S 2008: Will site specific nutrient management live up to expectation. *Research Journal of Agriculture and Forestry Sciences* **4(58)** 283-294.
- Bokhtiar S, Paul G, Alam K, Haque A, Hamidullah, Tirugnanasotkhi 2020: The effects of soil test based potassium application and manures on yield and quality of sugarcane grown on a typical calcareous soil of Bangladesh. *e-ific No.* **27** 9-13.
- Brohi AR, Karaman MR, Aktas A, Savasli E 1998: Effect of nitrogen and phosphorus fertilization on the yield and nutrient status of rice crop grown on artificial siltation soil from the Kelkit River. *Turkish Journal of Agriculture and Forestry* **22(6)** 585-592.
- BRRI 2014: Annual research report. Bangladesh Agril Res Inst, Joydebpur, Gazipur, 22-24.
- BRRI 2020: Year book of agricultural statistics of Bangladesh 2014. Dhaka, Bangladesh: Bangladesh Bureau of Statistics (BBS), Ministry of Planning, Government of the People's Republic of Bangladesh.
- Cheng Wang Da, Cheng Fang-Ming, Wu Wei, Lu JX, Zhao GP, Zhang GP 2000: Effects of NPK application ratios on yield and quality of directly shown late japonica rice. *Acta Agriculturae Universitatis Jiangxiensis*. **22(2)** 174-177.
- Clayton S 2010: "50 Years of Rice Science for a Better World- and It's Just the Start!" Rice Today. International Rice Research Institute, Philippines.
- Csatho P, Arendas T, Fodor N, Nemeth T 2009: evaluation of different fertilizer recommendation systems on various soils and crops in hungary. *Communications in Soil Science and Plant Analysis* **40** 1689-1711.
- Das S, Ali M M, Rahman M H, Khan M R, Hossain A, EL Sabagh A, Barutcular C, Akdeniz H 2018: Soil test based with additional nutrients increased the fertility and productivity of wheat-mungbean-T. Aman rice cropping pattern in the High Ganges River Floodplain soils of Bangladesh. *Bulgarian Journal of Agricultural Science* **24(6)** 992–1003
- Donald GM, Hamblin J 1976: The biological yield and harvest index of cereals as agronomic and plant breeding criteria. *Advances in Agronomy* **28** 361-405.
- Fageria NK, Santos AB, Coelho AM 2011: Growth, yield and yield components of lowland rice as influenced by ammonium sulphate and urea fertilization. *Journal of Plant Nutrition* **34** 371-386.
- FAO (Food and Agriculture Organization) 2006: FAO Production Yearbook, Food and Agriculture Organization, Rome, Italy. pp. 43-55.
- FAO (Food and Agriculture Organization) 2009: FAO Production Yearbook, Food and Agriculture Organization, Rome, Italy. pp. 56-77.
- FAO (Food and Agriculture Organization) 2016: Production, international trade, rice utilization and domestic rice. pp. 11-25.
- FAO (Food and Agriculture Organization) 2016: Production, international trade, rice utilization and domestic rice. *Rice Market Mon* **19(1)** 1-2.

- Hien, NL, Yoshihashi T, Sarhadi, WA 2006: Evaluation of aroma in rice using KOH method, molecular markers and measurement of 2-acetylpyrroline concentration. *Japanese Journal of Tropical Agriculture***50** 190-198.
- IRRI (International Rice Research Institute) 2009: Rough rice production by country and geographical region-USDA. Trend in the rice economy. In: world rice statistics. p. 589.
- IRRI (International Rice Research Institute) 2013: Annual Report Int. Rice Res. Inst. Los Banos, Philippines. p. 179.
- Issaka RN, Buri MM, Nakamura S, Tobita S 2014: Comparison of Different Fertilizer Management Practices on Rice Growth and Yield in the Ashanti Region of Ghana. *Agriculture, Forestry and Fisheries***3** 374.
- Jahan Mahs, Sarkar MAR, Barma NCD, Mondal MNA, Ferdousi MNS 2015: Grain yield, nutrient balance and economics of t. aman rice cultivation as influenced by nutrients management. *Bangladesh Journal of Agricultural Research* **40(1)** 17-34.
- Julfiquar AW, Haque MM, Haque AKGME, Rashid MA 2008: Current status of hybrid rice research and future program in Bangladesh. *Proc. workshop on use and development of hybrid rice in Bangladesh, held at BARC* 18-19.
- Kabir MS, Salam M, Islam A, Sarkar MAR, Mamun M, Rahman M, Nessa B, Kabir M, Shozib H, Hossain M, Chowdhury A, Nasim M, Iftakharuddaula K, Hossain M, Bhuiyan M, Karmakar B, Rahman M, Haque M, Khatun M, Ali M, Rabbi S, Biswas P, Rashid E, Rahman N 2021: Doubling rice productivity in Bangladesh: A way to achieving SDG 2 and moving forward. *Bangladesh Rice Journal* **24(2)** 1-47.
- Kiniry JR, McCauley G, Xie Y, Arnold J G 2001: Rice parameters describing crop performance of four U. S. cultivars. *Agronomy Journal***93** 1354-1361.
- Murugappan V, Latha MR 2001: Nutrient optimization strategy for rice by eliminating surprise nutrient deficiencies. *Madras-Agricultural-Journal***88(4-6)** 291-294.
- Mamun MAA, Haque MM 2016: Ways of managing fertilizers for improved rice production. *Journal of Applied Agriculture and Biotechnology***1** 1-13.
- Mamun MAA, Haque MM, Saleque MA, Khaliq QA, Karim AJMS, Karim MA 2018: Evaluation of different fertilizer management guidelines for boro rice cultivation in south central coastal region of Bangladesh. *Annals of Agrarian Science***16(4)** 466-475.
- Ministry of Finance (MoF) 2015: Ministry of food, *Monitoring Representative* 10-28.
- Momin SI, Husain M 2009: Technology development and dissemination to augment rice production in Bangladesh. *The Guardian*. pp. 33-35.
- Moniruzzaman M, Biswas JC 2013: Water and fertilizer productivity in dry season irrigated rice at farmers' field. *Bangladesh Rice J.***17(1&2)** 64-74.
- Nayak P, Saha MN 1999: Response and effect of N, P and K on rice grain yield and yield attributes under jute-rice-wheat sequence in a gangetic alluvial soil. *Environment and Ecology***17(2)** 467-470.
- Prasad R, Sharma SN, Singh S, Shivay YS 2002: Relative productivity, profitability and NPK removal from soil by some rice-based cropping systems. *Journal of Sustainable Agriculture***19(3)** 31-37.
- Rao AS, Srivastava S, Patil SK 2001: Nutrient mining in agro-climatic zones of Chhattisgarh. *Fertilizer-News*. **46(4)** 109-113.
- Rea RS, Islam MR, Rahman MM, Mix K 2019: Study of nitrogen use efficiency and yield of rice influenced by deep placement of nitrogen fertilizers. Department of Agriculture, Texas State University, San Marcos, USA, Department of Soil Science, Bangladesh Agricultural University, Mymensingh, Bangladesh. *SAARC Journal of Agriculture***17(1)** 93-103.

- Reddy MN, Sitaramayya M, Swamy SN, Sairam A, Kanth GK 1999: Productivity and soil fertility changes under continuous fertilization of rice (*Oryza sativa*)-rice cropping system. *Indian Journal of Agricultural Sciences* **69(6)** 395-398.
- Roy DR, Islam MS, Sarker BC, Tusher TR, Humayun MK, Muliadi, Mamun SA 2016: Impacts of fertilizer application on soil properties at Kaharole upazila of Dinajpur district in Bangladesh. *AGRIVITA Journal of Agricultural Science* **38(2)** 116-125.
- Saidou A, Kossou D, Acakpo C, Richards P, Kuyper TW 2012: Effects of farmers' practices of fertilizer application and land use types on subsequent maize yield and nutrient uptake in central Benin. *International Journal of Biological and Chemical Sciences* **6(1)** 365-378.
- Santhi R, Selvakumari G, Rani P, Perumal R 1999: Fertilizer calibrations for maximum and economic yields of rice in a rice-rice-blackgram sequence on Inceptisols. *Oryza* **36(2)** 136-138.
- Sarkar MIU, Rahman MM, Rahman GKMM, Naher UN, Ahmed MN 2016: Soil test based inorganic fertilizer and integrated plant nutrition system for rice (*Oryza sativa* L.) cultivation in inceptisols of Bangladesh. *The Agriculturists* **14(1)** 33-42.
- Shah AL, Islam MR, Haque MM, Ishaque M, Miah MAM 2008: Efficacy of major nutrients in rice production. *Bangladesh Journal of Agricultural Research* **33(3)** 639-645.
- Siddique MNA, Halim MA, Kamaruzzaman M, Karim D, Sultana J 2014: Comparative Insights for Investigation of Soil Fertility Degradation in a Piedmont Area which Cover the Anjamkhor Union of Baliadangi Upazila, Thakurgoan Bangladesh. *Journal of Environmental Science, Toxicology and Food Technology* **8(4)** 82-87.
- Singh KN, Prasad B, Sinha SK 2001: Effect of integrated nutrient management on a Typic Haplaquent on yield and nutrient availability in a rice-wheat cropping system. *Australian Journal of Agricultural Research* **52(8)** 855-858.
- Singh S, Jain MC 2000: Growth and yield response of traditional tall and improved semi-tall rice cultivars to moderate and high nitrogen, phosphorus and potassium levels. *Indian Journal of Plant Physiology* **5(1)** 38-48.
- Singh VK, Gautam P, Nanda G, Dhaliwal SS, Pramanick B, Meena SS, Alsanie WF, Gaber A, Sayed S, Hossain 2021: A. Soil Test Based Fertilizer Application Improves Productivity, Profitability and Nutrient Use Efficiency of Rice (*Oryza sativa* L.) under Direct Seeded Condition. *Agronomy* **11** 1756.
- Wade LJ, Amarante ST, Olea A, Harnpichitvitaya D, Naklang K, Wihardjaka A, Sengar SS, Mazid MA, Singh G, McLaren CG, Fukai S, Cooper M, Wade LJ 1999: Nutrient requirements in rainfed lowland rice. *Field Crops Research* **64(1-2)** 91-107.
- Wang DJ, Liu Q, Lin JH, Sun RJ 2000: Optimum nitrogen use and reduced nitrogen loss for production of rice and wheat in the Yangtse Delta region. *Environmental Geochemistry and Health* **26(2)** 221-227.