

Study of physiological growth indices of chickpea (*Cicer arietinum* L.) to soil and foliar application through integrated nutrient management practices

ABSTRACT

An investigation was conducted in Rabi season of 2023-24 at the instructional farm of Karunya Institute of Technology and Sciences, Coimbatore to evaluate the performance of physiological indices on chickpea under soil and foliar application. The experiment was established on a Factorial Randomized Block Design (FRBD). The results of the study revealed that morpho-physiological attributes of chickpea were statistically increased by the combined application of soil (NPK 75 per cent + Vermicompost 5 t ha⁻¹) combined with foliar (Nano-urea) nutrient management practices.

Comment [D1]: Please add results briefly to abstract

Keywords: Chickpea, Nano-DAP, Nano-Urea, Crop Growth Rate, Relative Growth Rate, Chlorophyll index

1. INTRODUCTION

Chickpea (*Cicer arietinum* L.) is the second major pulse crop, after common bean (*Phaseolus vulgaris*) in India chickpea is grown predominantly under rainfed conditions. It provides benefits to farming systems that includes lower carbon footprint owing to biological nitrogen fixation to increasing soil health[1]. Chickpea grown in an area of 15.1 million ha (hectares) all over the world, with a total production of 15.8 million tonnes [2]. India and Australia are the top producers contributing together 73 per cent of total area and production [3]. Pulses are cultivated in an area of 27.9 million hectares in India with a total production of 23.1 million tonnes. Chickpea is grown in an area of 13.7 million ha with total production of 10.2 million tonnes with productivity of 1447 kg ha⁻¹ in India and 40.6 lakh hectares of area with 37.67 lakh production and 9.23 lakh productivity in Tamil Nadu[4].

Chickpea can fix N up to 140 kg ha⁻¹ from the atmosphere and receives 80 per cent of its nitrogen (N) needs through symbiotic nitrogen fixation. For succeeding crop, it leaves a sizable quantity of residual nitrogen, and it also contributes as organic matter that helps maintain and improve soil health. The plant nutrients are the major sources for increasing quality and quantity in chickpea. Nutrient availability is one of the major constraints for food production and soil fertility, with unequal utilization of plant nutrients affecting crop growth, development, and yield [5]. Though the sole application of chemical fertilizers results in good crop production, it affects the soil health and status. Therefore, an effective nutrient management can be done by the integration of chemical fertilizers with organic manures such as farmyard manure and vermicompost.

Integrated Nutrient Management (INM) plays a crucial role in agricultural sustainability, encompassing the integration of synthetic fertilizers, organic manure, compost, biofertilizers, and micronutrients. By combining these elements, INM aims to optimize crop yields while minimizing nutrient losses, thereby ensuring agricultural profitability. Among the crop nutrients, nitrogen stands out for its pivotal role in plant growth and development. Consequently, the adoption of foliar application of nutrients emerges as a more efficient strategy compared to traditional fertilization methods. Foliar fertilization involves the direct spraying or application of liquid or water-soluble fertilizers onto plant leaves, facilitating rapid nutrient absorption by bypassing the soil uptake pathway. This supplementary feeding approach proves particularly beneficial during periods of nutrient deficiencies, stress, or accelerated growth, ensuring timely nutrient delivery to support plant health and productivity. Foliar nutrition minimises nutrient loss, boosts bioavailability, and economises crop output by lowering cultivation costs and reducing the quantity of fertilizer applied to crops [6]. Nano- Urea , a new agricultural input based on nanotechnology, has a particle size of 20 to 50 nm, offering a substantially higher surface area than regular urea prills. Liquid Nano- Urea , when directly sprayed onto leaves, enables absorption through stomata, providing crops with a targeted nutrient Nano- DAP , a white liquid fertilizer that provides phosphorous and nitrogen in a 2.5 :1 ratio to plants, improves crop growth and yields. Nano- DAP has great absorption capacity and easily penetrates plant tissues through stomata when used as a foliar spray [8]. Additional nutrient delivery is a critical aspect in boosting grain production in legumes. The degree of bloom drops impacts chickpea production contributing features and yield. The plant's retention of blossoms results in a bigger yield than projected[9]. The present study is aimed to determine the physiological attributes of chickpea through combined application of soil and foliar applications of nano-fertilizers.

2. MATERIALS AND METHODS

A field experiment was carried out at Karunya Institute of Technology and Science, Coimbatore, Tamil Nadu to study the response of chickpea (*Cicer arietinum* L.) to soil and foliar application through integrated nutrient management practices under the Coimbatore region during Rabi season of 2023-24. The climatic conditions under in the Coimbatore district of Tamil Nadu are subtropical. The total rainfall received during the crop-growing period was 128 mm. The weekly maximum and minimum temperatures during the experimental period ranged from 21.0°C to 35.1°C and 5.2°C to 15.1°C, respectively. The soil of the experimental plot was sandy clay loam with low organic carbon (0.42 percent), low in available nitrogen (164 kg ha⁻¹), high in available phosphorus (28.5 kg ha⁻¹), and low in available potassium (235 kg ha⁻¹). The soil reaction of the experimental field was nearly neutral (pH 8.10) with an electrical conductivity of 0.28 dSm⁻¹.

List 1. Treatment details:

A Factor (Soil)	Application
(A ₁)	NPK 100 percent
(A ₂)	NPK 100 percent + FYM 10 t ha ⁻¹
(A ₃)	NPK 100 percent + VC 5 t ha ⁻¹

Comment [D2]: Please add the statistical software you used

Comment [D3]: Refer to the table in the text

(A ₄)	NPK 75 percent + FYM 10 t ha ⁻¹
(A ₅)	NPK 75 percent + VC 5 t ha ⁻¹
B Factor (Foliar)	Application
(B ₁)	Nano-DAP
(B ₂)	Nano-Urea

The experiment was laid out in factorial randomized block design with soil application of NPK 100 percent (A₁), NPK 100 percent + FYM 10 t ha⁻¹ (A₂), NPK 100 percent + Vermicompost 5 t ha⁻¹ (A₃), NPK 75 percent + FYM 10 t ha⁻¹ (A₄) and NPK 75 percent + Vermicompost 5 t ha⁻¹ (A₅) combined with foliar applications, viz., Nano-DAP (B₁) and Nano-Urea (B₂). The chickpea seeds are sown in well-prepared land by dibbling method with seed rates of 90 kg ha⁻¹. A spacing of 30 × 10 cm was adopted. The crop was produced in an irrigated environment with one pre-planting irrigation applied 7 days before sowing and just one lifesaving irrigation. Hand weeding was done between crop rows and rows at 30 to 45 days after sowing. The crops were harvested manually using a sickle, wrapped into bundles with tags from each plot, and sun-dried. Threshing procedures were also carried out treatment-wise manually. The combination effect of different soil and foliar application was significant. This increase in production resulted from increased chickpea crop growth and development with the application of organic and inorganic fertilizers it might be owing to the higher availability of nutrients during crop growth, which eventually boosted growth. The data collected on various characters studied during the experiment were subjected to statistical analysis in a factorial randomized block design (FRBD). The significance of the difference was tested by the “F” test at a 5 percent level.

Crop growth rate

The crop growth rate (g⁻¹m⁻²day) for each specified stage was calculated using the standard formula given by Radford (1967)[10]below:

$$CGR = \frac{W_2 - W_1}{p(T_2 - T_1)}$$

Where,

W₂= Dry weight of crop plant at the time interval T₂

W₁ = Dry weight of crop plant at the time interval T₁

p = ground area occupied by the plant in m²

Relative growth rate

The relative growth rate (g⁻¹g⁻¹day) for each observational stage was worked out by substituting the corresponding dry matter accumulation values of that very stage in the formula was given by Radford (1967)[10]under:

$$RGR = \frac{\text{Log}_e W_2 - \text{Log}_e W_1}{T_2 - T_1}$$

Where,

W₂ and W₁ are the dry matter of plants at the time of T₂ and T₁

Chlorophyll Rate

The chlorophyll content of the crop was measured using an At LEAF Digital Chlorophyll Meter, which was very easy to handle, and the values were recorded quickly and perfect SPAD (Soil Plant Analysis Development) value were maintained properly.

3. RESULTS AND DISCUSSION

3.1. Physiological attributes of chickpea

3.1.1. Crop growth rate ($\text{g}^{-1}\text{m}^{-2}\text{day}$)

The result showed that the effect of ~~the~~ combined application of soil and foliar application with NPK 75 percent + Vermicompost along with Nano- Urea (A_5B_2) resulted in ~~the a~~ higher crop growth rate ~~of~~ 60 DAS to at harvest $2.36 \text{ g}^{-1}\text{m}^{-2}\text{day}^{-1}$. Whereas the lower crop growth rate was observed in the combination of 100 percent NPK with Nano- DAP $0.81 \text{ g}^{-1}\text{m}^{-2}\text{day}^{-1}$ (A_1B_1) (Table 1). The growing trend of CGR might be attributed to increased photosynthetic activity and a favourable response of CGR leads to increased plant population. Similar results were also reported by (Edwardset *al.*, 2005)[11] in maize (Figure- 1). CGR decreases after harvest owing to leaf senescence and a drop in leaf area index.

3.1.2. Relative growth rate ($\text{g}^{-1}\text{g}^{-1}\text{day}$)

The results showed that the effect of combined treatment of soil and foliar application with NPK 75 percent + Vermicompost along with Nano- Urea (A_5B_2) resulted in ~~a the~~ higher relative growth rate at 65 DAS- At harvest ($0.0230 \text{ g}^{-1}\text{g}^{-1}\text{day}$), whereas the lower relative growth rate was observed in the combination of (A_1B_1) 100 percent NPK with Nano- DAP $0.0200 \text{ g}^{-1}\text{g}^{-1}\text{day}$ (Table 1). The relative growth parameters are greatly influenced by the combined application of NPK 75 per cent + Vermicompost with Nano- Urea constitutes a multifaceted approach to augment plant growth and productivity. The treatment enhances nutrient availability, soil structure, and stress tolerance, thereby eliciting profound physiological responses, notably reflected in the RGR of plants. ~~With mMore number of~~ macronutrients, micronutrients, and organic matter, the treatment optimizes metabolic processes, energy utilization, and biomass accumulation. Uniformly released the Nano- Urea sustains nutrient availability throughout the growth cycle. A higher supply of nitrogen boosts plant growth and enhances physiological activities, thereby amplifying the production of growth and yield components (Figure. 2). These result findings were in close agreement with the findings of (Saithejaet *al.*, 2022) [12] on ~~the~~ green gram, (Omranet *al.*, 2018)[13] on mung bean.

3.1.3. Chlorophyll rate

The result showed that the effect of combined treatment of soil and foliar application with NPK 75 percent + Vermicompost along with Nano- Urea (A_5B_2) resulted in higher during peak flowering stage at 65 DAS - At harvest (0.0230), whereas the lower chlorophyll content at peak flowering stage was observed in the combination of 100 percent NPK with Nano- DAP (A_1B_1 -0.0200) (Table 1). The foliar

application of water-soluble fertilizers accelerated crop growth during the early stages, and gradually slowed as the crop matured. This phenomenon may be attributed to the rapid absorption of macro and micronutrients by the crop. Increased total chlorophyll and enzyme activities were observed, leading to enhanced photosynthesis. Due to the heightened leaf thickness, there is a likelihood of increased chlorophyll density within the leaves, thereby potentially sustaining more efficient photosynthesis (Figure. 2). These are similar findings corroborated with Kashiwagi *et al.*, 2010 [14] and Kumari *et al.*, 2018 [15] on sesame.

4. CONCLUSION

The findings of this study indicate that the combined application of soil and foliar NPK 75 percent, Vermicompost, and Nano-Urea exhibited remarkable efficacy in enhancing chickpea growth and productivity. Treatment led to significantly higher crop growth rates, relative growth rates, and chlorophyll rates compared to other combinations. These improvements can be attributed to the multifaceted approach of nutrient supply, enhanced soil structure, and stress tolerance. The sustained release of nutrients from nanoplays played a pivotal role in optimizing metabolic processes and biomass accumulation throughout the growth cycle.

REFERENCES

1. Bohra, A., Pandey, M. K., Jha, U. C., Singh, B., Singh, I. P., Datta, D., ... & Varshney, R. K. (2014). Genomics-assisted breeding in four major pulse crops of developing countries: present status and prospects. *Theoretical and Applied Genetics*, 127, 1263-1291.
2. FAOSTAT, (2021). Food and Agricultural Organization of the United Nations, FAO statistical databases. faostat.fao.org
3. Gaur, P. M., Samineni, S., Thudi, M., Tripathi, S., Sajja, S. B., Jayalakshmi, V., ... & Dixit, G. P. (2019). Integrated breeding approaches for improving drought and heat adaptation in chickpea (*Cicer arietinum* L.). *Plant Breeding*, 138(4), 389-400.
4. Directorate of Economics and Statistics, (2023). Agricultural Statistics at a Glance. Available at <http://desagri.gov.in/>
5. Siddiqui, S. N., Umar, S., Husen, A., & Iqbal, M. (2015). Effect of phosphorus on plant growth and nutrient accumulation in a high and a low zinc accumulating chickpea genotypes. *Annals of Phytomedicine*, 4(2), 102-105.
6. Srivastava, A., & Singh, R. (2023). Effect of Nitrogen and Foliar Spray of Urea and Nano Urea on Growth and Yield of Rabi Maize (*Zea mays* L.). *International Journal of Plant & Soil Science*, 35(18), 2037-2044.
7. Kumar, A., Ram, H., Kumar, S., Kumar, R., Yadav, A., Gairola, A., ... & Sharma, T. (2023). A Comprehensive Review of Nano-Urea vs. Conventional Urea. *International Journal of Plant & Soil Science*, 35(23), 32-40.

Formatted: Font: Italic

8. Maloth, A., Thatikunta, R., Parida, B. K., Naik, D. S., & Varma, N. (2024). Evaluation of Nano-DAP on Plant Growth, Enzymatic Activity and Yield in Paddy (*Oryzasativa* L.). *International Journal of Environment and Climate Change*, 14(1), 890-897.
9. Kamaleshwaran, R., &Karthiga, S. (2021). EFFECT OF FOLIAR NUTRITION ON YIELD AND GROWTH PARAMETERS OF GREENGRAM IN COASTAL AREA OF TAMILNADU (*Vignaradiata*) CV. Vamban 2. *International multidisciplinary e-Magazine*, 1(3), 22-27.
10. Radford, P. J. (1967). Growth analysis formulae-their use and abuse 1. *Crop science*, 7(3), 171-175.
11. Edwards, J. T., Purcell, L. C., &Vories, E. D. (2005). Light interception and yield potential of short-season maize (*Zea mays* L.) hybrids in the Midsouth. *Agronomy Journal*, 97(1), 225-234.
12. Saitheja, V., Senthivelu, M., Prabukumar, G., & Prasad, V. (2022). Maximizing the Productivity and Profitability of Summer Irrigated Greengram (*Vignaradiata* L.) by Combining Basal Nitrogen Dose and Foliar Nutrition of Nano and Normal Urea. *International Journal of Plant and Soil Science*, 34(22), 109-116.
13. Omran, A. H., Dass, A., Jahish, F., Dhar, S., Choudhary, A. K., &Rajanna, G. A. (2018). Response of mungbean (*Vignaradiata* L.) to phosphorus and nitrogen application in Kandahar region of Afghanistan. *Annals of Agricultural Research*, 39(1).
14. Kashiwagi, J., Upadhyaya, H. D., & Krishnamurthy, L. (2010). Significance and genetic diversity of SPAD chlorophyll meter reading in chickpea germplasm in the semi-arid environments. *Journal of food legumes*, 23(2), 99-105.
15. Kumar, A., Singh, S. N., & Khan, M. A. (2018). To know the effect of foliar application of thiourea and potassium nitrate on physiological growth at different stages of sesame (*Sesame indicum* L.). *Journal of Pharmacognosy and Phytochemistry*, 7(2), 2156-2158.

Comment [D4]: All letters are capital. Please correct them

Treatments	Crop growth rate (g ⁻¹ m ⁻² day ⁻¹)			Relative growth rate (g ⁻¹ g ⁻¹ day ⁻¹)			Chlorophyll rate		
	30-45 DAS	45- 60 DAS	60 DAS- Harvest	30-45 DAS	45- 60 DAS	60 DAS- Harvest	At vegetative	At initial flowering	At peak flowering
SOIL APPLICATION									
NPK 100 percent	0.38	0.84	1.21	0.0385	0.0425	0.0213	13.15	31.75	35.40
NPK 100 percent + FYM 10 t ha ⁻¹	0.43	0.94	1.35	0.0395	0.0430	0.0215	14.99	32.30	36.40
NPK 100 percent + VC 5 t ha ⁻¹	0.49	1.09	1.56	0.0395	0.0435	0.0218	15.90	33.40	36.96
NPK 75 percent + FYM 10 t ha ⁻¹	0.56	1.22	1.77	0.0400	0.0435	0.0218	16.00	35.45	37.85
NPK 75percent + VC 5 t ha ⁻¹	0.59	1.29	1.87	0.0400	0.0435	0.0218	16.60	36.77	40.90
S.E (d) ±	0.01	0.03	0.05	0.0002	0.0001	0.0001	0.42	0.23	0.38
CD (p=0.05)	0.03	0.07	0.10	0.0004	0.0002	0.0001	0.89	0.49	0.81
FOLIAR APPLICATION									
Nano-DAP	0.32	0.63	1.03	0.0364	0.0408	0.0204	10.28	28.29	481.29
Nano-Urea	0.66	1.52	2.08	0.0426	0.0456	0.0228	20.38	39.58	643.74
S.E.(d) ±	0.02	0.05	0.07	0.0003	0.0002	0.0001	0.67	0.37	0.61
CD (p=0.05)	0.05	0.12	0.15	0.0006	0.0004	0.0002	1.41	0.77	1.28
INTERACTION (A×B)									
S.E. ±	0.03	0.08	0.10	0.0004	0.0003	0.0001	0.95	0.52	0.86
CD (p=0.05)	0.07	0.16	0.22	0.0009	0.0005	0.0003	1.99	1.08	1.81

Table.1 Effect of Nano-fertilizer on crop physiological growth indices.




Figure. 1 Effect of Nano-fertilizers on crop growth rate




Figure. 2 Effect of Nano-fertilizers on Relative growth rate




Figure. 3 Effect of Nano-fertilizers on Chlorophyll rate