

## Original Research Article

### **Ingrown Toenail Management Across Mental Health and Other Health Facilities in Nigeria: A Review of Anesthetic and Surgical Management**

#### **Abstract**

##### *Background:*

The management of ingrown toenails in Nigeria is notably underdeveloped, posing potentially challenges for patients in mental health facilities and beyond. Addressing this issue requires an interdisciplinary approach and collaboration with other healthcare professionals. Furthermore, there is an important need for a comprehensive review of the treatment guidelines of the anesthetic and surgical techniques utilized in its treatment to improve the quality of patient care.

##### *Methods:*

We employed a structured narrative approach to identify and review available articles on ingrown toenail management in mental health and other health facilities in Nigeria. We aimed to evaluate the anesthetic and surgical techniques utilized in managing these patients.

*Results:* No documented treatment for ingrown toenails was identified among patients in mental health facilities across Nigeria. While various surgical techniques are employed for management in other health facilities, the specific anesthesia techniques utilized are unclear.

*Conclusions:* A unified management protocol/guideline for ingrown toenails and developing podiatry training in Nigeria are imperative. Developing and implementing an effective treatment protocol improves the quality of care and widens coverage to all patients. In addition,

establishing podiatry training across educational facilities could bridge this gap in ensuring optimal care for these patient populations.

**Keywords:** anesthesia, ingrown, mental, Nigeria, surgical, toenails

## Introduction

An ingrown toenail, or onychocryptosis, usually occurs when the nail plate punctures the periungual skin, causing substantial pain, inflammation, discomfort, and increased risk of infection if left untreated.<sup>[1]</sup> It is reported that ingrown toenails occur commonly in the hallux, presenting most often in teenagers and early adults.<sup>[2]</sup> However, it can present at any age, with several factors implicated as the possible contributory factors.<sup>[3]</sup> These factors include poor nail-cutting technique, hyperhidrosis, ill-fitting footwear, nail deformity, trauma, obesity, and peripheral edema. Patients with ingrown toenails might present late when the infection has set in.<sup>[4]</sup> It is exacerbated in individuals with mental health disorders, who may sometimes neglect seeking healthcare for other coexisting conditions.<sup>[5]</sup>

Ingrown toenails cause severe discomfort and pain in patients. Hence, it is a common reason for clinic consultations in Nigeria. An ingrown toenail is classified into three stages.<sup>[6]</sup> While the early stages can be managed conservatively, the late stage requires surgical treatment and partial or complete avulsion of the nail with or without nail matrix ablation. Ablation of the nail bed can be carried out chemically,<sup>[7]</sup> using laser,<sup>[8]</sup> surgically,<sup>[9]</sup> or electrocautery.<sup>[10]</sup> Chemical matricectomy has the lowest recurrence rate but has an increased risk of wound infection and prolonged healing time.<sup>[7]</sup>

Patients with mental health conditions sometimes appear physically inactive, which could be attributed to the side effects of some psychiatric medications.<sup>[11]</sup> However, some of the patients have co-existing undiagnosed ingrown toenails, which might be limiting their physical activities due to pain. The treatment choice for ingrown toenails varies based on the stage of the disease. However, the most significant factor in treatment choice stems from the available specialty. Podologists and pedicurists favor conservative management procedures using braces and similar devices, while some dermatologists use other noninvasive approaches, and the majority tend to perform surgery.<sup>[12]</sup>

In Nigeria, patients with ingrown toenails often present to surgeons rather than podiatrists, podologists, pedicurists, and dermatologists. While there are limited podiatrists and podologists in Nigeria, the few pedicurists and dermatologists do not usually treat ingrown toenails. As such, the most typical treatment method is surgery. The surgeries are amenable to various anesthetic techniques. Kim and colleagues<sup>[13]</sup> discussed using ultrasound-guided popliteal nerve block using short-acting lidocaine in the surgical treatment of ingrown toenail. However, digital blocks with local anesthetic agents without epinephrine are also employed.<sup>[14]</sup> Classical V-shaped and H-shaped blocks with anesthetic agents have also been described.<sup>[15]</sup>

This review aims to identify, review, and summarize existing literature on ingrown toenail management across mental and other health facilities in Nigeria. Our goal is to understand common anesthetics and surgical techniques and present a broad perspective on ingrown toenail management in Nigeria across various patients.

## **Methods**

A narrative approach was used to synthesize, interpret, and critique the findings of studies that reported the management of ingrown toenails across a range of patients in Nigeria. This approach was chosen in view of the paucity of literature on the subject in Nigeria. However, the review provides a framework for further studies and recommendations for a unified management protocol for ingrown toenails in Nigeria.

We conducted an electronic search of PubMed, Google, and Google Scholar for peer-reviewed, English-language articles published up until March 2024. Preliminary keyword searches included combinations of “anesthesia,” “ingrowing toenail,” “management,” “mental,” “Nigeria,” and “surgery.”

We limited our review to Nigeria, given the under-reporting of the existing modality for the management of ingrown toenails among a broad range of patients in Nigeria, including patients in mental health facilities. Furthermore, the training institutions have not developed specialty training in podiatry and podology as obtained in other parts of the world.

## **Results**

Our search resulted in 5 articles in total. During our search, we observed a notable absence of reported treatment for ingrown toenails provided to patients in mental health facilities in Nigeria. In the context of the paucity of literature on the subject in Nigeria, the narrative review aims to identify a few studies that describe the problem of interest. As seen in Table I, the identified studies fall under descriptive retrospective studies, including case reports, case series, and cross-sectional studies. Two hundred and eleven patients treated for ingrown toenails were reviewed, and the treated patients were reported using different methodologies. In a retrospective study<sup>[16]</sup> involving 52 patients, the anesthetic technique employed was digital block with 1% plain

Xylocaine and tourniquet. No anesthetic technique was mentioned in the case report <sup>[17]</sup>, case series<sup>[18]</sup>, and cross-sectional study.<sup>[19]</sup> A prospective study<sup>[20]</sup> among those reviewed mentioned that the procedure was done under local anesthesia, no specific anesthetic technique was mentioned. The surgical management employed for the early stage of the disease was a conservative approach in all the studies reviewed. However, there was a twenty-five percent recurrence in those managed conservatively within 1 year in the prospective study. Matricectomy was employed as the treatment of choice in stages II and III, while wedge resection plus partial matricectomy was used in the case series study.<sup>[18]</sup>

**Table I: Articles on ingrown toe nail management in Nigeria**

	<b>Surgical Method</b>	<b>Anesthesia</b>	<b>Patients Enrolled</b>	<b>Methodology</b>	<b>Findings</b>
Nwagbara <i>et al</i> 2022 <sup>[19]</sup>	Partial or complete great toenail excision plus debridement	Not stated	99	Cross sectional prospective study	Surgical treatment provides a better outcome with minimal recurrence for late stage or after failed conservative management.
Ogbeter and College 2020 <sup>[17]</sup>	Matricectomy	Not stated	1	Case report	Surgically treated medial and lateral folds of both halluces.
Benjamin <i>et al</i> 2017 <sup>[20]</sup>	Late: Suppuration, Ulceration, hypertrophic nail fold-Wedge resection, surgical matricectomy under local anaesthesia  Severe: Simple Nail Avulsion	Local Anesthesia-nonspecific	28	Prospective study	Patients (25%) who had conservative treatment and avulsion of the nail had recurrence.

Olakulehin et al., 2015 <sup>[18]</sup> .	Based on Heifetz staging and recurrence status, Stage II and III had Wedge resection of either or both medial and lateral toenail folds plus partial matricectomy.	Not stated	55	Case series	Within 1 year period, there were 2 recurrences out of 63 halluces treated in Stage II and III with Wedge resection plus partial matricectomy. In all, a 99% cure rate was recorded.
Akhator, 2011 <sup>[16]</sup>	Wedge resection of affected nail and matrix; total excision of total nail and matrix were done for late stages (stage 3). Earlier stages were managed conservatively.	Digital block with 1% plain Xylocaine and tourniquet	52	Retrospective study	After a 1-year period, there was no recorded recurrence. No recurrence in cases managed conservatively also.

## Discussion

In mental health facilities across Nigeria, there is a notable absence of reported treatment for ingrown toenails provided to patients. Research consistently demonstrates the heightened rates of morbidity and mortality among individuals struggling with mental illness.<sup>[21,22]</sup> The high incidence of undiagnosed medical conditions and the subsequent neglect of physical health concerns raise further alarm.<sup>[23]</sup> Compounding these challenges, some patients with mental health disorders in Nigeria are subjected to leg restraints,<sup>[24]</sup> elevating the risk of trauma and injuries to the lower extremities which is a risk factor for ingrown toenails. Delays in diagnosing and treating ingrown toenails can result in severe complications, potentially necessitating amputation.<sup>[25]</sup> Urgent measures are warranted to enhance the screening for foot disorders in mentally ill patients, thereby mitigating the prevalence of comorbidities within this vulnerable population.

Many patients in the index study had surgeries as the treatment modality. However, there was no clear description of the anesthetic techniques employed in most cases. Elsewhere, the anesthetic

techniques were well-defined. Kim and colleagues<sup>[13]</sup> talked about the use of short-acting lidocaine in ultrasound-guided popliteal nerve block for ingrown toenail surgical therapy. Nevertheless, digital blocks containing non-epinephrine local anesthetic drugs are also utilized.<sup>[14]</sup> This is concordance with one<sup>[16]</sup> of the studies in Nigeria describing the management experience of ingrown toenail. A digital block with 1% plain Xylocaine and a tourniquet was employed to achieve adequate anesthesia. There have been descriptions of traditional V- and H-shaped blocks containing anesthetic drugs.<sup>[15]</sup>

The peculiar observation in the surgical management of the patients in the index study is that orthopedic surgeons are the health professionals in Nigeria handling cases of ingrown nails. Practitioners who perform surgery on ingrowing toenails include orthopedic surgeons, general practitioners, podiatrists and podiatric surgeons.<sup>[26]</sup> The podiatric surgeons have recently made advances, employing minimally invasive surgery to treat ingrown toenail.<sup>[27]</sup> The health professional training is yet to develop a training program for pedology/podiatry in Nigeria. In terms, of the surgical technique, the surgeons in Nigeria adopt the conservative approach for the initial stage of ingrown toenails and matricectomy for the advanced stages in keeping with other health professionals outside Nigeria treating ingrown toenail.<sup>[28]</sup>

### **Limitations**

The paucity of literature on the topic constrains the scope of this review, while the relatively small patient populations included in various studies restrict the generalizability of findings. There is a concern that the management of ingrown toenails in Nigeria appears under-diagnosed and under-reported in healthcare facilities, potentially overlooked, particularly among individuals with mental health disorders.

## Conclusion

This paper provides an overview of the current management practices for ingrown toenails across various patients in Nigeria. Despite the scarcity of literature on the subject, it highlights the non-specific anesthetic techniques during surgical treatment. Notably, Nigeria appears not to have a well-developed management protocol for ingrown toenails. However, implementing a well-developed treatment protocol promises enhanced care and broader coverage for all patients, including those in mental health facilities.

There is a pressing need for policy initiatives to enhance mental health services in Nigeria, focusing on delivering comprehensive care to patients. This should encompass preventive and curative measures for lower extremity conditions, including ingrown toenails. It is imperative to establish clear guidelines regarding the use of anesthetic techniques during surgical interventions for ingrown toenails.

Furthermore, this review underscores the necessity for the Council of Medical Education in Nigeria to recognize the importance of establishing podiatry training and services in the country. Currently, orthopedic surgeons bear the brunt of managing ingrown toenails. Establishing podiatric surgery would alleviate this burden and ensure more specialized and high quality of care for patients with foot-related issues.



## REFERENCES

1. Bryant A., Knox, A. Ingrown toenails: the role of the GP. *Aust Fam Physician*. 2015; 44(3):102–5
2. Khunger N., Kandhari R. Ingrown toenails. *Indian J Dermatol Venereol Leprol*. 2012; 78(3):279–89.
3. Eekhof J.A., Van-Wijk B., Knuistingh Neven A., Van-der Wouden, J. Interventions for ingrowing toenails. *Cochrane Database Syst Rev*. 2012; (4): CD001541.
4. Coll C., Beech I. A rare complication following a Winograd Procedure: A case report and review of National sequellae, *Fastrc*. 2022; 2 (1):100131, 2667–3967.
5. Hahm H.C., Segal S.P. Failure to seek health care among the mentally ill. *Am J Orthopsychiatry*. 2005; 75(1):54–62.

6. Chabchoub I, Litaïem N. Ingrown Toenails. [Updated 2022 Sep 18]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK546697/>
7. Shaikh F.M., Jafri M., Giri S.K. Keane R. Efficacy of wedge resection with phenolization in the treatment of ingrowing toenails. *J Am Podiatr Med Assoc.* 2008;98(2): 118–22.
8. Ozawa T., Nose K., Harada T., Muraoka M., Ishii M. Partial matricectomy with a CO<sub>2</sub> laser for ingrown toenail after nail matrix staining. *Dermatol Surg.* 2005; 31(3): 302–5.
9. Hassan R E, Khan L, Shah S, et al. Surgical Strategies for Ingrown Toenails: A Comprehensive Review of Techniques, Outcomes, and Advancements. *Cureus* 2024; 16(1): e52501.
10. Selmy A, Waleed A, Saleh A., Khaled A., Soliman G., Mohamed S. Electrocautery with partial matrixectomy in comparison with partial matrixectomy alone in treatment of ingrown toenail. *Egypt J Surg* 2020; 39(4): 1095-1102
11. Vancampfort D., Firth, J., Schuch, F.B. et al. Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. *World Psychiatry.* 2017; 16(3):308-315.
12. Haneke E. Controversies in the Treatment of Ingrown Toenails. *Dermatol Res Pract.* 2012. Article ID 783924, doi:10.1155/2012/783924
13. Kim B.S., Kim K., Day J., Seilern-Und A.J., Kim, J. Ultrasound-Guided Popliteal Nerve Block with Short-Acting Lidocaine in the Surgical Treatment of Ingrown Toenails. *Int J Env Res Pub Health.* 2021; 11;18(10):5059. doi: 10.3390/ijerph18105059.
14. Pandhi D., Verma P. Nail Avulsion: Indications and Methods (surgical nail avulsion). *Indian J Dermatol VenereolLeprol.* 2012; 78: 299-308

15. Sánchez H.S., Giralt de-Veciana E. Comparison of 2 Anesthetic Techniques in Onychocryptosis Surgery of the Great Toe: A Randomized Clinical Trial, *Actas Dermo-Sifiliográficas*, 2022; 113, 4: T370-T375.
16. Akhator A. Surgical Matricectomy in the Management of Ingrown TOE Nail-Experience from Nigeria. *Biosci. Biotechnol. Res. Asia* 2011; 8(2): 603-605.
17. Ogbetere F.E., Erhuanga G.O. Onychocryptosis affecting the lateral and medial folds of both big toes: A rare case report and review of the literature. *Afr J Biol Med Res.* 2020; 3(3): 127-132.
18. Olakulehin O.A., Oluwadiya K.S., Akanbi O.O., Babalola O.R. A Physiological Approach to Manage Ingrown Toenails. *Sch J Appl Med Sc.* 2015; 3(1D):272-275
19. Nwagbara V.I., Ashindoitiang J., Asuquo I.M. Ingrown toenail (onychocryptosis or unguis incarnatus): Pattern and treatment outcome among patients seen in a secondary health facility over a three-year period. *Nig Q J Hosp Med.* 2022; 31:1-2
20. Benjamin T.I., Amefula T.E. Ingrown toenail, as seen in Bayelsa State, Nigeria. *Int J Adv Med.* 2017; 4(3):614-619.
21. Momen NC, Plana-Ripoll O, Agerbo E, et al. Mortality Associated with Mental Disorders and Comorbid General Medical Conditions. *JAMA Psychiatry.* 2022;79(5):444–453
22. Fiorillo A., Sartorius N. Mortality gap and physical comorbidity of people with severe mental disorders: the public health scandal. *Ann Gen Psychiatry* 2021; 20, 52
23. Druss B.G., Bradford D., Roseheck R., Radford M.J., Krumholz H.M. Quality of medical care and excess mortality in older patients with mental disorders. *Arch Gen Psychiatry*, 2001; 58, 565–572.

24. Omolayo A., Wada Y.H. Impact of traditional healers in the provision of mental health services in Nigeria. *AnnMed Surg.* 2022; 82: 104755, 2049-0801
25. Gupta A.K., Mays R.R. The impact of Onychomycosis on quality of life: a systematic review of the available literature. *Skin Appendage Disord.* 2018; 4(4):208–216.
26. Terrill A.J., Green K.J., Salerno A. *et al.* Risk factors for infection following ingrowing toenail surgery: a retrospective cohort study. *JFAR* 2020; 13, 48.
27. Ergün T., Korkmaz M., Ergün D., Turan K., Muratoğlu O.G., Cabuk H. Treatment of Ingrown Toenail with a Minimally Invasive Nail Fixator: Comparative Study with Winograd Technique. *J Am Podiatr Med Assoc.* 2024; 114(1):22-024.
28. Exley V., Jones K., O'Carroll G., Watson J., Backhouse M. A systematic review and meta-analysis of randomized controlled trials on surgical treatments for ingrown toenails part I: recurrence and relief of symptoms. *JFAR.* 2023; 10;16(1):35.