

MESO COMPOUNDS SYSTEMATIZATION –A CHEMICAL PARADOX AMONG INOSITOLS AND A GROUP OF SUPER-SYMMETRIC COMBINATIONS

ABSTRACT. *meso* Compounds have been defined by themselves, as well as by their relationship with another group of diastereomeric combinations, *C₂symmetrical* ones, isomer to *meso*. Every *meso* compound should present a homodimer equivalent able to display at least one *C₂symmetrical* isomer. Cahn-Ingold-Prelog test, however limited, is also valuable. A systematization of *meso* combinations includes the following types: (A) *meso* Homodimers possessing one *super-symmetrical* isomer or two real or envisaged *C₂symmetrical* enantiomers; (B) *meso* Heterodimers with a binding support having geometric or optical symmetry (e.g. >CR₂, etc.). (C) *meso* Heterodimers with a binding support devoid of symmetry (of the form >CRR', etc.), seeable as *meso*, and provable in this way not by themselves but by their equivalent *meso* homodimers. A paradoxical behavior has been shown at one of inositol isomers, in comparison with the others. Moreover, a new group of compounds has been disclosed, which are concomitantly *meso* and *C₂symmetrical*, and of this reason they have been called *super-symmetrical*.

Keywords: chemical paradox; *meso*; *C₂ symmetrical*; mirror plane of symmetry; linking support (matrix); homodimers; heterodimers; inositols; *super-symmetrical*.

1. Introduction

Chemical symmetry includes three types: *meso* [1], *C₂symmetrical* (*C₂symm.*) [2] and *centrosymmetrical* [3], and all operates with chiral carbons, contrary to geometrical symmetry that works with achiral carbons, and this property is decided by geometrical and not by chemical concepts. The molecule of a *meso* compound is formed either of two chiral enantiomeric halves uniformly linked with each other (homodimers) or linked on a mono- or polyatomic support (heterodimers). (The term uniform indicates that the linkage is made between the atoms of the same rank. A vast group of dimers, isomers to *meso* and *C₂symm.* is known, whose identical chiral halves are linked in a non-uniform manner [4]. The linking support can be symmetric (of the >CR₂ form, etc.) or non-symmetric (of the >CRR' form, etc.). *meso* Compounds are characterized by their mirror plane of symmetry and they are distinguished by the fact that in homodimers it intersects only bonds, while in heterodimers it intersects one or more atoms. When the two enantiomeric halves of *meso* combinations are alternatively dimerized, one or two *C₂symm.* isomers are produced, most frequently optically active and enantiomeric. Among the isomers of inositol, only one was found to produce two chiral, *C₂symm.* enantiomers. Alternative dimerization of the two halves of *meso* heterodimers includes the binding support when it is symmetric. However, duplication is preceded by removing of linking support when it is devoid of symmetry. The molecule of *C₂symm.* compounds is formed of either two identical chiral halves uniformly linked with each other or linked on a binding support, invariably of the >CR₂ form. Rotation of a *C₂symm.* compound by 180° around a defining axis arrives at the same structural context as initially. In this paper, the analysis of *meso* homo- and heterodimers when confronted with the above concepts has been made.

2. On the Planes of Symmetry

The plane of symmetry of 1,2-cis-dimethylcyclobutane cuts two bonds. Since at least one isomer, 1,2-trans-dimethylcyclobutane ($C_{2symm.}$), has been prepared [5,6] (Fig. 1), one can assert that the plane of symmetry of 1,2-cis-dimethylcyclobutane is a mirror plane of symmetry, and 1,2-cis-dimethylcyclobutane is a *meso* compound. The analysis of symmetry of cis- and trans-1,3-cyclobutane dicarboxylic acids [7] indicates the following. cis-1,3-Cyclobutane dicarboxylic acid has two planes of symmetry, one intersecting C-2 and C-4, and the other C-1 and C-3, but only the first one separates two enantiomeric halves. Rather as final results than a




Figure 1. A comparison of planes of symmetry of some disubstituted cycloalkanes.

general procedure, the two planes of symmetry have different ranks: the first one is a veritable chemical plane of symmetry and the other is a non-chemical, i.e. a geometrical plane of symmetry. trans-1,3-Cyclobutane dicarboxylic acid has no mirror plane of symmetry, consequently it is not *meso*; the trans isomer is in fact $C_{2symm.}$ However, cis,trans,trans-1,2,3,4-tetracarboxy-cyclobutane [(1 β ,2 β ,3 α ,4 β)-1,2,3,4-tetracarboxy-cyclobutane] [8] (see Fig. 6) is a *meso* heterodimer: its mirror plane of symmetry intersects C-1 and C-3 and some of the atoms linked to them. The same results are obtained with cis- and trans-1,3-dimethyl-cyclobutane [7] as well as with cis-1,4-dicarboxy-cyclohexane [9] and cis-1,5-dimethyl-cyclooctane [10]. C-1 and C-6 in 1,6-dimethyl-cyclodecane binds and shares a proton and the whole molecule becomes a carbocation [11].

(A) *meso* Homodimers and their $C_{2symmetrical}$ Isomers

Numerous results suggest that every *meso* homodimer (Fig. 2) presents at least one $C_{2symm.}$ isomer. Fischer and Hertz [12] reduced galactaric acid (a *meso* homodimer) with Na-amalgam and obtained a racemic mixture of D- and L-galactonic acid. They separated the two enantiomers of the mixture as salts of strychnine. Of every enantiomer they prepared the matching aldohexose, i.e. D- and L-galactose. As expected, both aldohexoses gave the same galactitol or galactaric acid, by reduction with Na-amalgam or oxidation with nitric acid, respectively. By this experiment, Fischer proved that the molecule of *meso* homodimers is formed of two enantiomeric halves. On the other hand, reduction of dextrotartaric acid with HI produced a single species of malic acid, that is D isomer [13]. Fischer had previously made similar experiments on D-mannitol: by oxidizing it kinetically, only one type of aldohexose, D-mannose, was obtained [14]. By these experiments, Fischer discovered a group of compounds whose molecule is formed of two identical chiral halves. Their distinct features were also noticed by Vickery [15]. Subsequently, these compounds have been gathered under the group called $C_{2symm.}$ [2,16-18]. NMR technique confirmed the perfect identity of the component halves of $C_{2symm.}$ homodimers: only half of the expected signals appeared [19].




Figure 2. *meso* Homodimers liable to present **chiral enantiomeric** C_2 *symm.* isomers.

Imaginary dimerization of the (R)-half of tartaric acid would produce (R,R)-tartaric acid, while (S)-half would give (S,S)-tartaric acid. A similar procedure applied to galactitol would produce D- and L- iditol. Other triads in carbohydrate chemistry are erythritol, (R,R)-threitol, (S,S)-threitol; allitol, D-mannitol, L-mannitol, as well as their aldaric acids. L-Cys-D-Cys forms a triad with L-Cys-L-Cys and D-Cys-D-Cys, and numerous triads of 2,5-diketopiperazines are known. At least seven triads of carotenoids of the type *meso*-zeaxanthin, (R,R)-zeaxanthin, (S,S)-zeaxanthin, have been found. Two triads of lignans are based on nordihydroguaiaretic and dihydroguaiaretic acids, and one of alkaloids includes *meso*-, (+)- and (–)-chimonanthine isomers. Diolmycin B2 is the C_{2symm} . isomer of (2S,3R)-diolmycin B1 (phenols), and daibudilactone B of daibudilactone C (terpenoids) [4]. As can be noticed, all C_{2symm} . isomers are optically active. Exposure of the (Z)-isomer of 2-(3,4,5-trimethoxyphenyl)-4-(3,4-methylenedioxyphenyl)-4-oxo-2-butenonitrile (β -cyanochalcone) in the solid state to sunlight led by dimerization to a symmetrical Z-Z dimer. Exposure of the (E)-isomer (4E) to the same conditions determined an unusual dimerization and produced an unsymmetrical E-Z dimer [20]. The structure of both dimers was established by X-ray crystallographic analysis. This diversity indicates the possibility to produce also a C_{2symm} . isomer. Synthesis of 52 and 53 *meso* diepoxy derivatives produced also two C_{2symm} . isomers, 51 and 54 [21]. The two C_{2symm} . isomers of *meso* isomer 176 [22] are known [23]. The configuration of chiral centers of 2,3-butanediols was correlated with the enantiomers of mannitol [24]. L,L-, L,D- And D,D-2,5-diketopiperazines of tryptophanes are known [25,26], and this opens the possibility for the synthesis of C_{2symm} . isomers of dimethylfelltanine A [27], felltanine A, felltanine C [28], felltanine D [29], dragmacidin B, trans-dragmacidin C [30]. An isomer of phenazostatin D, phenazostatin B, whose molecule is formed by dimerization of a half of phenazostatin D, is also known [31]. Structure comparison of *meso* cyclobutane derivatives endiandrin B, cinbalansan and heterotropan [32-34] indicate a metabolic relationship between them. On the other hand, endiandrin A corresponds to endiandrin B, di-O-methyl-endiandrin A to cinbalansan, and magnosalin to heterotropan, all as C_{2symm} . isomers. Some of the isomeric truxillic and truxinic acids have to be treated as homodimers and others as heterodimers [3]. β -Truxinic and ω -truxillic have a plane of symmetry intersecting two bonds and their pairs C_{2symm} . are δ - and μ -truxinic, respectively [35,36]. β -Truxinic acid is related also with β -N,N'-dimethyl truxindianilide [37] and caracasandiamide [38]. α -Truxillic acid is centrosymmetric [39]. γ -, epi-, ϵ - And peri-truxillic acids [3] have planes of symmetry intersecting atoms, hence their equivalent *meso* homodimer are (R,S)-tartaric acid or (R,S)-2,3-diphenylbutane. Both isochaejasmene (*meso*) and its isomer chaejasmene (C_{2symm} .) have been discovered [40]. All the above mentioned C_{2symm} . isomers are optically active. There isn't any doubt to question the possibility to construct chiral C_{2symm} . isomers of 2a and 12 [41], *meso* dimer [42], *meso* cyclohexene derivative [43], 4d [44], 1 [45], thymoquinone [46], *meso* DKP [47], and 72, 73, 75 [1] (Fig.2).

(B) *meso* Heterodimers with a binding support of the >CR₂ form, accompanied by at least one $C_{2symmetrical}$ isomer

meso Compounds based on a binding support of the >CR₂ form allow the existence of chiral C_{2symm} . isomers (Figs. 3 and 4), although they are not rigorously homodimers. E.g. all three asarolignans, as well as some of their isomers are known [48]. About 15 triads have been found especially among natural compounds: 2 carbohydrates, 2 amino acids, 5 lignans, 4 phenols, 2 polyols [4]. The things are also prepared for trehalose: since ent-glucose (L-glucose) has been synthesized [49,50], the following triad might be envisaged: trehalose (α -D-Glcp-1,1- α -D-Glcp), *meso*-trehalose (α -D-Glcp-1,1- α -L-Glcp), ent-trehalose (α -L-Glcp-1,1- α -L-Glcp). 3-Deoxyxylitol [51] or 3-deoxyribitol [52] with the two enantiomeric 3-deoxyarabinotols [53] should be added to carbohydrate list. L,D-Diaminopimelic acid and *meso* lanthionine are components of two triads of amino acids. 3,3'-Didemethoxynectandrin B, isonectandrin B, zuonin B and nectandrin B are *meso* isomers of lignans triads. Neolignans are represented by three asarolignans. Two *meso* hybocarpone, (3S,5R)-octahydrocurcumin and (3R,5S)-hannokinol are *meso* constituents of




Figure 3. *meso* and C_2 symm. isomers of asarolignans and neolignans [48].

phenol triads [4]. Where known, all C_2 symm. isomers are chiral.

Numerous heterodimers are known (Fig. 4) and their classification has been made:

- Of 3-deoxyxylitol or 3-deoxyribitol type: cis-1,2-dimethylcyclopropane [54], 1 [55], 15-18 [21], L,D-Diaminopimelic acid [56], 20, 22-24 [1], 40a, 42a, 83a [22], 2e, 2f [57], 23 [45], 9 [58], 2 [59], cyclopentenederivative [43], eurorubrin [60], daibudilactone D [61].
- Similar to 3-deoxy-3-ketoxylitol: 11 [62], 209a, 215b, 215c [22], cuscohygrine [63].
- Of *meso* 2,5-dimethyltetrahydrofuran type [64]: 14, 16 [65], 2 [66], 83 [67], glabrescol, teurilene [68], 50, 54 [22].
- Similar to cis-2,3-dimethylaziridine [69]: 36 [22].

(C) *meso* Heterodimers with a Binding Support Devoid of Symmetry (>CRR', etc.), Seeable as *meso*, and Provable in This Way not by Themselves but by their Equivalent *meso* Homodimers

Let's make the following theoretical experiment: the middle bond of erythritol [70] is broken and the two halves are linked on the >CHOH residue. It is obvious that xylitol or ribitol are produced, hence erythritol is the *meso* homodimer equivalent for both (Fig. 5). In this manner, by uniformly linking two enantiomeric halves to an arbitrary binding support, an unlimited number of compounds could be produced. The following derivatives of xylitol have been prepared: 1,5-anhydro-xylitol, 2,3,4-tri-O-acetyl-1,5-anhydroxylitol [71], 1,5-diacetyl-3-acetoxymethyl-2,4-methylene-xylitol and 2,4-methylene-xylitol [72]. Schmidt and Lieberknecht [73] synthesized *meso*-2,5-anhydro-3,4-isopropylidene-allaric acid. The latter compounds showed that the binding support can be either unitary or in a fragmented state (Fig. 5). Many *meso* heterodimers are produced by indirect paths, either chemically or biochemically, hence all of them have to be investigated *per se*, and not historically. Moreover, there is no connexion between the structure of the two enantiomeric halves, on one hand, and the binding support (Figs. 5-8).

1. Rules for finding *meso* homodimers equivalent to non-provable *meso* heterodimers. A number of rules have been elaborated for removing the binding support of *meso* heterodimers and finding their equivalent *meso* homodimers (equivalent, not isomer!). Contrary to *meso* heterodimers, the *meso* equivalents are able to present at least one C_2 symm. isomer. As all rules, the beyond rules will be better understood from their applications. In chemical terms, removing of linking support of *meso* heterodimers resembles to a surgical operation of medicine, probably. Here are some rules:

- When cut by the mirror plane of symmetry, an oxygen atom is equivalent with two hydroxy groups, a nitrogen atom with two amino, etc.
- A carbon apex is equivalent with two methyl groups or it is simply removed.
- Chiral atoms of *meso* heterodimers are carefully preserved in the obtained equivalent homodimers, by the analysis of their substituents.

If oxygen atom in cis-2,3-dimethyl-oxirane (Fig. 6) [74] is replaced by two hydroxy groups, (2R,3S)-butanediol is obtained. In cis-trimethyl-cyclopropane and trans-trimethyl-cyclopropane [75], the apex intersected by the mirror plane of symmetry cannot be replaced by two methyl groups since the chirality of the other two cyclopropane groups is annihilated. Consequently, that apex would be replaced by two imaginary R residues. In triethyl trans-cyclopropane-tricarboxylate [76] and triethyl trans-1,2,3-tricyano-cyclopropane-tricarboxylate [77],




Figure4. Heterodimers with symmetric binding support, able to form C_2 symm. isomers.




Figure 5. Bindingsupportof *meso* heterodimers can be also in a fragmented state.

replacement of the suitable apex by two methyl groups would produce *meso* dimethyl- and *meso* dimethyl-dicyano-succinic acid, hence the chiral character of the two carbons is preserved.

Four isomers are possible for tetrasubstituted cyclobutane with the same substituent, and all four are known [8,78-79]. Two of them – cis,cis,cis-1,2,3,4-tetracarboxycyclobutane and cis,trans,cis-1,2,3,4-tetracarboxy-cyclobutane – have mirror plane of symmetry intersecting bonds only, hence they have to be considered *meso* homodimers. The other two, cis,trans,trans-1,2,3,4-tetracarboxy-cyclobutane and trans,trans,trans-1,2,3,4-tetracarboxy-cyclobutane have to be treated as *meso* heterodimers. The *meso* homodimer of the latter two is *meso* tartaric acid. cis,trans,cis-1,2,3,4-Tetracarboxy-cyclobutane is *meso* and concomitantly it is the C_2 *symm.* isomer of cis,cis,cis-1,2,3,4-tetracarboxycyclobutane. Of this reason, we suggest the term *super-symmetric* for cis,trans,cis-1,2,3,4-tetracarboxy-cyclobutane and for similar compounds.

Four isomers are known for pentahydroxy-cyclopentane, and all are *meso* heterodimers [80]. Two *meso* homodimer equivalents can be found to the four cyclopentanepentols, by removing the suitable apexes, and they are (1 β ,2 β ,3 β ,4 β)-1,2,3,4-tetrahydroxy-cyclobutane (cis,cis,cis-1,2,3,4-tetracarboxycyclobutane) and (1 β ,2 β ,3 α ,4 α)-1,2,3,4-tetrahydroxy-cyclobutane (cis,trans,cis-1,2,3,4-tetracarboxy-cyclobutane). The first cyclobutane derivative has to be considered *meso*, although it gives only one C_2 *symm.* product, by alternative dimerization of its enantiomeric halves. The second equivalent, (1 β ,2 β ,3 α ,4 α)-1,2,3,4-tetrahydroxy-cyclobutane, is *super-symmetric*.

Nine inositols (hexahydroxycyclohexanes) are known [81,82], two are C_2 *symm.*, chiral and enantiomeric, and seven devoid of optical activity. Of all the latter, only cis-inositol and muco-inositol can be treated as homodimeric *meso*. Alternative dimerization of the two enantiomeric halves of cis-inositol gives neo-inositol. The latter is a special molecule: it has a mirror plane of symmetry containing some atoms, hence it is a heterodimer. On the other hand, it has an axis of symmetry, and when rotated by 180° around this axis, the same structural context is found, that is a feature of C_2 *symm.* compounds. Of this reason neo-inositol (Fig. 6) has to be considered *super-symmetric*. When treated as *meso* heterodimer, and its linking support removed, neo-inositol gives another *super-symmetric* combination, i.e. (1 β ,2 β ,3 α ,4 α)-1,2,3,4-tetrahydroxy-cyclobutane. epi-Inositol is a *meso* heterodimer and when its binding support removed, a *meso* homodimer, (1 α ,2 α ,3 α ,4 α)-1,2,3,4-tetrahydroxy-cyclobutane, is obtained. The latter is *meso* since its isomer is a *super-symmetric* derivative, (1 β ,2 β ,3 α ,4 α)-1,2,3,4-tetrahydroxy-cyclobutane. allo-Inositol is a homodimer with a mirror plane of symmetry. When its enantiomeric halves are alternatively dimerized, the symmetry is perturbed and two chiral, C_2 *symm.* enantiomers are produced, D-(+)-chiro- and L-(-)-chiro-inositol. allo Inositol is the only inositol isomer with a typical behavior of *meso* compound giving a triad. mio-Inositol is a *meso* heterodimer leading directly to (1 β ,2 β ,3 α ,4 α)-1,2,3,4-tetrahydroxy-cyclobutane, while the latter is obtained from muco-inositol (*meso*) via scillitol, scillitol being also *super-symmetric*.




Figure6. Illustration of finding of meso homodimer equivalents of meso heterodimers and of behavior of some homodimers.




Figure 7. *meso* Homodimer equivalents of polyols, epoxy compounds and anhydrides.




Figure8.Findingof*meso* homodimerequivalentsofcyclic*meso* heterodimers.

2. (C) *meso* Heterodimers with a Binding Support Devoid of Symmetry (>CRR', etc.). Compounds of different classes – linear (polyols), cyclic, bicyclic, etc., – have been approached (Figs. 7 and 8). Skipped polyols *per se* or intermingled with methyl or dimethyl groups have been synthesized since they are structural motifs of some natural compounds. Erythritol is not only a *meso* homodimer polyol for xylitol and ribitol as *meso* heterodimer polyols but also for 6

and 14 [83], 7 [1], 32 [22] etc. meso-Hexitol is homodimer for 3, 5, 6 [84], 56-59 [21], 27, 28 [1], etc. meso-Octitol is homodimer for 31, 86 [1], 34 [22], 85 and 86 [67], and meso-decitol for 60 [21], 35 and 56 [1], etc.

Epoxy group – in cyclohexene oxide [85], 201, 38 [22], 1, 18, 26 [86], 74 [67], 28a [45], XIII [79], 10, 12 [65], 2a [87] – is equivalent to two hydroxyls in homodimers. In all previous epoxy or anhydride compounds, the number of C atoms is the same in the equivalent *meso* homodimers. The *meso* homodimers of the next *meso* heterodimers would be found by elimination of the suitable apices and linking of the adjacent atoms. For a series of *meso* heterodimers, their *meso* homodimer equivalents are indicated (Fig. 7). Others could be found according to the rules explicitly presented or tacitly included. Dianhydride XIII [79] has **trans,trans,trans**-1,2,3,4-tetracarboxy-cyclobutane as equivalent *meso* homodimer (Fig. 6). 11 [65] and 199 [22] has cis-1,2-dicarboxy-cyclobutane and cis-1,2-dihydroxy-cyclobutane, respectively. Both 13 [50] and 39 [86] have cis-3,6-cyclohexene-1 as *meso* homodimer equivalent, and 46, a similar equivalent.

Di- or tetrasubstituted cyclobutane or disubstituted cyclobutene are the *meso* homodimer for: 5a [88], 15a [85], **meso** diester [1], 47 [22], 5 [59], 38 [1], 61a, 59, 45c [22]. For a substituted ring of eight atoms, 52 [22]; and for bicycle structures 211a and 215e [22], *meso* homodimer is based on a cyclohexane or cyclohexene ring. N Atom of a bicyclic alkaloid 75 [37], has been transformed in two amino groups. *meso* Homodimer equivalents of trans decaline derivatives – 69 and 70 [1], 4 [89], and a highly pyramidalized alkane 8 [90], are based on cyclooctane or cyclooctadiene. *meso* Homodimer equivalents of numerous bicyclic compounds – 3 and 8 [91], 2 and 5 [65], 64 [86], 102 [45], tropine [92] – are based on cis-1,4-disubstituted cyclohexane or cyclohexene (Fig. 8).

Also to this group belongs some compounds with a symmetric binding support, that are characterized by a relatively low structural plasticity, due to a rigid linking between the *meso* homodimeric unit and their binding **unit**. These seeable *meso* heterodimers appear especially when two rings, both containing fewer atoms than six, form bicyclic molecules. Although in relatively small number these molecules justify the existence of a distinct group. The quality of *meso* of the constituents of this group, although seeable, can be proved only by removing their linking support and then acting on their *meso* equivalent homodimer. Such molecules are: XIII [79], 2 [93], 9, 10, 11, 13 [65]; 2a [87], 199, 215d [22] (Fig. 7), 5a [88], 15a [85] (Fig. 8).

CONCLUSIONS

The two enantiomeric halves of *meso* compounds constitute the most important chemical duality. Of this reason these compounds constitute a(n) (unexploited) treasure for chemical duality, and it significantly reduces the huge advance of physics in comparison with chemistry as the choice and the most successful object of philosophy. The relationship between two groups of diastereomeric isomers – *meso* and *C₂symmetrical* – has been studied in detail and it disclosed a chemical paradox among inositols and a new distinct group of chemical combinations, the **super-symmetrical** ones. The chemical paradox of inositols found in this paper, harmoniously completes the definition of *meso* compounds as an ensemble of two enantiomeric halves. For the first time in chemical literature, probably, a systematization of *meso* compounds has been made. **The fundamental characteristic of a *meso* compound consists in its capacity to produce at least one *C₂symmetrical* isomer, by alternative dimerization of its enantiomeric halves.**

References

- [1] Hoffmann RW. *meso* Compounds: stepchildren or favored children of stereoselective synthesis? Angewandte Chemie-International Edition. 2003;42:1096-1109.

- [2] Pfaltz A, Drury III WJ. Design of chiral ligands for asymmetric catalysis: From C₂-symmetric PP- and NN-ligands to sterically and electronically nonsymmetrical PN-ligands. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;101:5723-5726.
- [3] Mallette JR, Casale JF. Rapid determination of the isomeric truxillines in illicit cocaine via capillary gas chromatography/flame ionization detection and their use and implication in the determination of cocaine origin and trafficking routes. *Journal of Chromatography A*. 2014;1364:234-240.
- [4] Iga DPI. meso Chemical symmetry definition by molecular modelling and application to inositols. *Asian Journal of Applied Chemistry Research*. 2023;14:44-62.
- [5] Gerberich HR, Walters WD. The thermal decomposition of cis-1,2-dimethylcyclobutane. *Journal of the American Chemical Society*. 1961; 83:3935-3939.
- [6] Gerberich HR, Walters WD. The thermal decomposition of trans-1,2-dimethylcyclobutane. *Journal of the American Chemical Society*. 1961;83:4884-4888.
- [7] Allinger NL, Tushaus LA. Conformational analysis. XLIII. Stereochemical studies in the cyclobutane ring system. *Journal of Organic Chemistry*. 1965;30:1945-1951.
- [8] Griffin GW, Veber D. cis,cis,cis-1,2,3,4-Tetracarbomethoxycyclobutane; structure of β -heptacyclene. *Journal of the American Chemical Society*. 1960;82:6417.
- [9] Kricheldorf HR, Schwarz G. New polymer syntheses. 17. Cis/trans isomerism of 1,4-cyclohexanedicarboxylic acid in crystalline liquid-crystalline and amorphous polyesters. *Makromolecular Chemistry*. 1987;188:1281-1294.
- [10] Shimada KN, Kawaguchi JM, Dobashi K et al. Efficient synthesis of selenocarbonyl compounds by treating carbonyl compounds with bis(1,5-cyclooctanediyloboryl) selenide. *Bulletin of the Chemical Society of Japan*. 1997;70:197-206.
- [11] Kirchen R, Ranganayakulu K, Rauk A. Dimethylcyclodecyl cations. Evidence for μ -hydride bridging. *Journal of the American Chemical Society*. 1981;103:588-596.
- [12] Fischer E, Hertz J. Reduction of galactaric acid. *Berichte der deutschen chemischen Gesellschaft*. 1892;25:1247-1261.
- [13] Fischer E. Configuration of tartaric acid. *Berichte der deutschen chemischen Gesellschaft*. 1896;29:1377-1383.
- [14] Fischer E, Hirschberger J. On Mannose IV. *Berichte der deutschen chemischen Gesellschaft*. 1889;22:3218-3224.
- [15] Vickery HB. Assignment of DL prefixes to the tartaric acids. *Journal of the Chemical Education*. 1957;34:339-341.
- [16] Dang TP, Kagan HB. The asymmetric synthesis of hydratropic acid and amino-acids by homogeneous catalytic hydrogenation. *Journal of the Chemical Society. Chemical Communications*. 1971;481-481.
- [17] Kagan HB, Dang TP. Asymmetric Catalytic Reduction with Transition Metal Complexes I. A Catalytic System of Rhodium (I) with (-)-23-(9-Isopropylidene-23-dihydroxy-14-bis(diphenylphosphino) butane a New Chiral Diphosphine. *Journal of the American Chemical Society*. 1972;94:6429-6433.
- [18] Ghosh AK, Mathivanan P, Cappiello J. C₂-Symmetric chiral bis(oxazoline)-metal complexes in catalytic asymmetric synthesis. *Tetrahedron Asymmetry*. 1998;9:1-45.
- [19] Shiono Y, Sasaki T, Shibuya F, Yasuda Y, Koseki T, Supratman U. Isolation of a Phomoxanthone A Derivative a New Metabolite of Tetrahydroxanthone from a *Phomopsis* sp. Isolated from the Mangrove *Rhizophora mucronata*. *Natural Products Communications*. 2013;8:1735-1737.
- [20] Ishii H, Sakurada E, (nee Kawanabe), Furukawa T, Koseki C, Ogata K, Koseki N, Ishikawa T, Harayama T. Photochemistry of 2-(3,4,5-trimethoxyphenyl)-4-(3,4-methylenedioxyphenyl)-4-oxo-2-butenonitrile (β -cyanochalcone): anomalous dimerization through isomerization in the solid state. *Chemical and Pharmaceutical Bulletin*. 1991; 39: 2173-2175.
- [21] Magnuson SR. Two-Directional synthesis and its use in natural product synthesis. *Tetrahedron*. 1995;51:2167-2213.
- [22] Garcia-Urdiales E, Alfonso I, Gotor V. Enantioselective Enzymatic Desymmetrizations in Organic Synthesis. *Chemical Reviews*. 2005;105:313-354.

- [23] Zhang L, Guo Z, Chen J, Xu Q, Lin H, Hu K, Guan X, Shen Y. Mechanism of 2,3-butanediol stereoisomers formation in a newly isolated *Serratia* sp T241. *Scientific Reports*. 2016;6:19257:1-12.
- [24] Rubin LJ, Lardy HA, Fischer HOL. Synthesis of the Optically Active Enantiomorphous 2,3-Butanediols. *Journal of the American Chemical Society*. 1952;74:425-428.
- [25] Huang R, Zhou X, Xuc T, Yang X, Liu Y. Diketopiperazines from Marine Organisms. *Chemical Biodiversity*. 2010;7:2809-2829.
- [26] Borthwick AD. 2,5-Diketopiperazines: Synthesis Reactions Medicinal Chemistry and Bioactive Natural Products. *Chemical Reviews*. 2012;112:3641-3716.
- [27] Yang S-W, Chan T-M, Terracciano J et al. Structure Elucidation of a New Diketopiperazine Sch 725418 from *Micromonospora* sp. *Journal of Antibiotics*. 2004;57:345-347.
- [28] Kozlovsky AG, Vinokurova NG, Adanin VM, Burkhardt G, Dahse H-M, Grafe U. New diketopiperazine alkaloids from *Penicillium fellutanum*. *Journal of Natural Products*. 2000;63:698-700.
- [29] Li S-M. Prenylated indole derivatives from fungi: structure diversity biological activities biosynthesis and chemoenzymatic synthesis. *Natural Product Reports*. 2010;27:57-78.
- [30] Garg NK, Stoltz BM. The formal total synthesis of dragmacidin B trans-dragmacidin C and cis- and trans-dihydrohamacanthins A. *Tetrahedron Letters*. 2005;46:2423-2426.
- [31] Maskey RP, Kock I, Helmke E, Laatsch H. Isolation and Structure Determination of Phenazostatin D a New Phenazine from a Marine Actinomycete Isolate *Pseudonocardia* sp B6273. *Zeitschrift fuer Naturforschung*. 2003;58b:692-694.
- [32] Yamamura S, Niwa M, Yukimasa T, Nonoyama M. The isolation and structures of novel neolignans and neosesquiterpenes from *Heterotropa takaoi* M. *Bulletin of the Chemical Society of Japan*. 1982;55:3573-3579.
- [33] Davis RA, Carroll AR, Duffy S, Avery VM, Guymier GP, Forster PI, Quinn RJ. Endiandrin A a potent glucocorticoid receptor binder isolated from the Australian plant *Endiandra anthropophagorum*. *Journal of Natural Products*. 2007;70:1118-1121.
- [34] Davis RA, Barnes EC, Longden J, Avery VM, Healy PC. Isolation structure elucidation and cytotoxic evaluation of endiandrin B from the Australian rainforest plant *Endiandra anthropophagorum*. *Bioorganic Medical Chemistry*. 2009;17:1387-1392.
- [35] Green BS, Rejtö M. μ -Truxinic acid. *Journal of Organic Chemistry*. 1974;39:3284-3285.
- [36] Isaza JH, Ito H, Yoshida T. Monochaetin a Di-hyperin Ester of Tetrahydroxy- μ -truxinic Acid from *Monochaetum multiflorum*. *Chemical and Pharmaceutical Bulletin*. 1999;47:1510-1511.
- [37] Dembitsky VM. Bioactive cyclobutane-containing alkaloids. *Journal of Natural Medicine*. 2008;62:1-33.
- [38] Fabricant DS, Nikolic D, Lankin DC et al. Cimipronidine a Cyclic Guanidine Alkaloid from *Cimicifuga racemosa*. *Journal of Natural Products*. 2005;68:1266-1270.
- [39] Smith MB, March J. *March's Advanced Organic Chemistry: Reactions Mechanisms and Structure* Sixth Edition Wiley: New York; 2007.
- [40] Li W D Z, Ma B C. A Simple Biomimetic Synthesis of dl-Chamaejasmine a Unique 3,3'-Biflavanone. *Organic Letters*. 2005;7:271-274.
- [41] Lautens M, Ma S, Belter RK, Chiu P, Leschzinner A. An Efficient Synthesis of 13-Cyclohexadienes from Oxabicyclo[2.2.1]heptenes via Tandem Ring Opening-Peterson Elimination Reactions. *Journal of Organic Chemistry*. 1992;57:4065-4066.
- [42] Eliel EL, Wilen SH. *Stereochemistry of Organic Compounds*: Wiley Interscience Chichester; 1994.
- [43] Sanchez-Sordo I, Chaves-Pouso A, Mateos-Gil J, Rivera-Chao E, Fananas-Mastral M. Desymmetrization of meso-dibromocycloalkenes by copper-catalyzed asymmetric boronate coupling with alkynes. *Chemical Catalysis*. 2023;3:100730.
- [44] Inoue M, Lee N, Kasuya S, Sato T, Hirama M, Moriyama M, Fukuyama Y. Total Synthesis and Bioactivity of an Unnatural Enantiomer of Merrilactone A: Development of an Enantioselective Desymmetrization Strategy. *Journal of Organic Chemistry*. 2007;72:3065-3075.

- [45] Wang Z, Chen Z, Sun J. Catalytic Asymmetric Nucleophilic Openings of 3-Substituted Oxetanes. *Organic and Biomolecular Chemistry*. 2014;12:6028-6032.
- [46] Poplata S, Tröster A, Zou Y-Q, Bach T. Recent Advances in the Synthesis of Cyclobutanes by Olefin [2+2] Photocycloaddition Reactions. *Chemical Reviews*. 2016;116:9748–9815.
- [47] Nonappa K, Ahonen, Lahtinen M, Kolehmainen E. Cyclic dpeptides: catalyst/promoter-free rapid and environmentally benign cyclization of free amino acids. *Green Chemistry*. 2011; Issue 5.
- [48] Ruolin Y, Zhonghong Y, Congying C, Jianhua L. Constituents and Activities of *Acorus tatarinowii*. *Medical Research Archives*. 2017;5:1-14.
- [49] Gal AEP, Pentchev G, Massey JM, Brady RO. L-Glucosylceramide: synthesis, properties and resistance to catabolism by glucocerebrosidase in vitro. *Proceedings of the National Academy of Sciences of the United States of America*. 1979;76:3083-3086.
- [50] Johnson CR, Golebiowski A, Steensma DH. Enzymatic Asymmetrization in Organic Media: Synthesis of Unnatural Glucose from Cycloheptatriene. *Journal of the American Chemical Society*. 1992;114:9414-9418.
- [51] Anderson PJ. Oxidation of 3-deoxyxylitol by L-iditol dehydrogenase. *Biochimica Biophysica Acta*. 1965;110:627-629.
- [52] Oka J, Ueda K, Hayaishi O, Komura H, Nakanishi K. ADP-Ribosyl protein lyase Purification properties and identification of the product. *Journal of Biological Chemistry*. 1984;259:986-995.
- [53] Boydell AJM, Jeffery J, Bürkstümmer E, Linclau B. Short Synthesis of Enantiopure C₂-Symmetric 12:45-Diepoxy pentane and “Pseudo”-C₂-Symmetric 3-Azido-12:45-diepoxy pentane from Arabitol. *Journal of Organic Chemistry*. 2003;68:8252–8255.
- [54] Setser DW, Rabinovitch BS. Singlet Methylene From Thermal Decomposition Of Diazomethane Unimolecular Reactions Of Chemically Activated Cyclopropane And Dimethylcyclopropane Molecules. *Canadian Journal of Chemistry*. 1962;40:1425-1451.
- [55] Wang Y-F, Chen C-S, Girdaukas G, Sih CJ. Bifunctional Chiral Synthons via Biochemical Methods 3 Optical Purity Enhancement in Enzymic Asymmetric Catalysis. *Journal of the American Chemical Society*. 1984;106:3695-3696.
- [56] Metzler DE, Metzler CM. *Biochemistry: the chemical reactions of living cells*: Elsevier, Amsterdam; 2003.
- [57] Gao X, Han H, Krische MJ. Direct Generation of Acyclic Polypropionate Stereopolyads via Double Diastereo- and Enantioselective Iridium-Catalyzed Crotylation of 1,3-Diols: Beyond Stepwise Carbonyl Addition in Polyketide Construction. *Journal of the American Chemical Society*. 2011;133:12795-12800.
- [58] Manna MS, Mukherjee, S. Org. Catalytic asymmetric desymmetrization approaches to enantioenriched cyclopentanes. *Biomolecular Chemistry*. 2015;13:18-24.
- [59] Annunziato G, Pieroni M, Benoni R, Campanini B, Pertinhez TA, Pecchini C. Cyclopropane-12-dicarboxylic acids as new tools for the biophysical investigation of *O*-acetylserine sulfhydrylases by fluorimetric methods and saturation transfer difference (STD) NMR. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2016;31:sup4:78-87.
- [60] Choudhary AL, Naughton M, Montánchez I, Dobson ADW, Rai DK. Current Status and Future Prospects of Marine Natural Products (MNPs) as Antimicrobials. *Marine Drugs*. 2017;15:1-42.
- [61] Wang YY. Chemical constituents and anti-inflammatory activity from the stem of *Neolitsea daibuensis*. [MSD thesis]: Kaohsiung Medical University, Kaohsiung Taiwan; 2011.
- [62] Wang Y-F, Sih CJ. Bifunctional chiral synthons via biochemical methods 4 Chiral precursors to (+)-biotin and (–)-a-factor. *Tetrahedron Letters*. 1984;25:4999-5002.
- [63] Novak M, Salemink CA, Khan I. Biological activity of the alkaloids of *Erythroxylum coca* and *Erythroxylum novogranatense*. *Journal of Ethnopharmacology*. 1984;10:261–274.
- [64] Naemura K, Takahashi N, Chikamatsu H. Enzyme-catalyzed asymmetric hydrolysis of meso-substrate. The facile synthesis of both enantiomers of cis-2,5-disubstituted tetrahydrofuran derivatives. *Chemical Letters*. 1988;17:1717–1720.

- [65] Jaeschke G, Seebach D. Highly enantioselective ring opening of cyclic *meso*-anhydrides to isopropyl hemiesters with Ti-TADDOLates: an alternative to hydrolytic enzymes? *Journal of Organic Chemistry*. 1998;63:1190-1197.
- [66] Mori Y, Hayashi H. Practical synthesis of 1,3-*o*-di-*tert*-butylsilylene-protected d- and l-erythritols as a four-carbon chiral building block. *Journal of Organic Chemistry*. 2001;66:8666-8668.
- [67] Newhouse T, Baran PS, Hoffmann RW. The economies of synthesis. *Chemical Society Reviews*. 2009;38:3010-3021.
- [68] Morimoto Y, Iwai T, Kinoshita T. Effective combination of two-directional synthesis and rhenium(VII) chemistry: total synthesis of *meso* polyether teurilene. *Journal of the American Chemical Society*. 1999;121:6792-6797.
- [69] Hamada J-I, Tsunashima S, Sato S. The reactions of the singlet NH radicals with C₄olefins in the liquid phase. *Bulletin of the Chemical Society of Japan*. 1983;56:662-666.
- [70] Levy DE, Fügedi P. (eds) *The Organic Chemistry of Sugars*: Taylor and Francis, London; 2006.
- [71] Fletcher Jr HG, Hudson CS. 1,5-Anhydro-xylitol. *Journal of the American Chemical Society*. 1947;69:921-924.
- [72] Hann RM, Ness AT, Hudson CS. 2,4:3,5-Dimethylene-DL-xylitol and 2,4-methylene-xylitol. *Journal of the American Chemical Society*. 1944;66:670-673.
- [73] Schmidt RR, Lieberknecht A. Functional D- and L-ribose derivative by retro racemic splitting. *Angewandte Chemie*. 1978;90:821-822.
- [74] Durig JR, Nease AB, Rizzolo JJ. Vibrational spectra and torsional analyses of *cis*-2,3-dimethyloxirane and *trans*-2,3-dimethyloxirane. *Journal of Molecular Structure*. 1983;95:59-84.
- [75] DePuy CH, Fuenfschilling PC, Andrist AH, Olson JM. Stereochemistry of the electrophilic ring opening of cyclopropanes 2 Reaction of *cis*- and *trans*-1,2,3-trimethylcyclopropane with deuterium(1+) ion. *Journal of the American Chemical Society*. 1977;99:6297-6303.
- [76] Kozhushkov SI, Leonov A, de Meijere A. Simple Large-Scale Preparation of 1,2,3-Tris-Acceptor Substituted Cyclopropanes. *Synthesis*. 2003;(6):0956-0958.
- [77] Papa AJ. Reaction of triethyl *trans*-1,2,3-tricyanocyclopropane-1,2,3-tricarboxylate with base Formation of a substituted cyclopropane anion radical. *Journal of Organic Chemistry*. 1968;33:2532-2534.
- [78] Griffin GW, Velluro AF, Furukawa K. The Chemistry of Photodimers of Maleic and Fumaric Acid Derivatives I Dimethyl Fumarate Dimer. *Journal of the American Chemical Society*. 1961;83:2725-2728.
- [79] Velluro AF, Griffin GW. Electrolytic Oxidation of Cyclobutane-1,3-dicarboxylic Acids An Electrochemical Synthesis of 2,4-Dicarbomethoxybicyclobutane. *Journal of Organic Chemistry*. 1966;31:2241-2244.
- [80] Collins PM. *Dictionary of Carbohydrates*: Chapman & Hall/CRC Taylor & Francis Group Boca Raton London New York; 2006.
- [81] Pigman W, ed. *The Carbohydrates: Chemistry, Biochemistry, Physiology* Academic Press New York; 1957.
- [82] Finar IL. *Organic Chemistry Vol 2*: Longmans Green and Co Ltd London; 1964.
- [83] Tullis J, Vares L, Kann N, Norrby P-O, Rein T. Reagent Control of Geometric Selectivity and Enantiotopic Group Preference in Asymmetric Horner-Wadsworth-Emmons Reactions with *meso*-Dialdehydes. *Journal of Organic Chemistry*. 1998;63:8284-8294.
- [84] Wang Z, Deschenes D. Enantiospecific Synthesis via Sequential Diastereofacial and Diastereotopic Group Selective Reactions: Enantiodivergent Synthesis of syn-1,3-Polyols. *Journal of the American Chemical Society*. 1992;114:1090-1091.
- [85] Bhuniya D, Datta Gupta A, Singh VK. Design Synthesis and Application of Chiral Nonracemic Lithium Amide Bases in Enantioselective Deprotonation of Epoxides. *Journal of Organic Chemistry*. 1996;61:6108-6113.
- [86] Woo S, Keay BA. "SN2' and "SN2' Like" Ring Openings of Oxa-n-Cyclo Systems". *Synthesis*. 1996;7:669-686.

- [87] Konowalchuk DJ, Hall DG. Divergent Synthesis of 1234-Tetrasubstituted Cyclobutenes from a Common Scaffold: Enantioselective Desymmetrization by Dual-Catalyzed Photoredox Cross-Coupling. *Angewandte Chemie-International Edition*. 2023;e202313503:1-6.
- [88] Asami M. A Novel Asymmetric Synthesis of Chiral Cyclopentanoid Building Blocks by the Use of Chiral Lithium Amide. *Bulletin of the Chemical Society of Japan*. 1990;63:1402-1408.
- [89] Hiroya K, Ogasawara K. A concise enantio- and diastereo-controlled synthesis of (–)-quinic acid and (–)-shikimic acid. *Chemical Communications*. 1998;2033-2034.
- [90] Camps P, Pujol X, Vazquez S, Pericas MA, Puigjaner C, Sola L. Cross-coupling of a functionalized highly pyramidalized alkene: DSC and NMR study of the [2+2] retrocycloaddition of cyclobutane cross products hyperstability and pyramidalization of the formed dienes. *Tetrahedron*. 2001;57:8511-8520.
- [91] Lautens M, Abd-El-Aziz AS, Lough A. Opening of Oxabicyclo[3.2.1]octenes with Organolithium Reagents A Route to Cyclic and Acyclic Compounds with High Stereocontrol. *Journal of Organic Chemistry*. 1990;55:5305-5306.
- [92] Al-Badr AA, Muhtadi FJ. Atropine. *Analytical Profiles of Drug Substances* 1985;14:325-389.
- [93] Reusch W, Priddy DB. vic-Cyclopropanediols from Lithium in Ammonia Reductions of Cyclic β -Diketones. *Journal of the American Chemical Society*. 1969;91:3677-3678.