

Response of onion (*Allium Cepa* L.) to nitrogen fertilizer rates and intra-row spacing in Southern Zone of Tigray, Ethiopia

ABSTRACT

Onion (*Allium cepa* L.) is a significant crop in terms of its economic value. However, its productivity is low due to different factors. Farmers apply excessive nitrogen fertilizers to increase yields, while others apply insufficient amounts, leading to reduced onion yields. Additionally, most farmers use different spacing, highlighting the need to determine the appropriate spacing in this area. Therefore, this study was carried out with the aim of evaluating the combined effect of nitrogen fertilizer rates and intra-row spacing on the growth, yield, and yield components of onion. A field experiment was conducted in the experimental field of Alamata Agricultural Research Center kara Adishabo in 2020 with experimental treatments (0, 23, 46, 69, and 92 kg N/ha) and intra-row spacing (5, 10, and 15 cm) using a factorial arrangement in a randomized complete block design with three replications. Results indicated that the interaction effect of nitrogen and intra-row spacing significantly swayed most of the yield and yield components of the onion where The highest maturity date was recorded from the 92 Kg N/ha and 15 cm (140 days) and 69 Kg N/ha and 10 cm (127 days) respectively. The highest (40.01t/ha) and lowest (15.03 t/ha) marketable yields were recorded from the treatment combinations of 69 kg N/ha and 5 cm and 0 kg N/ha and 15 cm respectively. Application of 69 Kg N/ha and 5 cm of intra-row spacing in the study area can lead to maximum yield.

Keywords: Onion, intra-row spacing, nitrogen, marketable bulb yield

INTRODUCTION

Onion (*Allium cepa* L.) is a significant crop in terms of its economic value. It belongs to the *Allium* genus of the Alliaceae family and is believed to have originated in Central Asia, specifically between Turkmenistan and Afghanistan. Some of its related species can still be found growing in the wild in this region (Bagali et al., 2012). Onions are known for being one of the highest dietary sources of flavonoids, which have been linked to a decreased risk of cancer, heart disease, and diabetes (Yemane et al., 2013)..

Multiple factors contribute to the low productivity of onion in Ethiopia. The inadequate onion yield in the country is attributed to incorrect fertilizer application, substandard agronomic practices, disease and pest infestations, weed growth, insufficient high-quality seed supply, limited pesticide availability, and inadequate irrigation water (Etana, 2019; Lemma and Shimelis, 2003). Nitrogen fertilization and plant spacing are among the most important factors that determine onion growth and yield. Nitrogen is crucial for achieving optimal onion yield as it is necessary for enhancing bulb size and overall crop productivity (Nasreen et al., 2007).

Regulating plant spacing is a cultural practice that influences bulb size, shape, and yield (Awat et al., 2010). A higher yield and better control of over or under-bulb size could be obtained if plants are grown at optimum density. As population density increased, bulb neck diameters, mean bulb weight, and plant height all decreased. When plants are grown too closely together, they compete for sunlight, essential nutrients, water, and air, resulting in the production of small, low-quality bulbs. Conversely, plants that are grown too far apart may produce vigorous individual plants but yield less per given area (Kihara et al., 2015; Winch, 2006).

In Raya Valley various types of vegetables had been introduced to the farmers with plots in the irrigation sites. Traditionally, farmers in this area have been cultivating onions (Berhane et al., 2010). However, onion yields have been low due to suboptimal fertilizer application and poor agronomic practices, particularly in plant spacing. Some farmers apply excessive nitrogen fertilizers to increase yields, while others apply insufficient amounts, leading to reduced onion yields. Additionally, most farmers use narrower intra-row spacing and the rest use wider spacing, highlighting the need to determine the appropriate intra-row spacing in this area. Lemma and Shimeles (2003) proposed that to maximize onion productivity, a comprehensive set of guidelines tailored to each specific growing region is necessary. Therefore, addressing this issue can be achieved by implementing optimal nitrogen fertilizer rates and suitable plant spacing in the study area. The overall aim of this study was to evaluate the combined effect of nitrogen fertilizer rates and intra-row spacing on the growth, yield, and yield components of onion.

MATERIALS AND METHODS

Description of the study area:

The research was conducted at the Alamata Agricultural Research Center experimental site, situated in the Southern Zone of Tigray, Raya Azobo Woreda, Ethiopia, specifically in Kara Adishabo. The area is characterized by a dry semi-arid climate (Araya et al., 2010). Geographically, it is positioned between 12°38'50" N and 12°44'36" N latitude and 39°35'10" E and 39°45'10" E longitude. The annual mean minimum, maximum, and average precipitation in the study area are 385.7 mm, 681.1 mm, and 543.6 mm, respectively, with a mean monthly

maximum temperature of 29.9 °C and minimum temperatures of 15.9 °C. The Woreda's altitude ranges from 1646 to 1670 m.a.s.l. It encompasses various soil types, including Vertisoils, Nitsoil, Combisols, and Luvisols. Vertisol (black soil with swelling characteristics) is the predominant soil type, covering over 70% of the study area (National Meteorology Agency, 2018 and Raya Azebo Woreda ARD Office, 2019).




Figure 1: Map of the study area

Experimental Treatments, Designs and procedures:

The experimental treatments included five nitrogen levels (0, 23, 46, 69, and 92 Kg/ha of N) and three intra-row spacing options (5, 10, and 15 cm). These factors were combined in a factorial design and arranged in a Randomized Complete Block Design (RCBD) with three replications. Spacing between blocks and plots was 1.5 m and 1 m, respectively with a 2 m x 3 m plot area. The spacing between furrows was kept at 40 cm and the spacing between the double rows in a furrow was 20 cm. The seedlings of the Bombay Red onion variety were grown on well-prepared nursery beds following the recommended management practices. This variety is well-suited and extensively grown in the study area. Prior to seedling transplanting, the experimental field was ploughed, pulverized, and leveled. Seedlings were transplanted to the

experimental plots when they reached a height of 12-15 cm or at the 3-4 true leaf stage, carefully uprooted from the nursery bed. Only healthy, vigorous, and uniform seedlings were transplanted, and any gaps were filled within a week after transplanting. All plots received a uniform application of Triple superphosphate (46% P_2O_5) at a rate of 92 kg P_2O_5 as a source of phosphorus. Additionally, half rates of urea (46% N) were applied during transplanting, while the remaining half rates of urea (46% N) were applied after 30 days of transplanting. Soil moisture was maintained at field capacity during planting, and furrow irrigation was used for water application in each plot. A four-day irrigation interval was upheld for the first four weeks, after which it was extended to five to seven-day intervals until 15 days before harvest, at which point irrigation was completely stopped (EARO, 2004).



Figure 2 : Field performance of onion

Soil Sampling:

Prior to planting, soil samples were collected from 10 random spots diagonally at six-meter intervals using an augur at a depth of 0-30 cm from the experimental site. The collected soil samples were combined into one composite sample. These composite soil samples were air-dried, thoroughly mixed, and ground to pass a 2 mm sieve size before laboratory analysis.

Data Collection

Soil data:

Prior to planting, the soil's physical and chemical properties, including soil texture, pH, electrical conductivity (EC), cation exchange capacity (CEC), exchangeable potassium and sodium, total nitrogen, total organic carbon, and total available phosphorus, were assessed and recorded.

Crop data:

Growth parameters (Number of leaves per plant, days to maturity and fresh biomass yield (g/plant), yield and yield components (Bulb diameter (cm), Dry biomass yield (g/plant), marketable bulb yield (t/ha) and unmarketable bulb yield (t/ha)) (See fig 2) of onions was collected from the eight central double rows (out of ten rows). Ten pre-tagged plants were randomly selected in each plot.

Data Analysis

Soil data analysis:

The collected soil samples were composite to one sample. The bulked soil samples were air-dried, thoroughly mixed, and ground to pass 2 mm sieve size before laboratory analysis. Then the samples were properly labeled, packed, and transported to Mekelle Soil Laboratory for the analysis of major physical and chemical properties.

Crop data analysis:

All data were presented as means and statistically analyzed using R software version 3.5.3. All the means data recorded of treatments were compared to each other for any significant differences using the least significant difference (LSD) method. The LSD (Least Significant Difference) was calculated based on the data, allowing for testing the differences among means. Separate analyses were performed for each data at the probability level for determination of significance ($P < 0.05$), where the analysis of variance indicated the presence of significant treatment differences.

RESULT AND DISCUSSION

Soil Physico-Chemical Properties of the Experiment

The results of the soil analysis (Table 1) before planting indicated that the soil pH value of the experimental site was 7.87 which were moderately alkaline (Murphy, 1968 and Tekalign, 1991).

Table 1 : Physico-chemical properties of the experimental soil

Soil properties	Unit	Experimental field value	Ratings	Sources
Particle size distribution	size %			
Sand	%	45		
Silt	%	35		
Clay	%	20		
Textural class			Loamy soil	Hazelton and Murphy, 2007
Soil Ph		7.87	Moderately alkaline	Tekalign,1991
Organic carbon	%	1.39	Low	Tekalign,1991
Organic matter	%	1.79	Low	Berhanu, 1980
Total Nitrogen	%	0.114	Low	Tekalign,1991
Total available phosphorus	mg /kg	14.89	Medium	Cottenie, 1980
Cation exchange capacity	Cmol(+)/kg	35.2	High	Hazelton and Murphy, 2007
Exchangeable Sodium	Cmol(+)/kg	1.83	High	FAO, 2006
Exchangeable potassium	Cmol(+)/kg	0.71	Medium	FAO, 2006
Electrical	dS/m	0.29	Medium	Hazelton and Murphy, 2007

Growth and phenology parameters

Leaf number per plant:

The highest leaf number (18.17) was obtained from the combination of 92 Kg N/ha and 15 cm while the lowest mean number of leaves per plant (4.33 and 6.83, respectively) was recorded at nil Kg N/ha and 23 Kg N/ha with an intra-row spacing of 5 cm for each (Table 2). The study's findings suggested that nitrogen significantly influenced the production of new shoots, leading to enhanced vegetative growth in onion plants and contributing to the increase in leaf number (Alebachew et al., 2019). The increase in leaf numbers due to higher nitrogen rates and wider intra-row spacing in this study is linked to reduced competition among plants for nutrients, moisture, and light. This is attributed to the greater production of auxiliary branches at wider intra-row spacing compared to closely spaced plants. These findings align with those of Tekle (2015).

Days to maturity:

Onion plants that were grown at a combination of 92 Kg N/ha and 15 cm, 92 Kg N/ha, and 10 cm, and those grown at 69 Kg N/ha and 15 cm and 69 Kg N/ha and 10 cm need the highest number of days to reach maturity stage. However, onion plants that were grown at a combination of 0 Kg N/ha and 5 cm and 23 Kg N/ha and 5 cm require fewer days to mature. The days to maturity of onion plants increased by 42% when grown with a nitrogen rate of 92 Kg N/ha and 15 cm, compared to plants grown without nitrogen (nil Kg N/ha) and 5 cm (Table 2). The duration to reach maturity was prolonged with higher nitrogen rates and wider intra-row spacing. Hence, plants treated with a higher rate of nitrogen and grown at wider intra-row spacing need more days to mature than those supplied with fewer rates of nitrogen and narrower intra-row spacing. The postponement of maturity resulting from higher nitrogen rates and wider intra-row spacing may be attributed to nitrogen excess, which delays onion maturity by prolonging the vegetative growth period through enhancing the biochemical processes of plants, rather than accelerating the maturity time (Alebachew et al., 2019).

Fresh Total Biomass Yield:

The highest fresh total biomass yield (116.93 g/plant) was obtained from the combined effect of 92 kg N/ha and wider intra-row spacing of 15 cm while the lowest was recorded from nil N fertilizer application combined with 5 cm intra-row spacing (32.33 g/plant) (Table 2). The observed rise in fresh total biomass yield per plant, resulting from higher nitrogen rates and wider intra-row spacing, can be attributed to enhanced leaf growth. This facilitated the accumulation of more assimilates in the bulbs, leading to increased bulb length, bulb diameter, mean bulb weight, root growth, and overall vegetative growth. These factors collectively contributed to the increased fresh biomass yield of onion plants under higher nitrogen rates and wider intra-row spacing, as they provided ample growth factors that allowed for greater assimilate accumulation (Yadav, 2010).

Dry Total Biomass Yield:

The highest dry total biomass yield (26.00 g/plant) was obtained from the combined effect of 92 Kg N/ha and 15 cm of intra-row spacing while the lowest dry total biomass yield (2.90 g/plant) was obtained from nil Kg N/ha and 5 cm of intra-row spacing (Table 2). The substantial increase in total dry biomass yield of onion plants with higher nitrogen fertilizer rates and wider intra-row spacing can be attributed to enhanced leaf growth, promoting the accumulation of more assimilates in the bulbs. This, in turn, led to increased bulb length, bulb diameter, mean bulb weight, root growth, and overall vegetative growth. These factors collectively contributed to the increased fresh biomass yield of onion plants under higher nitrogen rates and wider intra-row spacing, as they provided ample growth factors that allowed for greater assimilate production in the vegetative part and partitioning to the bulbs (Kamble and Kathmale, 2015). Furthermore, Plants grown at the widest spacing exhibited the highest dry total biomass, likely due to reduced competition for growth factors among them. This finding aligns with the results reported by Gessesew et al. (2015).

Table 2: Interaction effect of intra-row spacing and nitrogen fertilizer rates on leaf number per plant, days to maturity, fresh total biomass yield and dry total biomass

N rates (kg/ha)	Intra Row Spacing (cm)	Leaf Number Per Plant	Days To Maturity	Fresh Biomass Yield(g/plant)	Dry Biomass Yield(g/plant)	Total
92	15	18.17a	140.67a	116.93a	26.00a	
	10	16.33b	133.00b	104.30b	18.80b	
	5	14.67c	120.00de	97.77c	17.20bc	
69	15	14.00c	127.67c	91.40d	17.10bc	
	10	12.17d	121.67d	86.90e	13.90de	
	5	9.833e	116.67e	84.00e	12.74def	
46	15	9.17ef	120.33de	76.40f	14.80cd	
	10	8.83ef	111.67f	71.30g	11.90ef	
23	5	8.17fg	109.33fg	66.70h	10.90f	
	15	7.33gh	110.00fg	61.40i	10.30fg	
	10	6.93ghi	105.33ghi	50.80j	7.94gh	
	5	6.83hi	100.67ij	48.00j	7.50hi	
0	15	6.47hi	106.67gh	40.10k	5.00ij	
	10	6.00i	102.00hij	35.50l	3.80j	
	5	4.33j	99.33j	32.33l	2.90j	

LSD (5%)	1.26	4.81	4.38	2.69
CV (%)	7.6	2.5	3.7	13.3

Yield and yield components

Bulb diameter:

The widest bulb diameter was recorded from the combination of 92 Kg N/ha and an intra-row spacing of 15 cm. The narrowest bulb diameter was obtained from the combination of nil Kg N/ha and intra-row spacing of 5 cm (Table 3). The observed increase in bulb diameter with wider intra-row spacing and higher nitrogen fertilizer rates can be attributed to the positive impact of applied nitrogen on vegetative growth and enhanced photosynthesis in plants. This, in turn, promotes the translocation of photosynthates to the storage organ or bulb, resulting in an overall increase in bulb diameter (Thirupathi et al., 2014). The current finding is consistent with the results reported by Singh et al. (2018), who observed that the bulb diameter of onion increases with higher nitrogen fertilizer rates and wider intra-row spacing. The observed increase in bulb diameter at wider spacing can be attributed to the presence of ample nutrients and moisture, resulting from reduced competition effects. This favorable condition facilitates various physiological and metabolic processes, ultimately leading to an increase in dry matter production (Tegen et al., 2016).

Marketable Bulb Yield:

Increasing the rate of nitrogen application significantly increased the production of marketable bulb yield across the decreasing rate of the intra-row spacing. The combined application of 69 Kg N/ha with 5 cm of intra-row spacing produced 40.01 t/ha of marketable bulb yield and the lowest marketable bulb yield (15.03 t/ha) was obtained from treatments treated with nil Kg N/ha and wider intra-row spacing of 15 cm (Table 3). The observed increase in marketable bulb yield with closer spacing and higher nitrogen rates can be attributed to the greater number of bulbs with marketable size per unit area. Additionally, the higher nitrogen rates facilitated the uptake of readily available nutrients, enhancing growth and ultimately leading to improved assimilate partitioning to the storage organ, the bulb (Alebachew et al., 2019). The present study aligns with the results reported by Masrie (2019), who observed that the combination of higher nitrogen rates and narrower intra-row spacing (123 Kg N/ha and 6 cm) resulted in a higher marketable bulb yield.



Figure 3: differentiating marketable and unmarketable bulb yield of onion

Unmarketable Bulb Yield:

The highest unmarketable bulb yield of onion was recorded in nil nitrogen fertilizer application at the intra-row spacing of 5 cm followed by 23 Kg N/ha and 5 cm of intra-row spacing. But, the lowest unmarketable bulb yield of onion was attained in the combination of 92 kg N/ha with an intra-row spacing of 5 cm, 10 cm, and 15 cm, 69 Kg N/ha with an intra-row spacing of 5 cm, 10 cm, and 15 cm and 46 Kg N/ha with an intra-row spacing of 10 cm and 15 cm (Table 3). The observed high unmarketable bulb yield of onion in the treatment combination of no nitrogen and 5 cm intra-row spacing can be attributed to increased plant competition for nitrogen; those competitions reduces vegetative growth and assimilate production, resulting in the production of very small-sized bulbs that are not preferred by consumers (Yemane et al., 2013). The current study aligns with the results reported by Masrie (2019), which indicated that the combination of no nitrogen and 6 cm intra-row spacing resulted in the highest unmarketable yield of 1.05 t/ha.

Table 3: Interaction effect of intra-row spacing and nitrogen fertilizer rates on Intra row spacing, bulb diameter, neck thickness diameter, marketable bulb yield, and unmarketable bulb yield of onion

N rates (kg/ha)	Intra Row Spacing (cm)	Bulb diameter (cm)	Marketable bulb yield (t/ha)	Unmarketable bulb yield (t/ha)
92	15	8.83a	24.10cde	0.03f
	10	7.27b	29.63bc	0.04f
	5	5.80cde	38.30a	0.48ef
69	15	6.60bc	24.10cde	0.09f
	10	6.10cd	32.00b	0.14f
	5	5.63de	40.01a	0.2f
46	15	5.40def	22.20defg	0.16f
	10	5.00efg	23.30def	0.21f
	5	4.70fg	27.00bcd	1.53c
23	15	4.27gh	18.20fgh	0.88de
	10	4.23gh	18.75efgh	0.97d
	5	3.63h	19.42efgh	2.96b
0	15	2.50i	15.03h	0.76de
	10	1.39j	16.75gh	1.1cd
	5	1.03j	17.93fgh	4.96a
LSD (5%)		0.84	5.85	0.48
CV (%)		10.4	14.3	10.1

CONCLUSION

Onion (*Allium cepa* L.) is a commercially important bulb crop grown worldwide. Enhancing onion productivity and production can be achieved through various growth factors. In the study area, improper agronomic practices are a major constraint leading to low onion production and productivity. Among these practices, nitrogen fertilizer rates and intra-row spacing significantly impact onion bulb yield, leaf number per plant, fresh total biomass yield, dry total biomass yield,

days to maturity, bulb diameter, marketable bulb yield, and unmarketable bulb yield. The treatment combination of 69 Kg N/ha and 5 cm of intra-row spacing resulted in the highest marketable yield. Therefore, it can be concluded that planting Onion Bombay Red cultivar under irrigation with 5 cm of intra-row spacing and the application of 69 Kg N/ha in the Kara Adishabo area can lead to maximum yield

REFERENCES

- Berhane, G., Gebrehiwot, A., Berhe, K., & Hoekstra, D. (2010). Commercialization of vegetable production in Alamata Woreda, Northern Ethiopia: processes and impact. IPMS Case Study.
- Debele, B. (1982). The physical criteria and their rating proposed for land evaluation in the highland regions of Ethiopia [soil potentiality]. World Soil Resources Reports (FAO), (54).
- CIMMYT Economics Program. (1988). From agronomic data to farmer recommendations: an economics training manual (No. 27). CIMMYT.
- Cottenie, A. (1981). Soil and plant testing as a basis of fertilizer recommendations.
- Desalegne, L., & Aklilu, S. (2003). Research experiences in onion production.
- Etana, M.B., Mohammed, A. and Nebiyu, A., 2019. Effects of different level of nitrogen fertilizer application on growth, yield, quality and storage life of onion (*Allium cepa* L.) at Jimma, South Western Ethiopia. *Journal of Natural Sciences Research*, 9(10), pp.32-40.
- Etana, M.B., Mohammed, A., & Nebiyu, A. (2019). Effects of different levels of nitrogen fertilizer application on growth, yield, quality, and storage life of onion (*Allium cepa* L.) at Jimma, South Western Ethiopia. *Journal of Natural Sciences Research*, 9(10), 32-40.
- Ethiopian Agricultural Research Organization. (2004). Directory of released crop varieties and their recommended cultural practices.
- Gessesew, W.S., Mohammed, W., & Woldetsadik, K. (2015). Effect of nitrogen fertilizer rates and intra-row spacing on yield and yield components of onion (*Allium cepa* L. Var. cepa) under irrigation in Gode, South-Eastern Ethiopia. *International Journal of Plant Breeding and Crop Science*, 2(2), 46-54.
- Hazelton, P., & Murphy, B. (2016). Interpreting soil test results: What do all the numbers mean?. CSIRO Publishing.
- Kamble, B.M., & Kathmale, D.K. (2015). Effect of different levels of customized fertilizer on soil nutrient availability, yield, and economics of onion. *Journal of Applied and Natural Science*, 7(2), 817-821.

- Kihara, J., Tamene, L.D., Massawe, P., & Bekunda, M. (2015). Agronomic survey to assess crop yield, controlling factors, and management implications: a case-study of Babati in northern Tanzania. *Nutrient Cycling in Agroecosystems*, 102, 5-16.
- Masrie, B. (2019). Optimal rate of nitrogen and intra-row spacing for economical production of onion under irrigated farming system in Eastern Amhara Region, Ethiopia. *Current Research in Agricultural Sciences*, 6(2), 83-94
- Murphy, H.F. (1968). A report on the fertility status and other data on some soils of Ethiopia.
- Nasreen, S., Haque, M.M., Hossain, M.A., & Farid, A.T.M. (2007). Nutrient uptake and yield of onion as influenced by nitrogen and sulfur fertilization. *Bangladesh Journal of Agricultural Research*.
- Roy, R.N., Finck, A., Blair, G.J., & Tandon, H.L.S. (2006). Plant nutrition for food security: A guide for integrated nutrient management. *FAO Fertilizer and Plant Nutrition Bulletin*, 16(368), 201-214.
- Stephen, K., & Nicky, M. (2007). In-Service training on partial budgeting techniques. Mount Makulu, Lusaka, Zambia: Recommendations using simulation modeling. *Fertilizer Research*, 44, 37-49.
- Singh, R.P., Jain, N.K., & Poonia, B.L. (2000). Response of Kharif onion to nitrogen, phosphorus, and potash in eastern plains of Rajasthan. *The Indian Journal of Agricultural Sciences*, 70(12).
- Tadesse, T., Haque, I., & Aduayi, E.A. (1991). Soil, plant, water, fertilizer, animal manure, and compost analysis manual.
- Tegen, H., Jembere, M., Mihiretu, E., & Enyew, A. (2016). Influences of inter and intra-row spacing on yield, yield components, and morphological characteristics of onion (*Allium cepa* L.) at Western Amhara region. *African Journal of Agricultural Research*, 11(20), 1797-1804
- Tekle, G. (2015). Growth, yield, and quality of onion (*Allium cepa* L.) as influenced by intra-row spacing and nitrogen fertilizer levels in central zone of Tigray, Northern Ethiopia (Doctoral dissertation, Haramaya University).
- Thirupathi, R., Sunanda, E., Nagare, P.K., & Kumar, J.S. (2014). Effect of spacing and nitrogen levels on growth and yield of kharif white onion.
- United States Department of Agriculture. (1987). Textural soil classification study guide. National Employee Development Staff. Soil Conservation Service, United States Department of Agriculture, Washington DC, USA.
- Upper Awash Agro-Industry Enterprise. (2001). Progress Report 1996-2002, Addis Ababa, Ethiopia. Agricultural product 2001/2002, Addis Ababa, Ethiopia.
- Winch, T. (2006). Growing Food (pp. 1-103).

Yadav, R.L., Sen, N.L., & Yadav, B.L. (2003). Response of onion to nitrogen and potassium fertilization under semi-arid condition of Rajasthan. *Indian Journal of Horticulture*, 60(2), 176-178.

Yemane Kahsay, Y.K., Derbew Belew, D.B., &Fetien Abay, F.A. (2013). Effect of intra-row spacing on yield and quality of some onion varieties (*Allium cepa* L.) at Aksum, northern Ethiopia.

UNDER PEER REVIEW