

Case report

A case report of breast Myofibroblastoma: A bad looking benign tumor.

Abstract:

This study aims to analyze the relevance of the correct evaluation of the breast pathology, using a standardized method for reporting breast imaging studies, the Breast Imaging Report And Data System (BI-RADS). Myofibroblastoma(MFB) of the breast is a rare, benign, mesenchymal tumor of breast, can be a diagnostic challenge for the non-experienced general radiologist or radiology resident, due to its clinical, mammographic and ultrasonographic characteristics. In this report we present a case of 57-year-old women with a breast lump and non-specific imaging findings, through her mammographic and ultrasonographic evaluation, requiring histopathological correlation, making the diagnoses of MFB, therefore, lumpectomy was indicated as treatment.

Key words:Myofibroblastoma of the breast; benign breast tumor; BI-RADS; mammography; ultrasound.

1. INTRODUCTION

Breast cancer is the most diagnosed cancer in women in the United States of America and some other countries. Imaging techniques and categorization systems have advanced by great steps in the last decades, however there are some lesions that can still cause confusion and mimic malignancy. We present a case of a women with a left breast lump and pain; ultrasound and mammography were consistent for perform a histopathological correlation.

2. PRESENTATION OF THE CASE

57-year-old women with unremarkable health story presented to the Family Medicine Clinic for a 1 year of mild pain in the left breast, as well in the last month she notices the presence of a lump. Physical examination revealed a solid andmobile tumor in the upper external quadrant of the left mammary gland, a diagnostic mammogram and ultrasound was requested.

Mammogramfindings were negative for a nodules, malignant calcifications, asymmetries, or pathological lymph nodes (Image 1), otherwise the ultrasound revealed a hypoechogenic mass in the 02:00 Radio in the left breast (Image 2), the shape was irregular, and margins were non-circumscribed, parallel orientation, posterior shadowing was present and absent vascularity (Image 3).




Image 1.- Bilateral mammogram with breast heterogeneously dense.




Image 2.- Grayscale ultrasound in the 02:00 Radio in the left breast the shape was irregular, and margins were non-circumscribed, parallel orientation, posterior shadowing was present (Blue arrow).




Image 3.- Doppler Color ultrasound with absent vascularity.

Imaging findings suggest a probably malignant etiology, so it was decided to take an ultrasound guided biopsy of the tumor (Image 4). Three tissue samples of a filiform solid and white tissue were analyzed (Image 5).




Image 4.- Grayscale ultrasound guided biopsy of the tumor shows the needle (With the arrow) and the nodule (Blue arrow).



Image 5.-Macroscopic view shows three solid and white tissues.

The histological sections after hematoxylin and eosin staining show a myofibroblastic reactive proliferation without atypia. No malignant cells were found (Image 6 and 7).




Image 6.- Biphasic fibroepithelial proliferation of ducts and stroma, hematoxylin and eosin 40x.




Image 7.- Ductal epithelial cells without atypia, hematoxylin and eosin 400x.

Myofibroblastic tumor of the breast was diagnosed, and patient underwent lumpectomy without complications.

3.- DISCUSSION

Breast cancer is the most common cancer in women, the incidence of which continues to increase worldwide. Imaging screening has contributed to substantial reductions in breast cancer mortality, resulting in an increased prevalence of benign biopsies statistically [1].

Irregular hypoechoic masses on breast ultrasound are usually considered suspicious lesions. If the lesions combine other features of malignancy such as spiculated margin, nonparallel orientation, and posterior shadowing, they are considered moderate and highly suspicious for malignancy (BI-RADS categories 4b and 4c) or highly suggestive of malignancy (BI-RADS category 5) [2]. Such lesions are initially determined to be suspicious, at which point sonographically guided core needle biopsies are performed [3]. In this case we found a mass suspicious of malignancy.

MFB is a rare, benign, mesenchymal tumor of breast [4]. MFB was first described by Wargotz et al in 1987 as a distinct stromal tumor of the breast [5]. There have been <90 case reports of mammary MFB reported till date after being first described as a distinct entity in 1987 [6]. More recently, these tumors have been noted at extramammary sites as well, and the term Mammary-type MFB is often used when referring to this group of tumors [5]. While earlier studies reported a

male predominance, this tumors may also occur in female patients [5], as in the case of this patient.

Clinically, these tumors present as slow growing, painless masses, without evidence of local lymphadenopathy, in middle-aged patients [4].

The typical imaging appearance of breast MFB is a well-circumscribed, gently lobulated mass with macroscopic fat and variable density on mammography [7].

Breast MFB demonstrates similar benign imaging findings on ultrasound, a parallel, circumscribed, heterogeneous or hypoechoic mass with variable posterior features, soft elastography features and mild internal vascularity. Posterior features, if present, are generally posterior acoustic shadowing secondary to acoustic impedance caused by the increased internal cellular density of the mass relative to the surrounding normal fat lobules and fibroglandular tissue. The vascularity is reported in the literature as predominantly peripheral vessels [7]. We found a mass with shape was irregular, and margins were non-circumscribed, parallel orientation, posterior shadowing was present and absent vascularity

Microscopic examination typically reveals uniform, slender spindle cells morphologic features of myofibroblast admixed with broad bands of hyalinized collagen[5]. Lesional cells of breast MFB show immunoreactivities for myofibroblastic markers, such as desmin, smooth muscle actin, and muscle specific actin. Besides, most MFB are positive for CD34, BCL-2, vimentin and hormonal receptors like estrogen receptor, progesterone receptor and androgen receptor[7].

Though MFB of the breast is a benign tumor and local excision is most likely curable[7].

Spindle cell lesions of the breast comprise a wide variety of conditions, either benign, locally aggressive, or malignant. Among the differential diagnoses are Pseudoangiomatousstromalhyperplasia, Fibromatosis, Spindle cell lipoma, Leiomyoma, Invasive lobular carcinoma and Metaplastic spindle cell carcinoma

4.- CONCLUSION

Imaging studies are constantly evolving, prioritizing the most important public health problems, such as breast cancer.

The American College of Radiology has proposed a standardized method for reporting breast imaging studies, using the BI-RADS, currently based on the fifth edition, categorizing lesions, and providing predictive values for malignancy.

Breast MFB is a rare, benign, and asymptomatic tumor, however, it can sometimes show characteristics of malignancy, through mammographic and ultrasonographic evaluation.

The clinical and imaging characteristics evaluated in this patient, were suggestive for malignancy, so it was required histopathological correlation, and lumpectomy was subsequently performed as definitely treatment.

This case is a clear example of the multidisciplinary work that must be carried out, for the diagnosis and treatment of breast pathology.

Consent

We as authors declare that written informed consent was obtained for publication of this case report and accompanying images.

Ethical approval

As per international standard or university standard written ethical approval has been collected and preserved by the author.

References

- [1] Kim MJ, Kim D, Jung W, Koo JS. Histological analysis of benign breast imaging reporting and data system categories 4c and 5 breast lesions in imaging study. *Yonsei med j.* 2012 Nov 1;53(6):1203-10. Doi: 10.3349/ymj.2012.53.6.1203. Pmid: 23074123; pmcid: pmc3481383.
- [2] Kim YR, Kim HS, Kim HW. Are irregular hypoechoic breast masses on ultrasound always malignancies?: a pictorial essay. *Korean j Radiol.* 2015 nov-dec;16(6):1266-75. Doi: 10.3348/kjr.2015.16.6.1266. Epub 2015 oct 26. Pmid: 26576116; pmcid: pmc4644748.
- [3] Cho SH, Park SH. Mimickers of breast malignancy on breast sonography. *J ultrasound med.* 2013 Nov;32(11):2029-36. Doi: 10.7863/ultra.32.11.2029. Pmid: 24154908.
- [4] Abdul-ghafar J, Ud din N, Ahmad Z, Billings SD. Mammary-type myofibroblastoma of the right thigh: a case report and review of the literature. *J med case rep.* 2015 Jun 2;9:126. Doi: 10.1186/s13256-015-0601-0. Pmid: 26033228; pmcid: pmc4470027.
- [5] Fritchie KJ, Carver P, Sun Y, Batiouchko G, Billings SD, Rubin BP, Tubbs RR, Goldblum JR. Solitary fibrous tumor: is there a molecular relationship with cellular

angiofibroma, spindle cell lipoma, and mammary-type myofibroblastoma? Am j clin pathol. 2012 jun;137(6):963-70. Doi: 10.1309/ajcpqeg6ynn6cna1. Pmid: 22586056.

[6] khatib Y, Pandey V, Khade AL, Pandey R. Myofibroblastoma of the breast: a rare cause of breast lump in a postmenopausal woman. J midlife health. 2018 jan-mar;9(1):47-49. Doi: 10.4103/jmh.jmh_59_17. Pmid: 29628731; pmcid: pmc5879850.

[7] Yan M, BomeislP, Gilmore H, Sieck L, Kuchta Z, HarbhajankaA. Clinicopathological and radiological characterization of myofibroblastoma of breast: a single institutional case review. Ann diagnpathol. 2020 oct;48:151591. Doi: 10.1016/j.anndiagpath.2020.151591. Epub 2020 aug 15. Pmid: 32829069.

UNDER PEER REVIEW