

Effect of different levels of nitrogen and potassium on yield and quality of potato

Abstract

The research experiment was carried out at Nalanda College of Horticulture, Noorsara during 2021-22 to find out the effect of different levels of nitrogen and potassium on growth and yield of potato. Findings of research showed that among the individual nitrogen and potassium levels, N_5 and K_4 performed better in respect of, number of tubers per plant (7.85 and 6.80), average tuber weight (62.28 g and 67.27 g), tuber weight per plant (656.71 g and 473.45 g), tuber yield (270.95 q/ha and 216.73 q/ha), diameter of tuber (5.21 cm and 3.98 cm), vitamin C content in tuber (14.15 mg/100g and 11.89 mg/100g), fresh weight (63.68 and 109.79), dry weight (12.89 and 22.12), moisture content of fresh tuber (76.09 and 91.07). In case of interaction effect of $N \times K$, the treatment of $N_5 \times K_4$ pronounced significantly to number of tubers per plant (11.44), single tuber weight (110.19 g), plant tuber weight (962.00 g), tuber yield (360 q/ha), diameter of tuber (8.10 cm), vitamin C content in tuber (18.50 mg/100 g), fresh weight (109.79 g), dry weight (22.12 g), moisture content of fresh tuber (91.07 g). Potato var. Kufri Lima yielded the maximum (360.76 q/ha) in interaction of $N_5 \times K_4$ when applied with 160 kg/ha nitrogen and 120 kg/ha potassium.

Key word: potato, nitrogen, solanum.

Introduction

Potato (L. (*Solanum tuberosum* L.), having diploid chromosome number ($2n=48$), a member of the solanaceae family, is the most important crop in the vegetable class. It is the most extensively grown tuber crop ranking fourth after rice, wheat and maize in world. It is a vital source of nutrients for human population and is taken almost daily by over a billion people, fresh or refined (FAOSTAT, 2017). It is believed to be used as global food security. About 40%, 35% and 25% of potato are grown in Europe, different developed countries and rest of the world respectively (Gull *et al.*, 2011). Being starchy crop, it produces high production of starch per unit area per unit time. The 100 gm of fresh tuber contains 70-80% water, 20.6%

carbohydrates, 2.1% protein, 0.3% fat, 1.1% crude fibre and 0.9% ash. Potatoes are rich source of carbohydrate, vitamin C, minerals and essential amino acids like leucine, tryptophan and isoleucine (Bist and Sharma, 1997). Potato is considered as a highly nutritious food and it gives high productivity per unit area and time as compared to many food grains. The crop matures faster within 90 to 100 days, and provide good tuber yield even in just 60 days. Overall, potato production is one of the most efficient means of converting plant, land, water and labour into a palatable and nutritious food (Sahadevan, 2007). Due to its uniqueness, potato has been declared as “Future food crop” by Food and Agricultural Organization, Rome in the year 2008 (Hussain, 2016). The potato tuber is an enlarged swollen underground stem with variable shapes and sizes. The swelling of the tuber is due to the translocation and storage of photosynthetes (carbohydrates), which retain its highest with maturation of the aerial element of the plant. Hence, tuber boom and improvement of potato in general depends on the presence of ample foliage that produces the fundamental assimilates which directly depend on fertilization.

The most important nutrient in fertilizer is nitrogen which is a growth stimulant for plants. Nitrogen is an essential constituent of various metabolically active compounds like amino acids, proteins, nucleic acids, pyrimidines, flavines, purines, nucleoproteins, enzymes, protein, protoplasm and chlorophyll (Panda, 2006). So its availability is important to plant growth and yield. Nitrogen applications can increase dry matter and protein content of tuber, quality, bulking rate, haulm growth and total tuber yield (Belanger *et al.*, 2002). There is a positive impact of N on photosynthetic rate, leaf expansion, total number of leaves, canopy growth and shoot dry matter. The higher nitrogen content can be attributed to increased root proliferation due to effect of nitrogen on cellular activities and the translocation of some growth-stimulating compounds to roots, resulting in improved tuber growth and nutrient absorption (Sharma and Sood, 2002).

Potassium (K) is the 3rd essential nutrient element for plant growth and taken up from the soil solution by the plant roots in the form of potassium ion (K^+). Potassium is very mobile within the plant and its deficiency symptoms in plants appear first in the older leaves (Tisdale *et al.*, 1993). Potassium is directly involved in enzyme activation, maintenance of water status, energy relations, and translocation of assimilates and protein synthesis. K regulates cellular turgid pressure to avoid wilting, which in turn controls the stomata opening and hence greatly enhances drought tolerance (McCarty, 2005). The response of crop to potassium increases significantly in the presence of nitrogen (Sharma, 1989). Potatoes require high amounts of potassium fertilizer for optimum growth, production and tuber quality. Potassium promotes leaf development, especially in the early stages of growth and for longer periods of time during tuber bulking. The nutrients (Al-Moshileh *et al.*, 2004) management plans including their economic cost and return were pondered to obtain optimum yield and maximum profit. Potassium affects tuber quality characteristics such as ascorbic acid, total soluble solids, specific gravity, and tuber bulking rate, in addition to enhancing resistance to pests and diseases, which results in a better market price for the produce. Potato plants require much more potassium than many other vegetable crops; therefore, it is also regarded as an indicator crop for K availability (Al-Moshileh, 2004).

Positive effects of N and K interaction in terms of crop productivity and economics, but the balance of N and K application is not appropriately practiced in many parts of the world. Inadequate application of potassium (K) combined with over application of nitrogen (N) is a serious problem in modern intensive agricultural production systems. Higher yields and crop quality can be obtained at optimal N:K nutritional ratios. Optimal N and K application is essential for best nutrient management in agriculture. Application of K facilitates the uptake and transport of nitrate towards aerial parts of the plant, which in turn enhances the activities of nitrate assimilating enzymes (K cycling and recycling play an

important part in NO_3^- translocation from root to shoot as counter ion and assimilate loading in the phloem (Maathuis, 2007). Cycling and recycling of K^+ increased with increasing shoot growth rate, which is in accordance with the suggested role of K^+ for charge balance of NO_3^- in the xylem and organic acids in the phloem (Engels and Kirkby, 2001). The present research was undertaken to study the effect of combined and individual effect of nitrogen and potassium on growth and yield of potato and also to determine the optimum levels of nitrogen and potassium individual or in combination, which would be best for the growth of potato.

Materials and method

The present experiment was carried out during Rabi season of 2022-23 at the Vegetable Research Field, Department of Vegetable Science, Nalanda College of Horticulture, Noorsarai, Nalanda (Bihar). The experiment was laid out in factorial randomized block design with twenty treatments in three replications. Potato cultivar Kufri Lima was taken as test crop. The treatments T_1 - $\text{N}_0:\text{P}_0:\text{K}_0$, T_2 - $\text{N}_0:\text{P}_{80}:\text{K}_{80}$, T_3 - $\text{N}_0:\text{P}_{80}:\text{K}_{100}$, T_4 - $\text{N}_0:\text{P}_{80}:\text{K}_{120}$, T_5 - $\text{N}_{100}:\text{P}_{80}:\text{K}_0$, T_6 - $\text{N}_{100}:\text{P}_{80}:\text{K}_{80}$, T_7 - $\text{N}_{100}:\text{P}_{80}:\text{K}_{100}$, T_8 - $\text{N}_{100}:\text{P}_{80}:\text{K}_{120}$, T_9 - $\text{N}_{120}:\text{P}_{80}:\text{K}_0$, T_{10} - $\text{N}_{120}:\text{P}_{80}:\text{K}_{80}$, T_{11} - $\text{N}_{120}:\text{P}_{80}:\text{K}_{100}$, T_{12} - $\text{N}_{120}:\text{P}_{80}:\text{K}_{120}$, T_{13} - $\text{N}_{140}:\text{P}_{80}:\text{K}_0$, T_{14} - $\text{N}_{140}:\text{P}_{80}:\text{K}_{80}$, T_{15} - $\text{N}_{140}:\text{P}_{80}:\text{K}_{100}$, T_{16} - $\text{N}_{140}:\text{P}_{80}:\text{K}_{120}$, T_{17} - $\text{N}_{160}:\text{P}_{80}:\text{K}_0$, T_{18} - $\text{N}_{160}:\text{P}_{80}:\text{K}_{80}$, T_{19} - $\text{N}_{160}:\text{P}_{80}:\text{K}_{100}$, T_{20} - $\text{N}_{160}:\text{P}_{80}:\text{K}_{120}$. The crop was planted with a spacing of 50cm×20cm and plot size of 3.0m×3.0m. Uniform cultural practices recommended for broccoli was followed. The recorded data of the trial were subjected to statistical analysis and the results were documented, analysed and presented in tabular form.

Results and discussion

Effect of N, K and N × K on growth and yield parameters of potato

The maximum number of tubers per plant (9.56) was recorded significantly with N_5 (160 kg/ha) and minimum number of tubers (6.84) in the absence of nitrogen N_1 . With the application K_4 (120 kg/ha) the maximum mean value of number of tubers per plant (8.92) was significantly

observed. The minimum value of number of tubers per plant (7.46) was showed in K_1 (0 kg/ha). The maximum number of tubers per plant (11.44) was recorded with the application of $N_5 \times K_4$ (160 \times 120 kg/ha). However, the minimum number of tubers per plant (6.74) was noted in the $N_2 \times K_1$ ($N_{100} \times K_0$).

Table-1: Effect of different levels of nitrogen and potassium on number of tuber per plant

Treatment	K_1 (C=0kg/ha)	K_2 (80kg/ha)	K_3 (100kg/ha)	K_4 (120kg)	Mean
N_1 (C=0kg/ha)	6.87	6.84	6.86	6.80	6.84
N_2 (100kg/ha)	6.74	7.83	8.14	8.19	7.72
N_3 (120kg/ha)	7.91	7.83	8.70	8.61	8.26
N_4 (140kg/ha)	7.96	8.78	9.33	9.56	8.91
N_5 (160kg/ha)	7.85	9.11	9.84	11.44	9.56
Mean	7.46	8.08	8.57	8.92	
	SEm(\pm)		CD		
N	0.20		0.57		
K	0.18		0.51		
N \times K	0.40		1.15		

The maximum value of single tuber weight (91.54 g) was noticed significantly with N_5 (160 kg/ha) whereas the minimum single tuber weight (57.55 g) was recorded in the absence of nitrogen (N_1). With the application of K_4 (120 kg/ha), the value of single tuber weight (91.07 g) was significantly observed. The minimum value of single tuber weight (55.27 g) was showed in K_1 (0 kg/ha). The maximum value of single tuber weight (110.19 g) was recorded with the application of $N_5 \times K_4$ (160 \times 120 kg/ha) which was at par with $N_4 \times K_4$ and $N_5 \times K_2$. However, the minimum value of single tuber weight (46.60 g) was noted in the $N_1 \times K_1$ ($N_0 \times K_0$).

Table-2: Effect of different levels of nitrogen and potassium on single tuber weight (g)

Treatment	K_1 (C=0kg/ha)	K_2 (80kg/ha)	K_3 (100kg/ha)	K_4 (120kg)	Mean
N_1 (C=0kg/ha)	46.60	53.82	62.52	67.27	57.55
N_2 (100kg/ha)	52.00	65.57	71.47	77.50	66.64
N_3 (120kg/ha)	56.67	74.14	81.94	94.25	76.75
N_4 (140kg/ha)	58.80	82.53	92.87	106.13	85.08
N_5 (160kg/ha)	62.28	98.77	94.93	110.19	91.54
Mean	55.27	74.97	80.75	91.07	
	SEm(\pm)		CD		

N	2.01	5.74
K	1.79	5.14
N×K	4.01	11.48

The maximum mean value of tuber weight per plant (811.18 g) was found significantly with N₅(160 kg/ha). The minimum mean tuber weight per plant(382.38g) was noticed in the absence of nitrogen(N₁). With the application of K₄(120 kg/ha) the maximum mean value of tuber weight per plant (742.34 g) was significantly observed. The minimum mean value of tuber weight per plant(464.73g) was showed in K₁(0Kkg/ha). The maximum mean value of tuber weight per plant (962 g) was observed with the application of N₅× K₄(160 kg/ha × 120 kg/ha) and N₅× K₂. However, the minimum mean value of tuber weight per plant (280.95 g) was noted in the N₁×K₁(N₀×K₀).

Table-3: Effect of different levels of nitrogen and potassium on tuber weight per plant

Treatment	K₁(C=0kg/ha)	K₂(80kg/ha)	K₃(100kg/ha)	K₄(120kg)	Mean
N₁(C=0kg/ha)	280.95	354.91	420.20	473.45	382.38
N₂(100kg/ha)	327.51	488.61	634.88	642.89	523.47
N₃(120kg/ha)	422.20	678.41	691.22	798.29	647.53
N₄(140kg/ha)	636.87	697.54	802.76	835.06	743.06
N₅(160kg/ha)	656.11	769.13	857.46	962.00	811.18
Mean	464.73	597.72	681.30	742.34	
	SEm(±)		CD		
N	16.88		48.33		
K	15.10		43.23		
N × K	33.76		NS		

The maximum mean value of tuber yield (339.35 q/ha) was found significantly with N₅(160kg/ha) which was at par with N₄(140kg/ha). The minimum mean tuber yield(178.55 q/ha) was noticed in the absence of nitrogen(N₁). The maximum mean tuber yield(323.02q/ha) was significantly observed in K₄(120 kg/ha). The minimum mean tuber yield (227.03 q/ha) was recorded without application of potassium(K₁). The maximum mean value of tuber yield (372.36 q/ha) was found with the application of N₄× K₄(140×120 kg/ha) which was at par with N₅× K₂, N₃× K₄, N₅× K₄, N₄× K₃ and N₅× K₃. However, the minimum mean value

of tuber yield(130.01q/ha)was recordedinthe $N_1 \times K_1(N_0 \times K_0)$.

Table-4: Effect of different levels of nitrogen and potassium on tuber weight (q/ha).

Treatment	K ₁ (0 kg/ha)	K ₂ (80 kg/ha)	K ₃ (100 kg/ha)	K ₄ (120 kg)	Mean
N ₁ (0 kg/ha)	130.01	167.98	199.49	216.73	178.55
N ₂ (100 kg/ha)	239.37	251.54	284.34	297.53	268.20
N ₃ (120 kg/ha)	242.27	274.86	308.38	367.70	298.31
N ₄ (140 kg/ha)	252.55	303.21	358.40	372.36	321.63
N ₅ (160 kg/ha)	270.95	369.85	355.85	360.76	339.35
Mean	227.03	273.49	301.29	323.02	
	SEm(±)		CD		
N	6.71		19.22		
K	6.00		17.19		
N × K	13.43		38.44		

It is found that application of N and K fertilizers significantly increased growth and yield of tuber. Increased application of different levels of N and K found significant effect on growth and yield of tuber in potato. Interaction effect of N×K was also found to be significant on growth and yield of tuber. Similar findings are supported by Sezeket *et al.*, (2018). Plant height is an indicator of vegetative growth. The addition of nutrients enhances the soil fertility and productivity levels which results in healthy crop. The minimum plant height might be due to the result of unavailability of nitrogen and other nutrients required by the plants for their normal growth and development. It was observed that an increase in nitrogen levels positively affected the plant height character which might be due to the role of nitrogen for cell division, cell enlargement and protein synthesis. Potassium nutrient in metabolism and many processes needed to promote plant vegetative growth and development. Positive response was registered by Ali *et al.* (2003) and Hossain *et al.* (2009) who found that increasing potassium fertilization increased shoot height, number of leaves per plant and shoot fresh weight. Application of K increases plant height, crop vigour and imparts resistance against drought, frost and diseases. Potassium increases leaf expansion particularly at early stages of growth and extends leaf area duration by delaying leaf shedding near

maturity. It increases both the rate and duration of tuber bulking. Its application activates a number of enzymes involved in photosynthesis, carbohydrate and protein metabolism, and assists in the translocation of carbohydrates from leaves to tubers. Potassium increases the size but not the total number of tubers (Trehan *et al.*, 2001). Potassium application thus increases yield by the formation of larger sized tubers. Zhao-Hui *et al.* (2008) found that K increased the yields of tomato at most N rates. Nevertheless, the N_0K_0 treatment produced the lowest yields. The interaction between N and K showed that with higher rates of N application, the effect of K became more significant with increasing application rate. This clearly indicated that as soon as a higher yield potential through better N supply was implemented, there was a rapid depletion of soil K. According to Mullins *et al.* (1991) and Cassman *et al.* (1992), the relationship between N and K determines the balance between vegetative growth, fruit quality and reproductive processes. Increased N and K forced higher yield which is also supported by Taya *et al.* (1994) and found that the highest tuber yield (>75 g) t/ha was related to the third treatment (i.e. 150 kg N/ha) and the lowest tuber yield was related to the first treatment (100 kg N/ha). Higher fertilizer response may be linked to the increase in total leaf area which in turn increased the amount of solar radiation intercepted and more photo-assimilates might have been produced and assimilated to the tubers.

Effect of N, K and of N × K on quality parameters of potato tuber

The maximum mean value of diameter of tuber (6.63 cm) was found significantly with N_5 (160 kg/ha). The minimum mean diameter of tuber (3.88 cm) was noticed in the absence of nitrogen (N_1). The maximum mean value of diameter of tuber (6.13 cm) was significantly observed in K_4 (120 kg/ha) which was significantly superior to all potassium levels. The minimum mean diameter of tuber (4.53 cm) was showed in 0K kg/ha (K_1). The maximum mean value of diameter of tuber (8.10 cm) was found with the application of $N_5 \times$

$K_4(160 \times 120 \text{ kg/ha})$ which was significantly superior to all $N \times K$ interaction. However, the minimum mean value of diameter of tuber (3.73 cm) was recorded in the $N_1 \times K_1 (N_0 \times K_0)$.

Table-5: Effect of different levels of nitrogen and potassium on diameter of tuber (cm)

Treatment	$K_1(C=0\text{kg/ha})$	$K_2(80\text{kg/ha})$	$K_3(100\text{kg/ha})$	$K_4(120\text{kg})$	Mean
$N_1(C=0\text{kg/ha})$	3.73	3.84	3.96	3.98	3.88
$N_2(100 \text{ kg/ha})$	3.82	4.77	5.12	5.51	4.81
$N_3(120 \text{ kg/ha})$	4.80	4.91	5.66	6.19	5.39
$N_4(140 \text{ kg/ha})$	5.07	5.81	6.33	6.88	6.02
$N_5(160 \text{ kg/ha})$	5.21	6.29	6.94	8.10	6.63
Mean	4.53	5.12	5.60	6.13	
	SEm(\pm)		CD		
N	0.14		0.40		
K	0.13		0.36		
$N \times K$	0.28		0.81		

The maximum mean value of vitamin C content (15.97 mg/100 g) was observed significantly with $N_5(160 \text{ kg/ha})$. The minimum mean vitamin C content (11.17 mg/100 g) was noticed in the absence of nitrogen (N_1). The maximum mean value of vitamin C content (14.09 mg/100 g) was significantly noticed in $K_4(120\text{kg/ha})$ which was at par with $K_3(100\text{kg/ha})$. The minimum mean vitamin C content (12.68 mg/100 g) was noted in $K_1(0\text{kg/ha})$. The maximum mean value of vitamin C content (18.50 mg/100 g) was found with the application of $N_5 \times K_4(160 \times 120 \text{ kg/ha})$ which was significantly superior to all $N \times K$ interactions. However, the minimum mean value of vitamin C content (10.46 mg/100 g) was recorded in the $N_1 \times K_1 (N_0 \times K_0)$.

Table-6: Effect of different levels of nitrogen and potassium on vitamin c content (mg per 100 gm) in tuber

Treatment	$K_1 (0 \text{ kg/ha})$	$K_2 (80 \text{ kg/ha})$	$K_3 (100 \text{ kg/ha})$	$K_4 (120 \text{ kg})$	Mean
$N_1 (0 \text{ kg/ha})$	10.46	10.97	11.36	11.89	11.17
$N_2 (100 \text{ kg/ha})$	11.22	12.05	12.34	12.98	12.15
$N_3 (120 \text{ kg/ha})$	13.44	13.54	13.87	11.83	13.17
$N_4 (140 \text{ kg/ha})$	14.12	14.42	14.94	15.26	14.68
$N_5 (160 \text{ kg/ha})$	14.15	15.57	15.67	18.50	15.97
Mean	12.68	13.31	13.64	14.09	

	SEm (\pm)	CD
N	0.32	0.92
K	0.29	0.83
N \times K	0.64	1.84
N \times K	3.46	9.91

The maximum mean value of fresh weight of tuber (91.43 g) was observed significantly with N_5 (160 kg/ha). The minimum mean fresh weight of tuber (57.78g) was noticed in the absence of nitrogen (N_1). The maximum mean value of fresh weight of tuber (91.01 g) was significantly noticed in K_4 (120 kg/ha). The minimum mean fresh weight of tuber (55.51g) was noted in K_1 (0 kg/ha). The maximum mean value of fresh weight of tuber (109.79g) was found significantly with the application of $N_5 \times K_4$ (160 \times 120 kg/ha) which at par with $N_4 \times K_4$. However, the minimum mean value of fresh weight of tuber (45.94 g) was shown in the $N_1 \times K_1$ ($N_0 \times K_0$).

Table-7: Effect of different levels of nitrogen and potassium on fresh weight of tuber.

Treatment	K_1 (C=0kg/ha)	K_2 (80kg/ha)	K_3 (100kg/ha)	K_4 (120kg)	Mean
N_1 (C=0kg/ha)	45.94	54.23	62.66	68.29	57.78
N_2 (100kg/ha)	51.56	65.70	71.64	78.46	66.84
N_3 (120kg/ha)	56.75	74.45	81.96	93.83	76.75
N_4 (140kg/ha)	59.61	82.71	92.20	104.69	84.80
N_5 (160kg/ha)	63.68	96.99	95.25	109.79	91.43
Mean	55.51	74.82	80.74	91.01	
	SEm (\pm)		CD		
N	1.73		4.96		
K	1.55		4.43		
N \times K	3.46		9.91		

The maximum mean value of dry weight of tuber (18.61g) was found significantly with N_5 (160 kg/ha). The minimum mean dry weight of tuber (12.08g) was recorded in the absence of nitrogen (N_1). The maximum mean value of dry weight of tuber (18.13g) was significantly noticed in K_4 (120kg/ha). The minimum mean dry weight of tuber (10.91g) was with K_1 (0 kg/ha). The maximum mean value of dry weight of

tuber(22.12 g) was found significantly with the application of $N_5 \times K_4$ (160 × 120 kg/ha) which atpar with $N_4 \times K_4$. However, the minimum mean value of dry weight of tuber (9.08g) was showed in the $N_1 \times K_1$ ($N_0 \times K_0$).

Table-8: Effect of different levels of nitrogen and potassium on dry weight of tuber (g).

Treatment	K_1 (C=0kg/ha)	K_2 (80kg/ha)	K_3 (100kg/ha)	K_4 (120kg)	Mean
N_1 (C=0kg/ha)	9.08	11.07	13.10	15.05	12.08
N_2 (100kg/ha)	9.65	13.01	15.06	17.49	13.80
N_3 (120kg/ha)	10.78	15.02	17.65	16.28	14.93
N_4 (140kg/ha)	12.16	16.15	17.13	19.69	16.28
N_5 (160kg/ha)	12.89	20.04	19.37	22.12	18.61
Mean	10.91	15.06	16.46	18.13	
	SEm(±)		CD		
N	0.36		1.03		
K	0.32		0.93		
N × K	0.72		2.07		

The maximum mean value of moisture matter content in tuber (82.71 %) was found significantly with N_1 (0 kg/ha) which was at par with N_2 (79.49 %), N_3 (78.31 %) and N_5 (78.31%). The minimum mean moisture matter content in tuber (74.30%) was recorded in the (N_4). The maximum mean value of moisture matter content in tuber (81.98%) was significantly noticed in K_1 (0 kg/ha). The minimum mean moisture matter content in tuber (78.86%) was with 120 kg/ha (K_4). The maximum mean value of moisture matter content in tuber (93.03%) was found significantly with the application of $N_1 \times K_1$. The minimum mean value of moisture matter content in tuber (70.87 %) was recorded in the $N_4 \times K_3$ ($N_{140} \times K_{100}$ kg/ha).

Table-9: Effect of different levels of nitrogen and potassium on moisture content in tuber.

Treatment	K_1 (C=0kg/ha)	K_2 (80kg/ha)	K_3 (100kg/ha)	K_4 (120kg)	Mean
N_1 (C=0kg/ha)	93.03	81.14	77.00	79.69	82.71
N_2 (100kg/ha)	83.10	78.45	79.35	77.05	79.49
N_3 (120kg/ha)	79.15	76.25	75.51	74.88	76.45
N_4 (140kg/ha)	78.52	76.19	70.87	71.60	74.30
N_5 (160kg/ha)	76.09	74.83	71.26	91.07	78.31
Mean	81.98	77.37	74.80	78.86	
	SEm(±)		CD		

N	1.76	5.05
K	1.58	4.52
N×K	3.53	10.10

It is quite clear that individual effect of N and K responded significantly on ascorbic acid, fresh weight, dry weight and moisture content of tuber. This might be due to increase in uptake of nutrients nitrogen and potassium which promoted ascorbic acid content which supported by Tomar and Singhal(2007). The positive influence of potassium was due to the close relationship between carbohydrates metabolism and formation of ascorbic acid. Awaadet *al.* (2016) found that applications of N and K at higher levels significantly increased the fresh and dry weights of the lettuce plants. These results are in agreement with the findings of Gülser (2005) who indicate that the yield of spinach was increased by increasing the rates of N fertilizer. Nurzyńska-Wierdak (2009) indicated that the increase in the amounts of N and K application generally contributed to an increase in fresh leaf weight and yield. Abdel-Motagally and Osman (2010) found that increasing N and K fertilizer rates resulted to a significant increase in yield compared to the other treatments. Application of K has a direct influence on tuber quality (Trehanet *al.*, 2001).

References

- Abdel-Motagally, F.M.F. and Osman E. A., 2010. Effect of nitrogen and potassium fertilization combinations on productivity of two sunflower cultivars under East-of El-ewinate conditions. *American-Eurasian Journal of Agricultural & Environmental Sciences*, 8(4): 397-401.
- Ali, M. A., Hossain, M. A., Mondal, M. F. and Farooque, A. M., 2003. Effect of nitrogen and potassium on yield and quality of carrot. *Pak. J. Biol. Sci.*, 6: 1574-157.
- Al-Moshileh, A. M. and Errebi, M. A., 2004. Effect of various potassium sulphate rates on growth, yield and quality of potato grown under sandy soil and arid

- conditions. Potassium and fertigation development in West Asia and North Africa, pp. 24-28. Rabat, Morocco: IPI Regional Workshop.
- Awaad, M.S., Badr, R.S., Badr, M.A. and Abd-elrahman, A.H., 2016. Effects of different nitrogen and potassium sources on lettuce (*Lactuca sativa* L.) yield in a sandy soil. *Eurasian Journal of Soil Science*, Vol. 4 (5):299-306.
- Belanger, G., Walsh, J.R., Richards, J.E., Milburn, P.H. and Ziadi, N. 2002. Nitrogen fertilization and irrigation affects tuber characteristics of two potato cultivars. *Am. J. Pot. Res.*, 79:269-279.
- Bist, B.S. and Sharma, H.C., 1997. Potato statistics India and the World Technical Bulletin, CPRI, Shimla. 40:124.
- Cassman, K.G., Roberts, B.A., Bryant, D.C., 1992. Cotton response to residual fertilizer potassium on vermiculitic soil, organic matter and sodium effects. *Soil Sci. Soc. Am. J.*, 56:823-830.
- Engels, C. and Kirkby, E.A., 2001. Cycling of nitrogen and potassium between shoot and roots in maize as affected by shoot and root growth. *J. Plant Nutr. Soil Sci.*, 164:183-191.
- FAOSTAT, 2017. World Potato Production Quantity (Tonnes), Yields and Harvested Areas for 2017. <http://fenix.fao.org/faostat/internal/en/#data/QC/visualize>
- Gul, Z., Khan, A.A., Jamil, K. Review, 2011. Study of Potato leaf roll virus (PLRV) potato in Pakistan. *Canadian Journal on Scientific and Industrial Research*, 2(1):24.
- Gulser, F., 2005. Effects of ammonium sulphate and urea on NO_3^- and NO_2^- accumulation, nutrient contents and yield criteria in spinach. *Scientia Horticulturae*, 106 (3): 330-340.
- Hussain, A. K., Islam, M. M., Bari, M. R., Amin, M. S., M. H. A. and Kabir, M. A., 2009. Effect of mulching and levels of potassium on growth and yield of carrot. *Bangladesh Res. Pub. J.*, 32: 963-970

- Hussain, T., 2016. Potatoes: ensuring food for the future. *Adv Plants Agric Res.*, 3(6): 178-182.
- Maathuis, F.J.M., 2007. Monovalent Cation Transporters; Establishing a Link Between Bioinformatics and Physiology. *Plant Soil*, 301:1-5.
- Mc Carty, L.B., 2005. Best golf course management practices. 2nd Edition, Prentice Hall Inc. Upper Saddle River, New Jersey, USA. 896 p. Meter as Decision Tool for Nitrogen Management of Potato. *Agron J.*, 98: 496-506.
- Muliins, G.L., Burmester, C.H., Reeves, D.W. 1991. Cotton response to the deep placement of potassium on Alabama soils. In: Proceedings of the Beltwide Cotton Conferences, January 8-12. pp. 922-924. San Antonio, National Cotton Council of America, Memphis.
- Nurzyński, J Michałojc, Z., 1998. Plonowanie pomidora uprawianego w węłmineralnej w zależności od nawożenia potasowego. *Zesz. Nauk AR Kraków*, 333(57): 235-239.
- Nurzyńska-Wierdak, R., 2009. Growth and yield of garden rocket (*Eruca sativa* Mill) affected by nitrogen and potassium fertilization. *Acta Scientiarum Polonorum – Hortorum Cultus*, 8(4): 23-33.
- Panda, S.C., 2006. Management and Organic farming. Agrobios (India). Jodhpur.
- Pervez, M., Ayyub, M., Shabeen, M., and Noor, M. 2013. Determination of Physiochemical Characteristics of Potato Crop Regulated by Potassium Management. *Pakistan Journal of Agriculture Sciences.*, 50(4), 611-615
- Sahadevan, K.G. 2007. Advantages of Commodity Futures Trading Through Electronic Trading Platform for Farmers of Uttar Pradesh: A Study of Potato and Mentha, Multi Commodity Exchange of India Limited, Mumbai.

- Sezek, M., E. Ozturk and V. Aslan, 2018. The effect of different nitrogen fertilizer form and sowing date on some morphological characters of oilseed sunflower in high elevation condition. *Fresenius Env. Bull.* 27(12):8486- 8492.
- Sharma R.C., Sood, M.C. 2002. Nitrogen and potassium interaction on the tuber yield, quality and organic carbon status of Shimla soils. *In Potato, Global Research & Development*. Proceedings of the Global Conference on Potato, New Delhi, India, Indian Potato Association 1999, 2002;2:843-851
- Sharma, U.C., 1989. Potato yield and nutrient uptake as influenced by fertility levels at different location in Meghalaya. *Journal of Indian potato Association*. 16: 91-95.
- Taya, J.S., Malik, Y.S., Pandita, M.L., Khurana, S.C., 1994. Fertilizer management in potato based system: Growth and yield of potato. *J. Ind. Potato Assoc.*, 21(3-4): 184-188.
- Tisdale, S.L., Nelson, W.L., Beaton, J.D. and Halvin, J.L. 1993. *Soil Fertilizers* (5th Edn.). Macmillan Publishing Co. Inc. New York, pp. 109-229.
- Tomar, I.S. and Singhal, H.C., 2007. Effect of varying levels of nitrogen and different dates of planting on fruit quality of Rabi tomato. *JNKVV. Res. J.* 41 (2): 219-223.
- Trehan, S.P., Roy, S.K. and Sharma, R.C., 2001. Potato variety differences in nutrient deficiency symptoms and responses to NPK. *Better Crops International*, 15: 18-21.
- Zhao-Hui, LIU., Li-Hua, LI., Xiao-Lin JIANG., HARDTER R., Wen-Jun ZHANG., Yu-Lan ZHANG., and Dong-Feng ZHENG., 2008. Effect of N and K Fertilizers on Yield and Quality of Greenhouse Vegetable Crops. *Pedosphere*, 18(4): 496–502.

UNDER PEER REVIEW