

# Phenotypic Characterization of Indigenous Turkeys (*Meleagris gallopavo*) in the Ashanti, Ahafo, Bono and Bono East Regions of Ghana

## Abstract

A study was conducted to assess the effects of variety, region, and sex on body measurements and phenotypic correlations between various body measurements.

Three hundred (300) adult ~~indigenous~~ Indigenous Turkeys consisting of 195 males and 105 females, age of 6-7 months ~~old~~ from each of the four middle-belt regions were sampled. Data on qualitative and quantitative traits from 3 Turkeys consisting of two (2) males and one (1) female randomly selected per farmer ~~was taken (195 males and 105 females)~~. The Morphometric traits such as body weight, beak length, body length, shank length, thigh length, head length, and sternum length were measured and recorded in kilograms (kg) and centimeters (cm). The Qualitative characters (colour) were determined using colour chart.

White, Bronze, Black, Black White, and Buff were ~~the Turkey plumage identified~~ identified Turkey plumage. The overall mean body weights for the five colour varieties were: White ( $4.792 \pm 0.201$  kg), Black ( $4.744 \pm 0.112$  kg), Bronze ( $4.783 \pm 0.284$  kg), Black & white ( $4.547 \pm 0.113$  kg) and Buff ( $5.059 \pm 0.292$  kg). Turkeys from the Bono East region had much higher body weight than Turkeys from the Ashanti, Bono, and Ahafo regions. Cases of genotype-environment interaction were observed. The effect of Sex and Region was a highly significant ( $P < 0.05$ ) ~~source of variation for most of the~~ variation source for most traits. There were positive, moderate-to high correlations between the various body measurements. Body measurement was highly correlated ( $P < 0.05$ ) with body weight, ranging from 0.69 for head length to 0.96 for wing length. This study indicated that there are five (5) indigenous Turkey lines ~~and each showed distinct physical variations for both qualitative and quantitative traits which~~. Each showed distinct physical

**Commented [EM1]:** which traits were most affected by sex and region

[variations for qualitative and quantitative traits that](#) can be harnessed for future genotypic and molecular characterization.

**Keywords:** Turkey; [Region Ghana](#); Correlation; Characterization, Quantitative

## Introduction

Characterization of Animal Genetic Resources (AnGR) is of enormous importance to every nation since it provides employment, income, and food (Mogre, 2010). It includes chickens, turkeys, ducks, geese, quails, guinea fowls, and other domesticated birds (FAO, 2014). Ghana's upsurge in population has led to [an augmented urge increased demand](#) for meat (Osei *et al.*, 2012). This has necessitated the need for producers to [often frequently](#) import improved AnGR (Mogre, 2010). In most developing countries, the demand for improved breeds has led to indiscriminate crossing leading to dilution or loss of the adapted breeds (Serem, 2014). This strategy has led to undiscovered genes being lost forever. Genetic depletion of local animal variety has placed 20 percent of the world's breeds at risk of extermination (Osei *et al.*, 2012). This [feat-issue](#) is prevalent in most African countries, where there has been little effort to conserve the local poultry breed or lines (Manyelo *et al.*, 2020), which is also [the a](#) challenge among Ghanaian breeds and Turkey is no exception. To overcome the problem of breed annihilation and contain the loss of important undiscovered genes, conservation and sustainable development of Farm Animal Genetic Resources (FAnGR) relying on the several varieties that live well in the low external input agriculture typical of developing countries is suggested (Rexhaj *et al.*, 2018). Turkey is gaining much consideration in Ghana (Boschloo, 2019). Most of the nation's conventional livestock comes from the Ashanti, Brong Ahafo, and Upper West Region (Taylor, 2023). It is a source of protein for the rural

**Commented [EM2]:** The introduction mentions several studies and facts without fully explaining their relevance to the specific study on turkeys.

**Commented [EM3]:** Introduce the importance of turkeys in Ghana earlier in the introduction, and provide more detail on their current status and the challenges they face.

**Commented [EM4]:** kindly ensure that the most recent and relevant studies are cited, references from 2010 might be outdated, but newer studies could provide a more current perspective.

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

populace, employment and generate family income (MOH, 2010). The Turkeys in the commercialized sector have lost several abilities for survival in the wild; and can no longer exist without human maintenance. The decline in the production of Turkeys is normally due to the sensitivity of this animal to infectious diseases and the strain to cope with an unfavorable environment and also due to their low reproductive performance (Marchewka *et al.*, 2013). Thus, the indigenous Ghanaian Turkey is one of the AnGRs in Ghana that requires much attention and improvement towards increase in production. FAO (2014) reported that Turkeys in Ghana generally exhibited various feather colors including bronze, white/black, black, and red/buff as a whole or in combination. However, body weight, beak length, body length, shank length, thigh length, head length, sternum length, interactive effects, and correlation were not actualized in the available studies. Specifically, while this study was on phenotypic characterization, the available studies in Ghana were on perceptions and documentation surveys (FAO, 2014). Hence, it is of fundamental importance to estimate the frequency, investigate the effect of breed, colour variety, region, and sex and also determine the phenotypic correlations between selected morphometric traits of the indigenous Turkeys in Ghana.

The purpose of this study was therefore aimed at determining the phenotypic characterization of indigenous turkeys (*Meleagris gallopavo*) in the Ashanti, Ahafo, Bono, and Bono East Region of Ghana.

## Materials and methods

### *Location and duration of the experiment*

The survey was conducted from October 2020 to April 2021. The study took place in the Ashanti, Ahafo, Bono, and Bono East regions (Figure .1). These regions lie between latitudes 6.7470 N, 7.000 N, 7.3900 N, and 7.430 N, and longitudes 1.5208 W, 1.729 W, 1.8429 W and 1.9268 W respectively (GSS, 2019). The vegetation of these regions consists of deciduous, moist semi-deciduous forests and the soils are very fertile. The areas experience two seasons annually. The wet season is between April and November and the dry season is between December and April. The average temperatures are 23.9 °C with a minimum of 20.3 °C, a maximum of 37.8 °C, and an average rainfall of 1276 mm with Humidity at 72 % (WorldData, 2020).

### *Selection of the study area*

Formatted: Highlight

Commented [EM5]: could be clarified by explaining why these traits are important to study in turkeys.

Commented [EM6]: Please expand on the specific objectives, explaining what new insights this research aims to provide and why these insights are important.


The study areas were selected from the four (4) zones found on the Map of Ghana, namely: Ashanti, Ahafo, Bono, and Bono East Region. These administrative zones were chosen based on purposive sampling. A total of six (6) districts were selected from each of the four regions or study zones. Ashanti (Adansi Asokwa, Afigya Kwabre North, Ejura Sekyedumase, Ahafo Ano South West, Akrofuom District, and Adansi North), Ahafo (Asunafo North Municipal, Asunafo South, Asutifi North, Tano North, Asutifi South, and Tano South), Bono (Banda, Berekum West, Dorma Central, Jaman South, Sunyani, and Wenchi) and Bono East Region (Atebubu Amantim, Pru East, Kintampo South, Sene East, Techiman Municipal, Nkoranza) were purposely chosen from the four zones representing the Middle-belt of Ghana. The towns selected for the study were Asokwa, Boaman, Ejura, Adugyama, Akrofuom, Fomena, Goaso, Kukuom, Kenyasi, Duayaw nkwanta, Hwidiem, Bechem, Banda Ahinkro, Jinijini, Dorma Ahinkro, Drobo, Sunyani, Wenchi, Atebubu, Yeji, Kintampo, Kajaji, Tuobodom and Busunya. The snowballing technique (Coleman, 1958) was used in identifying the farms, thus a total of 154 farms were used to generate morphological and phenotypic characteristics of the indigenous Turkeys.

Visual appraisal of the external features of 300 indigenous Turkeys using a longitudinal design was done. Hence, morphologically distinct indigenous Turkeys were sampled using a random sampling technique to collect data on qualitative traits (plumage colour) and quantitative traits such as body weight, shank length, and sternum length following the standard descriptor (FAO, 1986).

#### ***Morpho-biometric data collection***

Three hundred (300) adult indigenous Turkeys of 6-7 months old from each of the four middle-belt regions were sampled. Data on qualitative and quantitative traits from 3 Turkeys consisting of two (2) males and one (1) female randomly selected per farmer was taken (195 males and 105 females). The following morphometric traits were measured using a 50kg capacity scale with a precision of 10g, a tape measure, and a colour chart. Measurements were recorded in kilograms (kg) and centimeters (cm). The qualitative characters were determined by direct observation of each Turkey with the help of a colour chart (Au-IBAR, 2015).

Picture 1 : Morphology of Indigenous Turkeys (*Meleagris gallopavo*)



The quantitative data measured include:

- A. **Beak length**: distance between the ends of the upper mandible and commissure of the down and upper mandibles
- B. **Body length (BL)**: distance between the tip of the upper mandible and the tail.
- C. **Wing Length** was taken from the shoulder joint to the extremity of terminal phalanx.
- D. **Thigh Length (TL)**: distance between the hock joint and the pelvic joint
- E. **Sternum length (SL)**: distance between both vertices of the sternum (processus carinae)
- F. **Shank length (SKL)**: distance between the calcaneus and the ankle and processus xiphoideus leaning the bird on its back
- G. **Head Length**: end of the neck to start of the beak.
- H. **Body weight (BWT)**: Birds were weighed using the 50kg of capacity with a precision of 10g and their weights were read and recorded

Qualitative Parameters included:

**Feather colours**: Various feather colours were identified visually and their frequency and detailed description were recorded with the aid of a colour chart.

### Data analysis

The effects of variety, region, and sex on body measurements. The data was subjected to least squares analysis of variance using the GLM procedure Type III of GenStat Discovery Edition 11 (2008) on the following fixed model

$$Y_{ijkl} = \mu + V_i + R_j + S_k + VR_{ij} + VS_{ik} + SR_{kj} + VRS_{ijk} + e_{ijkl} \dots \dots \dots (1)$$

$Y_{ijkl}$  = body weight, beak length, wing length, body length, thigh length, sternum length, shank length, head length.

$\mu$  = the overall mean

$V_i$  = the effect of the  $i^{\text{th}}$  variety of Turkey,  $i = 1, 2, 3$  and  $4$

$R_j$  = the effect of the  $j^{\text{th}}$  region,  $j = 1, 2, 3$  and  $4$

$S_k$  = the effect of the  $K^{\text{th}}$  sex of the Turkey,  $k = 1$  and  $2$

$VR_{ij}$  = is the interaction effect between  $i^{\text{th}}$  variety and the  $j^{\text{th}}$  region

$VS_{ik}$  = is the interaction effect between  $i^{th}$  variety and the  $K^{th}$  sex

$SR_{kj}$  = the interaction effect between  $K^{th}$  sex and the  $j^{th}$  region

$VRS_{ijk}$  = the interaction effect between  $i^{th}$  variety, the  $j^{th}$  region, and  $K^{th}$  sex

$e_{ijkl}$  = the random error term assumed normally and independently distributed,  $(0, \sigma^2 e)$

Differences among means of significant effects were separated by the probability of difference using the same software.

## Results and Discussion

### Qualitative Traits (Colour Frequency)

From a total sample of 300 adults-adult (6-7 months-old) Turkeys from the middle belt segment of Ghana, five colour varieties of Turkeys were identified. The colour varieties identified were white, black & white, black, bronze, and buff varieties (Table 1).

### Distribution of Turkey Plumage in the Ashanti, Ahafo, Bono, and Bono East Region of Ghana

**Table 1:** Percentages of the five Colour Varieties of Turkey in the Ashanti, Ahafo, Bono and Bono East Region of Ghana

Variety	Ashanti Region (n=84)	Bono Region (n=71)	Ahafo Region (n=79)	Bono East Region (n=66)	Total/Variety
---------	-----------------------------	--------------------------	---------------------------	-------------------------------	---------------

White	36.1	22.2	30.6	11.1	100
Black	29.3	25.8	25.0	19.8	100
Bronze	38.9	16.7	27.8	16.7	100
Black& white	22.1	24.8	27.4	25.7	100
Buff	29.4	11.8	17.7	41.2	100

The wide variation ~~of the in~~ plumage colour of ~~indigenous-Indigenous~~ Turkeys ~~in the various across different~~ regions indicates the existence of genetic variability (Marelli *et al.*, 2022). This suggests that the Ghanaian indigenous poultry resources ~~are still have~~ not ~~been highly heavily~~ diluted by exotic breeds (Cocou *et al.*, 2023). These colour frequencies are in disagreement with Naceur *et al.* (2014) who reported the highest frequency for white/black (29.8 %), followed by bronze (26.4 %), black (22.9 %), and red/buff (20.9 %). Turkeys are of different plumage colors: black, white/black, white, red/buff, and bronze, these results are in agreement with the results of Halbouche *et al.* (2010). These colour frequencies are also in disagreement with Savage and Zakrzewska (2006) who found a frequency of 30 % for bronze as well as black (26 %) followed by black and white (22 %). The wide variation of plumage colour of local Turkey ~~could be the result likely reflects the genetic diversity that has of the genetic variation~~ accumulated ~~through over~~ decades of domestication ~~of the species~~ (Amado, 2023)

**Commented [EM7]:** I think the most recent and relevant studies cited would be beneficial.

### Quantitative Traits (Body Measurement)

*Body weight and linear body measurements of Turkeys in the middle belt of Ghana*

**Table 2:** Effect of region on linear body measurement and body weight of turkeys in the middle belt of Ghana

Variable	Ashanti Region	Bono Region	Ahafo Region	Bono East Region	P-value	LSD
BWT (kg)	4.908±0.08 <sup>b</sup>	4.997±0.09 <sup>b</sup>	4.978±0.08 <sup>b</sup>	5.356±0.09 <sup>a</sup>	0.001	0.242
BKL (cm)	2.424±0.03	2.528±0.03	2.490±0.03	2.498±0.03	0.059	0.079

HL (cm)	6.393±0.20	6.408±0.23	6.278±0.22	6.303±0.24	0.968	0.616
BL (cm)	30.74±0.37 <sup>ab</sup>	31.62±0.40 <sup>a</sup>	29.55±0.38 <sup>b</sup>	30.46±0.41 <sup>b</sup>	0.002	1.080
THL (cm)	19.26±0.15 <sup>b</sup>	20.37±0.17 <sup>a</sup>	19.57±0.15 <sup>b</sup>	19.40±0.17 <sup>b</sup>	<0.001	0.431
WL (cm)	27.97±0.41 <sup>b</sup>	30.35±0.45 <sup>a</sup>	29.09±0.43 <sup>a</sup>	29.24±0.46 <sup>a</sup>	0.002	1.219
SHK (cm)	11.82±0.18 <sup>b</sup>	12.49±0.20 <sup>a</sup>	12.30±0.19 <sup>a</sup>	11.91±0.21 <sup>b</sup>	0.047	0.276
SL (cm)	12.61±0.25 <sup>b</sup>	13.88±0.26 <sup>a</sup>	13.26±0.26 <sup>a</sup>	12.35±0.29 <sup>b</sup>	<0.001	0.749

*P-Value = probability value, BWT= body weight, BKL= beak length, HL = head length, BL= body length, THL= thigh length. WL= Wing length, SHK= shank length and SL= sternum length*

*NB: Means between regions with different superscripts are significantly different ( $p \leq 0.05$ ).*

At maturity, ~~which is~~ typically around 18-24 weeks for commercial turkey breeds, there is a ~~substantial-significant~~ increase in body weight and muscle development, ~~crucial for meat production. By this stage, turkeys~~ ~~Turkeys have generally are raised for meat production, and at this stage, they have~~ reached their full adult size. The muscle ~~is well-developed~~, especially ~~in key morphometric traits such as~~ body weight, beak length, head length, body length, and thigh length. ~~Wing-wing~~ length, shank length, and sternum length ~~are well-developed to meet the demands of meat production~~ (Sogut *et al.*, 2016)

The location had a significant ( $P \leq 0.05$ ) influence on the body measurements of Turkeys in most of the traits. Body weight of Turkeys in the Ashanti, Bono, and Ahafo regions were similar ( $P \geq 0.05$ ) and these were lower ( $P \leq 0.05$ ) than values recorded in the Bono East region.

Beak length and Head length values were similar ( $P \geq 0.05$ ) across all the regions, ~~indicating no significant regional variation in these traits~~. For body length, Ahafo and Bono East recorded similar ( $P \geq 0.05$ ) values. However, these were ~~significantly~~ lower ( $P \leq 0.05$ ) than ~~those values~~ recorded in Bono and Ashanti regions. Thigh lengths for Turkeys in the Ashanti, Ahafo, and Bono East regions were similar ( $P \geq 0.05$ ) and these were lower ( $P \leq 0.05$ ) than values in the Bono region. Wing lengths for Turkeys in Bono East, Bono, and Ahafo ~~region-regions~~ were similar ( $P \geq 0.05$ ) and these were higher ( $P \leq 0.05$ ) than those in the Ashanti region. The shank ~~length-lengths~~ of



Turkeys in the Ashanti and Bono East regions were similar ( $P > 0.05$ ) and these were lower ( $P < 0.05$ ) than values in the Bono and Ahafo regions which also had similar values ( $P > 0.05$ ). The sternum lengths of Turkeys in the Bono and Ahafo regions were similar ( $P > 0.05$ ) and these values were higher ( $P < 0.05$ ) than those in the Ashanti and Bono East regions which also recorded similar values. Environmental factors play an important role in the variation of the size of birds. According to Naceur *et al.* (2014), the variation in the size of the bird is due to the collective effects of [environmental factors including](#) humidity, temperature, and altitude which play an important role in the availability of feed in each region, subsequently influencing the body weight and linear measurement of the birds.

**Table 3:** Effect of sex on for linear body measurements and body weight of Turkeys in the middle belt of Ghana

VARIABLE	SEX	MEAN±STANDARD ERROR	P-VALUE	LSD
BL (cm)	Male	32.53±0.15	0.001	0.506
	Female	26.94±0.21		
BKL (cm)	Male	2.60±0.013	0.512	0.044
	Female	2.25±0.018		
WL (cm)	Male	31.39±0.16	0.001	0.418
	Female	24.87±0.22		

THL (cm)	Male	20.41±0.07	0.006	0.218
	Female	18.20±0.09		
HL (cm)	Male	7.20±0.11	0.001	0.161
	Female	4.762±0.15		
SL (cm)	Male	14.38±0.11	0.001	0.357
	Female	10.50±0.15		
SKL (cm)	Male	13.18±0.07	0.001	0.217
	Female	10.17±0.09		
BWT (kg)	Male	5.521±0.03	0.003	0.101
	Female	4.182±0.04		

M± E S: Mean± Standard error.

*P-Value = probability value, BWT= body weight, BL= body length, WL= Wing length, HL = head length, SL= sternum length, BKL= beak length, SHK= shank length and THL= thigh length*

Sexual dimorphism was [evident](#) in favor of the male ( $P < 0.05$ ) ~~as expressed in across~~ all traits studied. The mean body weight of indigenous Ghanaian Turkeys found in the Middle belt was 5.52±0.03 and 4.18±0.04 kg for the males and females respectively. This trend was also similar to a report by Naceur *et al.*, (2014) who recorded weight for adult turkeys kept under the semi-intensive system of between 6.70 and 8.90 kg in males and between 2.90 and 3.14 kg in females. [Similarly, Ogah \(2011\) observed comparable trends in 20-week-old turkeys from Nigeria, with males and females weighing 3.38 kg and 2.65 kg, respectively.](#) ~~Ogah (2011) reported similar trends on 20-week-old turkeys from Nigeria (3.38 Kg and 2.65 Kg for male and female, respectively).~~ However, the ~~values-measurements for~~ [body length of the body](#), [beak length of the beak](#), [shank length of the shank](#), [thigh length of the thigh](#), and [sternum length of the sternum](#) of adult females were lower than those observed by Naceur *et al.*, (2014) [for turkeys from in Tunisia](#). ~~in Turkeys.~~ This dissimilarity in size of the male and female is associated with the effect of various hormones that ~~leads-lead~~ to a different growth rate (Naceur, 2014).

Formatted: Strikethrough

**Table 4: Interaction effect of fixed factors on morphometric traits**

Type Interaction effect	BL	BKL	WL	THL	HL	SL	SHK	BWT
<b>Region*Colour</b>	ns	ns	ns	ns	ns	ns	ns	ns
<b>Region*Sex</b>	*	*	*	*	*	*	*	*

<b>Colour*Sex</b>	*	*	*	*	*	*	*	*
<b>Region*Colour*Sex</b>	*	*	*	*	*	*	*	*

Ns= Not significant, \* = Significant

BWT= body weight, BL= body length, WL= Wing length, HL = head length, SL=sternum length, BKL= beak length, SHK= shank length and THL= thigh length.

#### Interaction effect of fixed factors on morphometric traits

Region and phenotype interaction effects on all the traits were not significant ( $P > 0.05$ ). All other interaction effects on all traits were significant ( $P < 0.05$ ) as shown in Table 4. The interaction effects between region and phenotype on all traits were not significant ( $P > 0.05$ ), while all other interaction effects were significant ( $P < 0.05$ ), as detailed in Table 4.

The significant effect of sex as well as and region on most measurements indicates that the measurement of the performance depends on the region in which where the  $\chi$  turkeys were assessed. This is a case of genotype-environment interaction (Erdem, 2021). This implies that selection for improvement of body weight and size should be done within regions provided the same genotypes exist (Erdem, 2021). Colour and sex had a significant ( $P < 0.05$ ) effect on all the variables. From a morphological point of view, it is important to consider the type of variety/color to use in most genetic improvement programmes and the design of the breeding objectives. This is because body measurements can be influenced by the sexual hormones. Similar effects of sexual hormones on linear body measurement and live weight have been reported in other livestock species, such as broilers (Olanrewaju *et al.*, 2019) and Turkeys (Nabtiti, 2017)

**Table 5: Phenotypic Correlation coefficients of body weight and body measurements of male (above diagonal) and female (below diagonal) Turkeys**

	BWT	BKL	WL	THL	SL	SHK	BL	HL
BWT		0.88***	0.96*	0.95***	0.85***	0.86***	0.91***	0.69***
BKL	0.35***		0.91*	0.74***	0.78***	0.82*	0.88***	0.69***

WL	0.46***	0.61***		0.87***	0.79***	0.83***	0.90*	0.70***
THL	0.88***	0.58***	0.87***		0.79***	0.76*	0.87***	0.57***
SL	0.78**	0.51***	0.58***	0.87***		0.71***	0.71***	0.73***
SHK	0.84***	0.32***	0.75***	0.85***	0.84***		0.96***	0.74***
BL	0.79**	0.67***	0.76***	0.44**	0.36*	0.52***		0.68***
HL	0.64***	0.32***	0.56**	0.69***	0.45***	0.49***	0.40***	

\*= $P \leq 0.05$ , \*\*= $P < 0.01$ , \*\*\*= $P < 0.001$ .  $R \leq 0.30$ : Low, 50; moderate,  $\geq 51$  strong

BWT= body weight, BL= body length, WL= Wing length, HL = head length, SL= sternum length, BKL= beak length, SHK= shank length and THL= thigh length.

The body measurement is highly correlated ( $P < 0.05$ ) with body weight, [with correlation coefficients](#) ranging from 0.69 for head length to 0.96 for wing length. Similarly, relationships between all the traits were positive and significant. The high and significant phenotypic correlations between body measurements and body weights in both sexes suggest high predictability between the traits in both male and female Turkeys. Ogah, (2011) recorded a similar trend between body weight and principal body measurements in Turkey, and thus indicated that selection for body weight may lead to an increase in other body measurements given that the majority of genes influencing the body weight and body measurements of Turkey are of common attainment. The implication here is that it can be helpful as a selection criterion. [This also indicates that there is a good relationship between weight and body parameters. The implication here is that body measurements can serve as effective selection criteria, indicating a strong relationship between weight and body parameters.](#)

## Conclusion

The findings of this research [on turkey characteristics across different regions](#) highlight [significant the regional distinctions in turkey morphometric traits](#) notably in body weight, body length, thigh length, wing length, shank length, beak length, and head length [with turkeys Turkeys](#) from Bono East region consequently [having have](#) heavier body weights (carcass yield) compared to the other regions and this may inform strategies for optimizing turkey production [in specific geographic](#)

areas. ~~Also~~ Additionally, selection for increased body weight (carcass yield) may concurrently result in an enhancement of other body measurements. ~~Also, the p~~ Pronounced divergence (sexual dimorphism) in physical attributes highlights a notable disparity between sexes within the studied population and this provides a fundamental basis for targeted intervention and advancement. The strong correlation between various body measurements and body weight in both ~~males and females sexes~~ underscores the intricate association within the studied characteristics and these findings contribute valuable insights into the interplay of body measurement and weight, potentially informing future research on the subject. [The strong correlations between various body measurements and body weight in both sexes provide valuable insights into the complex relationships among these traits, potentially guiding future research and breeding programs.] – as an alternative sentence

Further research ~~works~~ should ~~be steered~~ focus on ~~the a comprehensive~~ characterization of the indigenous Turkeys in ~~the same~~ these regions including Genetic, molecular, and immunological characterization.

#### Ethics approval and consent to participate

All authors hereby declare that “Principle of laboratory animal care” (NIH publication No.85-23, revised 1985) ~~were was~~ followed as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee (state the approval document number)

**Commented [EM8]:** Ethical approval from an animal care and use committee may be necessary

#### Consent for publication

All authors declared that written informed consent was obtained from the approved parties for publication if this article and accompanying images

#### Reference

- [1] Adedibu I.I and Ayorinde, K.L. (2011). Sexual Dimorphism in Predicting Body weight of two Broiler Strains.
- [2] Amado M. C, María E. C, Antonio H. B, Juan V. D, Vincenzo L, Amparo (.year.) MGenetic diversity in 10 populations of domestic Turkeys by using microsatellites markers, Poultry Science, Volume 102, Issue 1.

**Commented [EM9]:** Year ?

- [3] Boschloo R. (2019) Analysis poultry sector Ghana: An update on the opportunities and Challenges. Netherlands Enterprise Agency.
- [4] Brown, M. M., Alenyorege, B., Teye, G. A., & Roessler, R. (2017). Phenotypic diversity, major genes and production potential of local chickens and guinea fowl in Tamale, northern Ghana. *Asian-Australasian journal of animal sciences*, 30(10), 1372.
- [5] Cocou C.K, Yaah A.E.K., Oumbortime N, Frederich M.H, Kokou T. and Oyegunle E.O. (2023). The productivity and resilience of the indigenous chicken in the tropical environments: improvement and future perspectives, *Journal of Applied Animal Research*, 51:1,456-469
- [6] Coleman J. S. (1958-1959). Rational Analysis: The study of social organization with survey methods. *Human Organization*.
- [7] Erdem H.S. (2021). Genotype-environment interaction in layer chickens in the growing stage: comparison of three genotype at two different feeding levels with or without red mite (*Dermanyssus gallinae*) infestation. *Arch Anim Breed*
- [8] FAO (2012). Phenotypic characterization of animal genetic resources. FAO Animal Production and Health Guidelines. No. 11. Rom
- [9] FAO. (2014). Poultry Sector Ghana. FAO Animal Production and Health livestock Country Reviews. No. 6. Rome
- [10] FAO. (2014). Chicken genetic resources used in smallholder production systems and opportunities for their development, by P.Sorensen. FAO smallholder poultry Production Paper No. 5 Rome
- [11] Halbouche M, Mouats A, Belhadi A, Homrani A, Didi M (2010). Ponte, Reproduction et

**Commented [EM10]:** Please be careful in writing the reference (kindly be consistent)

Développement Sexuel Chez la dinde Domestique (*MeleagrisGallopavo*) en Algérie.  
Influence du Phénotype et du Niveau Protéique des Aliments. European Journal of  
Scientific Research ISSN 1450-216X Vol.47 No.3, pp.473-483

- [12] Marelli SP, Zaniboni L, Strillaci M.G, Madeddu M, Cerolini S. (2022) Morphological Characterization of two light Italian Turkey Breeds. *Animals* (Basel).
- [13] Marchewka, J, Watanabe T.T.N, Ferrante V, Estevez I. (2013). Review of the social and environmental factors affecting the behavior and welfare of Turkeys (*Meleagris gallopavo*). *Poultry Science* 92:1467–1473.
- [14] Mbelayim, H. S.N.H (2015). Phenotypic and morphological characterization of Indigenous Ghanaian rabbit (*oryctolagus cuniculus*) resources in northern Ghana
- [15] Ministry of Food and Animal, (2000). Constraints and Recommendations on Improving the Competitiveness of the Ghanaian Poultry Industry; prepared by a committee appointed by MOFA with membership Directorate, The veterinary Services Directorate, the Animal Research Institute (CSIR) and the Livestock Planning and Information Unit, Accra
- [16] Mogre, J. W. S. (2010). Phenotypic and Morphological Characterization of indigenous Guinea fowl resources in northern Ghana. An M. Sc. dissertation submitted to the Department of Animal Science University of Ghana. Pp: 4-60
- [17] Mogesse, H (2007). Phenotypic and genetic characterization of indigenous chicken populations in Northwest Ethiopia. Thesis submitted to the Faculty of Natural and Agricultural Sciences Department of Animal, Wildlife and Grassland Sciences University of the Free State, Bloemfontein, South Africa. In partial fulfilment of the requirements for the degree Philosophy Doctor (PhD)
- [18] MOH (2010). Dietary and Physical Activity Guidelines for Ghana. Ministry of Health, Ghana.

**Commented [EM11]:** What does MOH stand for, write it in full

**Formatted:** Highlight

- [19] Nabtiti A.E (2017). *Effect of stocking density, Breed, Age of Bird and their interaction on Turkeys Weight Gain, Feed Consumption and FCR*. International Journal of agricultural Sciences and Veterinary Medicine.
- [20] Naceur, M, Djebbi. A, Haddad. I, Chriki. A (2014) Phenotypic Characterization of the indigenous Turkey in the North West Regions of Tunisia
- [21] Osei, D.Y., Apori, S.O. & Osafo, E.L.K. (2012). Rabbit Production in Selected Urban Areas of Middle belt Ghana: Status and Implications for Policy and Research. *Journal of Animal Production*, 14(2):131 -140
- [22] Ogah, D. M. (2011). Assessing size and conformation of the body of Nigerian indigenous Turkey.
- [23] Olanrewaju H.A., Miller W.W., Maslin W.R., Collier S.D., Purswell J.L., Branton, S.L. (2019) Interactive effects of light-sources, photoperiod, and strains on growth performance, carcass characteristics, and health indices of broilers grown to heavy weights, *Poultry Science*, Volume 98, Issue 12,
- [24] Rexhaj, N., Papa, L., Hoda, A., & Kume, K. (2018). The State of Local Cattle of Busha Type in Albania and Kosovo. Their Identification and Morphometric Description. *ANGLISTICUM. Journal of the Association-Institute for English Language and American Studies*, 7(1), 78-86.
- [25] Savage T.F and Zakrzewska E.I. (2006). Single Gene Traits of the Turkey. <<http://oregonstate.edu/>
- [26] Serem, J. K. (2014). Characterization of Rabbit Production Systems. In Central, Coastal, Eastern and Rift Valley Regions of Kenya (Doctoral dissertation, University of Nairobi).



- [27] Sogut, B & Çelik, Ş, Ayasan, T and İnci, H (2016). Analyzing Growth Curves of Turkeys Reared in Different Breeding Systems (Intensive and Free-Range) with some Nonlinear Models. *Revista Brasileira de Ciência Avícola*. 18. 619-628. 10.1590/1806-9061-2016-0263.
- [28] Taylor J. (2023). Ghana Livestock Voluntary. United States Department of Agriculture, Foreign Agricultural Service.
- [29] Manyelo T. G, Selaledi L, Hassan Z. M and Mabelebele M (2020). Local Chicken Breeds of Africa: Their Description, Uses and Conservation Methods. *Animals (Basels)*.
- [30] World Data, (2020). The Climate in Ghana. Available at [www.worlddata.info/Africa/ghana/climate.php](http://www.worlddata.info/Africa/ghana/climate.php). Accessed 18/10/21