

Original Research Article

Germination and Seedling Performance of Watermelon as affected by Seed Priming

ABSTRACT

An experiment was conducted to study the germination and seedling performance of three watermelon cultivars viz. Asian 2, Pakorea F1 and Black Red with six seed priming treatments such as no priming, hydropriming with distilled water, halopriming with 1.5N NaCl and 3% KNO₃, Osmopriming with 3% PEG (Polyethylene Glycol) and oxidative priming with 0.1N HCl solution. The seeds were primed overnight with the aforesaid liquids and were allowed to grow both in petridishes in dark condition within laboratory and in pot soil under natural condition in an open grill house. All three watermelon cultivars showed significant difference due to seed priming with aforesaid primers. Germination Index (GI) was found higher in Asian 2 and Black Red cultivars due to HCl priming and in Pakorea F1 due to NaCl priming, while priming with PEG and KNO₃ inhibited the GI of respective cultivars. Seedling Vigor Index (SVI) was found higher in Pakorea F1 with HCl priming whereas lower in Asian 2 and Black Red cultivars with PEG priming. Both in dark (Lab) and light (open field) conditions, total seedling dry matter was found higher in Asian 2 and Black Red cultivars due to HCl priming and lower in Pakorea F1 due to PEG priming. Results showed that low concentrated (0.1N) HCl solution can be used as priming medium to enhance germination and seedling growth of watermelon seeds.

Key words: HCl, Hydropriming, Osmotica, Polyethylene Glycol, Primer, Vigour index

1. INTRODUCTION

Watermelon (*Citrullus lanatus* L) is a scrambling and trailing vine-like flowering plant species of the Cucurbitaceae family. It is a highly cultivated fruit and grown from tropical to temperate regions worldwide. Edible watermelon is a large fruit, which is a Berry with a hard rind and no internal divisions, and is botanically called a Pepo. The fruit can be eaten raw or pickled, and the rind is edible after cooking. It may also be consumed as juice or as an ingredient in mixed beverages. Watermelon is grown in 122 countries and around 1,200 cultivars are grown on all cultivated continents (Tibor, 1993; Adeoye et al., 2011). China currently produces the largest quantity, followed by Iran, Turkey and Brazil (FAO, 2017). In Bangladesh, annually watermelons are cultivated in near about 15,740 hectares of land and 2,74,000 metric tons of watermelons are produced with an average production of 23.33 Mt/ha (Biswas, 2020).

Watermelon is mainly cultivated in Patuakhali, Bhola, Borguna, Noakhali, Pirojpur, Bagerhat, Sunamganj, Pabna, Natore, Panchagarh, Gopalganj, Jhenaidah, Chuadanga, Feni and Satkhira districts of Bangladesh (Biswas, 2020). In the Southern part of Bangladesh, it is

Commented [Reviewer1]: Please add the keywords such as seed priming and seed germination

Commented [Reviewer2]: Is it a suitable keyword for this article?
I am not sure if it can be a keyword in this article.

Commented [Reviewer3]: Please focus more on seed priming and review more related articles.

extensively cultivated and it is one of the major crops grown there and watermelons of this region have a high demand in the market due to its high quality and bigger size. Both hybrid and open pollinated cultivars are cultivated by the Bangladeshi growers but the demand of hybrid cultivars is much higher than the open pollinated cultivars.

Watermelon is cultivated in the various constraints in Bangladesh where seed germination and early crop establishment is a major barrier such as hard soil crust, drought, excessive rainfall, water logging, nutrient deficiency and so on. However, no study on the effects of different priming agents or osmotica on the performance of watermelon seed germination and seedling growth has been reported in Bangladesh.

Priming is basically a physiological process in which the seeds are presoaked before planting which, by itself, allows partial imbibition though preventing the germination (Nascimento and Aragão, 2004). During the priming, several processes including storage, material handling, activation and synthesis of a number of enzymes and nucleic acids, repair and build up, ATP synthesis, and the cytoplasmic membrane repair in treated seeds will all start to develop (Hosseini and Koocheki, 2007). Priming generally induces faster and more uniform seed germination especially in adverse physical conditions (Nascimento, 2003). It is nowadays being extensively used to improve seed germination and seedling emergence in a wide range of crop species (Hosseini and Koocheki, 2007). It is reported to practice in wheat, cabbage, pepper, muskmelon, mungbean etc. crop for better germination and early establishment of growing seedling in home and abroad (Aloui, 2014; Awal et al., 2024). Priming with some osmoticas was reported to produce uniform and healthy seedling of watermelon (Sivritepe *et al.*, 2003). So far, no study is conducted on seed priming in watermelon in Bangladesh although initial establishment of plantlets of this crop is a tuff job at farmer's level. Therefore, the aim of the study is to find out the effects of priming with some chemical agents on seed germination and early seedling growth of watermelon.

Commented [Reviewer4]: Please write the importance of this study.

2. MATERIALS AND METHODS

The study was conducted during the period from May 2019 to September 2019 in the Plant Ecology Laboratory of the Department of Crop Botany, Bangladesh Agricultural University, Mymensingh, and in a grill house in open natural condition outside the Department (24°75' N latitude and 90°50' E longitude). The experiment comprised two factors. Factor A: six seed priming treatments such as (i) no priming, (ii) hydropriming with distilled water, (iii) halopriming with 1.5N NaCl, (iv) halopriming with 3% KNO₃, (v) Osmopriming with 3% PEG (Polyethylene Glycol) and (vi) oxidative priming with 0.1N HCl; Factor B: three watermelon cultivars viz. Asian 2, Pakorea F1, and Black Red. The experiment was laid out in a Factorial Randomized Block Design with four replications.

The watermelon cultivar Asian 2 is a F1 hybrid imported from Bangkok, Thailand. Pakorea F1 is a hybrid Korean cultivar which is marketed in Bangladesh by DAEIL Seed Co. Ltd. Black Red is originated in Thailand and marketed by SEEDLINE Co. Ltd. Seeds of the aforesaid watermelon cultivars were collected Dhaka and Mymensingh seed markets. Before using for investigation, the seeds were kept in chilled condition of -20°C in a refrigerator.

The priming was done by immersing seeds of each cultivar in 100 mL of respected priming solutions in 500 mL glass beakers, covered with aluminum foil and kept overnight (for 12h) at 20°C in aerated dark chamber. After 12 hours, primed seeds were rinsed with distilled deionized water for two minutes with an aim to wash-off the priming solutions from the

surface of the seeds and then dried in room condition to the original moisture level and left overnight at room temperature to their original weight.

The clean and dried seeds were placed in the petri-dishes as per treatment design. The petri-dishes with 9 cm diameter and Whatman No. 1 filter paper were used for seed germination. The petri-dishes were kept in a dark condition in the Laboratory. The seeds in the petri-dishes were observed daily and the number of seeds passed for germination was recorded. After completion of germination, the lengths and dry weight of plumule and radicle were recorded. The dry weight was determined by drying the seedlings in an oven at $80 \pm 2^\circ\text{C}$ till attained a constant weight.

Besides the seeds placed in the petri-dishes, the primed seeds were also sown in the pot soil in the grill house (open or natural condition) that is situated outside the Department. In each pot, 25 seeds from each cultivar were sown at a depth of 3 cm. The pots were saturated in water by surface irrigation. During plantlet growth, pots were irrigated daily by spraying with water until water would drain from the bottom of the pot. After complete emergence, lengths and dry weight of plumule and radicle were recorded.

Germination Index (GI) was calculated as the number of seed germination observed or counted that day is divided by the number of days after seed placement (Ranal *et al.*, 2009):

$$GI = \sum Gt/Dt \quad \text{Eq [1]}$$

where, Gt = is the number of germinated seeds on day t and Dt = is the time corresponding to Gt in days.

Seedling Vigor Index (SVI) was calculated following Abdul-Baki and Anderson (1973):

$$\text{Seedling Vigor Index (SVI)} = [\text{Mean root length } (L_r) + \text{Mean shoot length } (L_s)] \times \text{Percentage of seed germination } (GP) \quad \text{Eq [2]}$$

where, GP is the seed germination percentage (%), ratio of the number of germinated seeds to initial fifteen test seeds; L_r is the root length (L); L_s is the shoot length (L).

The collected data on different parameters were statistically analyzed to obtain the level of significance using MSTAT-C Package Programme. The mean differences were compared with Duncan's Multiple Range Test.

3. RESULT AND DISCUSSION

3.1 Germination Index (GI)

Germination Index (GI) showed significant differences for the priming treatments in all watermelon cultivars (Fig. 1). In Asian 2, GI was maximum in HCl and minimum in control treatment, while in Pakorea F1, GI was higher in NaCl treatment and lower in KNO_3 treatment. The GI was higher in HCl and lower in PEG in Black Red cultivar.

Commented [Reviewer5]: You need to explain different things here:

- 1.The concentration of priming treatments.
- 2.How did you prepare the priming solutions?
- 3.For PEG (Polyethylene Glycol), did you use PEG6000 or PEG8000?
- 4.How did you calculate the concentration?
- 5.How did you decide the duration of seed priming, I think 12 hours is not enough for priming the watermelon seeds. Could you provide some references showing that 12 hours is efficient. I think the minimum should be 24 to 48 hours.

Commented [Reviewer6]: Please improve this part. Its written very poor.

Commented [Reviewer7R6]: You need to discuss your results with several other reports in the discussion part.




Fig. 1: Effect of seed priming on Germination Index (GI) in three cultivars of watermelon in Lab condition.

Commented [Reviewer8]: Please add the significance using Tukey's on the bars.

An improvement in the germination process with seed pre-priming agents has been commonly reported in several crops. Ebrahimi *et al.* (2014) reported that the osmopriming may enhance the seed germination and growth rate of tomato cultivars under salt stress conditions. Batista *et al.* (2015) reported that both hydropriming and osmopriming improved

germination and seed germination rate index of pepper seeds compared to unprimed seeds. Matias *et al.* (2018) indicated that hydropriming improved tolerance of sunflower plants under saline environments. According to Kubala *et al.* (2005), the priming can improve germination rate through metabolic activation involving the synthesis of proteins, nucleic acids, and enzymes, and increasing water uptake, respiratory activity, and reserve mobilization.

Germination rate index (GRI) of melon seeds was reduced with the increase of the salt stress level; however, this negative effect can be fully reversed with the hydropriming and osmopriming under mild salt stress. Under severe salt stress, the GRI ranged from 0.99 to 3.10 seed day⁻¹, and was significantly greater when the seeds were primed with water, followed by KNO₃ priming, and lower for unprimed seeds (Oliveira *et al.*, 2019).

3.2 Seedling Vigor Index (SVI)

Seedling Vigor Index (SVI) showed significant differences among the priming treatments in Asian 2, Pakorea F1 and Black Red cultivars. The results revealed that, SVI was maximum in Pakorea F1 with HCl treatment whereas minimum in Black Red cultivar with PEG treatment (Fig. 2). HCl treatment also influenced SVI in Black Red but not vary significantly in Asian 2. However, Asian 2 showed the maximum SVI with HCl treatment and minimum in PEG. In both Pakorea F1 and Black Red cultivars, maximum and minimum SVIs were observed in HCl and PEG, respectively.




Fig. 2: Effect of seed priming on Seedling Vigor Index (SVI) in three watermelon cultivars in Lab condition.

Commented [Reviewer9]: For unprimed treatment, you can control and explain it in the materials and methods.

Seedling vigor is a measure of the extent of damage that accumulates as viability declines, and the damage accumulates in seeds until the seeds are unable to germinate and eventually die (Kaya *et al.*, 2006; Farhoudi *et al.*, 2007; Marcos-Filho, 2015). SVI is used as a

phytotoxicity index to evaluate the effect of heavy metal, chemical etc. on seedling growth (Srivastava and Thakur, 2006; Kabir *et al.*, 2008). Salehzade *et al.* (2009) showed that osmopriming improved germination and seedling vigor than that unpriming in wheat (*Triticum aestivum* L.). Mamun *et al.* (2018) found that priming agent exerted significant effect ($p < 0.01$) on seed germination rate, germination pattern and seedling vigor of winter rice. In general, seed priming showed a positive effect on seed germination and seedling vigor of winter rice. NaCl at any concentration reduced emergence rate, while CaCl_2 priming enhanced germination rate the most. Priming with CaCl_2 , KCl and PEG found the best for germination index, while CaCl_2 , KCl and CuSO_4 resulted in the most vigorous seedlings (Zheng *et al.*, 2016). Seed priming with KNO_3 can cause a significant increase in seedling vigor of the wheat crop as compared to hydro-priming or dry broadcasting (Basra *et al.*, 2003).

3.3 Seedling dry weight

Seed priming showed significant influence on the total dry weight of watermelon seedling (Fig. 3). There was significant variation among the priming agents. In all three watermelon cultivars, total dry weight was found higher due to HCl priming and lower due to PEG priming both in dark condition under Laboratory and natural open pot sowing conditions.

Oliveira and Steiner (2017) reported that inhibiting action of salt stress on the initial growth of cucumber seedlings was increased with the rise of salinity levels, and the exposure of seeds to severe salt stress resulted in the decrease of total dry matter. Acosta-Motos *et al.* (2017) showed that the total dry matter production severely was severely inhibited by salinity; however, this effect was fully reversed with the use of seed priming with water or KNO_3 . This showed that seeds primed with water or KNO_3 may adjust the salinity stress.

Commented [Reviewer10]: Please check for spelling.

Commented [Reviewer11]: Please improve this part. You can cite the bellow articles in your discussion. Please find more articles related to your work and compare them with your results.

1. <https://www.doi.org/10.3390/stresses4020012>
2. <https://www.doi.org/10.3390/plants12112187>




Fig. 3: Effect of seed priming on total dry weight of watermelon in Lab (upper) and pot sowing (below) condition.

Commented [Reviewer12]: Does it mean root+shoot?

4. CONCLUSION

It may be concluded that seed priming with HCl showed better and priming with PEG showed poor germination, seedling vigour and dry matter production in all three watermelon cultivars studied both in dark condition under laboratory and pot sown under natural condition. So, low concentrated (0.1N) HCl solution can be used as best seed primer to enhance germination and seedling growth of watermelon seeds.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

~~Authors hereby~~ Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTEREST

Authors have declared that no competing interests exist.

REFERENCES

- Abdul-Baki AA, Anderson JD. Vigor determination in soybean seed by multiple criteria. Crop Science. 1973;13:630-633.
- Acosta-Motos JR, Ortuño MF, Bernal-Vicente A, Diaz-Vivancos P, Sanchez-Blanco MJ, Hernandez JA. Plant responses to salt stress: adaptive mechanisms. Agronomy. 2007; 18.
- Adeoye I, Olajide-Taiwo F, Adebisi-Adelani O, Usman J, Badmus M (2011). Economic analysis of watermelon based production system in Oyo State, Nigeria. ARPN Journal of Agricultural and Biological Science. 2011; 6:53-59.
- Aloui H, Souguir M, Latique S, Hannachi C. Germination and growth control and primed seeds of pepper as affected by salt stress. Cercetari Agronomice în Moldova. 2014;47: 83-95.
- Awal MA, Hossain MA, Iqbal MA, Soufan W, Erman M, Ammar H, Elsabagh A. Zinc biofortification of mungbean (*Vigna radiata* L.) cultivars through zinc chemo-priming. Pakistan Journal of Botany, 2024;56(5); [http://dx.doi.org/10.30848/PJB2024-5\(21\)](http://dx.doi.org/10.30848/PJB2024-5(21)).
- Basra S, Pannu I, Afzal I. Evaluation of seedling vigor of hydro and matriprimed wheat (*Triticum aestivum* L.) seeds. International Journal of Agricultural Biology. 2003;5:121-123.
- Batista TB, Binotti FF, Cardoso ED, Bardivieso EM, Costa E. Aspectos fisiológicos e qualidade de mudas da pimenteira em resposta ao vigor e condicionamento das sementes. Bragantia. 2015;74:367-373.
- Biswas SC. New improved variety development of rice & non rice crops of Bangladesh along with the generation of sustainable technologies of vegetable production. Journal of Crop Botany. 2020;15:102-115.
- Ebrahimi R, Ahmadizadeh M, Rahbarian P. Enhancing stand establishment of tomato cultivars under salt stress condition. South Western Journal of Horticulture, Biology and Environment. 2014;5:19-42.
- FAO. FAOSTAT. Food and Agriculture Organization of the United Nations, Rome, Italy, 2017. <http://www.fao.org/faostat/en/#data> (accessed 18 January 2019).

Commented [Reviewer13]: Please check the journal instruction on how to write the references.

- Farhoudi R, Sharifzadeh F, Poustini K, Makkizadeh MT, Kochakpor M. Effects of NaCl priming on salt tolerance in canola (*Brassica napus*) seedlings grown under saline conditions. *Seed Science and Technology*. 2007;35:754-759.
- Hosseini A, Koocheki A. The effect of different priming treatments on germination percent and mean germination time of four varieties of sugar beet. *Journal of Agronomy Research*. 2007;5:69-76.
- Kabir M, Iqbal MZ, Shafiq M, Farooqi Z. Reduction in germination and seedling growth of *Thespesia populnea* L., caused by lead and cadmium treatments. *Pakistan Journal of Botany*. 2008;40:2419-2426.
- Kaya MD, Okcu G, Atak M, Ckh Y, Kolsarc O. Seed treatments to overcome salt and drought stress during germination in sunflower (*Helianthus annuus* L.). *European Journal of Agronomy*. 2006;24:291-295
- Kubala S, Garneczarska M, Wojtyła Ł, Clippe A, Kosmala A, Żmieńko A, Lutts S, Quinet M. Deciphering priming-induced improvement of rapeseed (*Brassica napus* L.) germination through an integrated transcriptomic and proteomic approach. *Plant Science*. 2015;231:94-113.
- Mamun AA, Naher UA, Ali MY. Effect of seed priming on seed germination and seedling growth of modern rice varieties. *The Agriculturists*. 2018;16:34-43.
- Marcos-Filho J. *Fisiologia de sementes de plantas cultivadas*. Londrina: Abrates, 2015, 660p.
- Matias JR, Torres SB, Leal CC, Leite MD, Carvalho S. Hydropriming as inducer of salinity tolerance in sunflower seeds. *Revista Brasileira de Engenharia Agrícola e Ambiental*. 2018;22:255-260.
- Nascimento WM, Aragão FAS. Muskmelon seed priming in relation to seed vigor. *Scientia Agrícola*. 2004; 61:114-117.
- Nascimento WM. Muskmelon seed germination and seedling development in response to seed priming. *Scientia Agrícola*. 2007; 60:71-75.
- Oliveira CE, Steiner F. Potassium nitrate priming to mitigate the salt stress on cucumber seedlings. *Scientia Agraria Paranaensis*. 2017; 16:454-562.
- Oliveira CEDS, Steiner F, Zuffo AM, Zoz T, Alves CZ, De Aguiar VCB. Seed priming improves the germination and growth rate of melon seedlings under saline stress. *Ciência Rural*. 2019; 49:1-11.
- Ranal MA, Santana DG, Ferreira WR, Mendes-Rodrigues C. Calculating germination measurements and organizing spreadsheets. *Brazilian Journal of Botany*. 2009; 32: 849-855.
- Salehzade H, Izadkhah Shishvan M, Ghiyasi M, Forouzin F, Abbasi Siyahjani A. Effect of seed priming on germination and seedling growth of wheat (*Triticum aestivum* L.). *Research Journal of Biological Sciences*. 2009; 4:629-631.

- Sivritepe N, Sivritepe HO, Eris A. The effects of NaCl priming on salt tolerance in melon seedlings grown under saline conditions. *Scientia Horticulturae*. 2003; 97:229-223.
- Srivastava S, Thakur IS. Evaluation of bioremediation and detoxification potentiality of *Aspergillus niger* for removal of hexavalent chromium in soil microcosm. *Soil Biology and Biochemistry*. 2006; 38:1904-1911.
- Tibor F. Watermelon: *Citrullus lanatus* (Thunb.) Matsum. & Nakai. "Genetic Improvement of Vegetable Crops", 1993, pp. 295-311. <https://doi.org/10.1016/B978-0-08-778040-826-2.50025-4>.
- Zheng M, Hussain S, Tao Y, Jiang Q, Peng S, Huang J, Cui K, Nie L. Seed priming in dry direct-seeded rice: consequences for emergence, seedling growth and associated metabolic events under drought stress. *Plant Growth Regulation*. 2016; 78:167-178.