

# FACTORS ANALYSIS FOR DEPRESSION AND SUICIDAL IDEATION AMONG UNIVERSITY STUDENTS USING APRIORI ASSOCIATION ALGORITHM

## Abstract.

Depression stands as a prominent and prevalent mental health issue, representing a significant global public health concern. Its emergence can be attributed to diverse factors. Suicide stands as a prominent global cause of death, eliciting concern on a widespread scale. This study was to find the factors for depression among university students in Bangladesh. There are so many factors such as Loneliness, Hopelessness, Helplessness, Relationship Issues, Grade problems, Academic Pressure, Parental problems, Money problems, Social Comparison, Social Media Influence, Family Expectations, Lack of Sleep, Uncertain Future, Health Issues, Bullying, Substance Abuse and Unemployment etc. These factors vary among male and female students. Apriori association algorithm were used to calculate support, confidence and lift of factors sets. Frequent factors sets and relationship were found from the work using association algorithm. The research participants are Bangladeshi university students, Google Form were used to collect the data. Data collection carried out by online questionnaire. The findings from data analysis indicated that academic pressure, uncertain future, hopelessness, family expectation, loneliness, financial crisis and unemployment are the key factors. The prevention of suicides is achievable. Hence, identifying depression and forecasting the potential for suicide risk serves as a means to prevent instances of self-harm within the university student population.

**Keywords:** Suicidal Ideation, Suicidal Tendency, Depression, Factors, Apriori, University Students.

## 1 Introduction

Depression is one of the leading and common mental health problems. It is the most alarming global public health issue. It may arise from various factors, including a lack of understanding about the condition. Depression is marked by enduring feelings of sadness, hopelessness, diminished self-worth, heightened anxiety, sleep disturbances, feelings of guilt, tendencies toward self-harm, and ideation of suicide [1-3]. Depression not only affects one's ability to live normally, but it also is a burden to society in general [4,5].

Early detection of depression is an efficient solution. It would help student to take proper steps against depression. Suicide means actually killing themselves. Suicidal behaviors are frequently accompanied by hopelessness, depression, or self-destructive behavior. Suicidal behaviors are individuals experiencing repeated thoughts of killing themselves life (suicidal ideation), planning to kill themselves (suicide plan), and actual efforts to kill themselves (suicide attempt) [6,7].

Suicide is a leading cause of death worldwide. It is a concerning issue nowadays for all of us. The suicidal incident is a curse for the family and society. The impact of suicide incidents on family and society is inevitable. In most cases, suicide ideation is the result of depression [8,9].

Recently, Bangladeshi university students have been considered to be highly suicide-prone. A large number of students committed suicide in Bangladesh. All of them were meritorious and talented. It is a major concern for all of us that Bangladeshi university students are making such a dangerous decision to commit suicide.

Suicides are preventable. So, detecting depression and predicting suicide risk is a way to stop suicidal incidents among university students [10].

This research employs Apriori association algorithm to identify and analysis the factors for depression and suicidal tendency among university students.

In this work, data analysis and association used to detect depression and improve suicide risk predictions among Bangladeshi university students to save lives and make precautions.

### **1.1 Motivation**

The value of a life is invaluable. Every person has a right and wants to live a peaceful life in this beautiful world. University students are the most meritorious person in any country. They are the assets of a nation. A nation will not progress without the contributions of university students. They are proud of their own family. So, the value of the life of students is priceless.

Nowadays, a large number of students are suffering from depression. There are many factors behind this problem like a failure, family, economic instability, high expectation, etc. Depression destroys the student's career, happiness, and dream. It is a big problem for a person, family, and nation.

It's widespread and daily news of suicide incidents by Bangladeshi university students. In most cases, suicide ideation is the result of depression. We lost a meritorious life in a suicide incident, a family lost their beloved member, and deprived a nation of the services.

Day by day, suicidal cases are increasing among university students. It's high time to be aware and concerned to stop it.

We can save many lives by stopping suicide cases. If we can predict the suicidal incident, it's very much possible to stop it by precaution. As depression is a significant reason for suicide, we can detect depression and its level among students by analyzing the related factors to make it easier for suicide prediction.

This work is significant and well-timed for Bangladeshi university students to discover the factors for depression, detect depression among them and predict suicide risk. This work will help save lives and take precautions against students' depression and suicide ideation.

### **1.2 Objectives**

Depression and suicidal ideation seize our young generation. University students are noticeably suffered from deadly depression and later on commit suicide. We hope our young and future generations will live peaceful and active life.

The objectives of this study are to stop suicidal tendency among Bangladeshi university students by identifying the factors of depression, detecting the level of depression and predicting suicide risk. This work will predict suicide risks that will help to take precautions for victim students.

This work will be effective and helpful:

- i. To predict suicide risk among university students in Bangladesh.
- ii. To detect depression among students.
- iii. To measure depression among university student.
- iv. To identify the factors which are responsible for suffering from depression.
- v. To list primary reasons and factors for depression.
- vi. To prevent suicide incidents by predicting suicide risk.
- vii. To save the valuable lives of meritorious university students.
- viii. To take care of the mental health of students.
- ix. To create opportunities to arrange motivation sessions or counseling for proper and needed students.
- x. To stop the premature death of students.
- xi. To save the career of students who are suffering from depression.
- xii. To make the ultimate fulfillment of the dreams of students and their families.

## **2 Literature Review**

There are some previous works based on factors detection of depression and suicidal ideation. Various models and approaches are used in these, which are mentioned here.

M Ashraful Islam et al. [11] in 2018, studied to explore the factors of depression among Malaysian university students. They found that second year students are more depressed than first year students. They also found that family background is a key factor of depression.

Karl Peltzer et al. [12] in 2014, studied the risk factors of depression among Western Nigeria. They surveyed 820 university students and found sleeping problem as a risk factor of depression.

In 2020, Kumar et al. [13] exhibited various models to detect early suicidal ideation using Twitter data through sentiment analysis and supervised learning methods. This work used 60,188 positive and negative tweets, and among these, 86.42% were suicidal tweets. They proposed additional keywords of n-grams as search key phrases and used Vader sentiment analysis for data separation.

Sparsh Sharma & Surbhi Sharma in 2020 [14] analyzed the depression and suicidal tendencies using sentiment analysis on social networking sites of persons affected by Corona Virus Disease called COVID-19's lockdown. They proposed a Sentiment Analysis Framework that can extract the emotions and sentiments of people from Twitter and report the location and required pieces of information to the authority to save lives from the pandemic.

In 2021, Jung et al. [15] implemented a suicidality detection model for Twitter data using a machine learning approach. They randomly selected 20,000 tweets and analyzed metadata and text features to build this effective model. They studied Metadata features to understand their possibility and importance in suicidality detection models. By integrating metadata and text features, they obtained the model of good performance (F1 score of 0.846) that can help humans to detect suicidality using social media posts.

In 2021, Roy et al. [16] developed SAIPH to predict future risk of suicidal thought by using Twitter data. They used the Tweepy package in python and accessed the Twitter API, allowing access to the past nine days of publicly available Twitter content for a given query.

In 2018, Katchapakirin et al. [17] developed a depression detection algorithm for the Thai language on Facebook, which people use as a tool for sharing opinions, feelings, and life events using Natural Language Processing (NLP) techniques. They studied that Facebook behaviors could predict depression levels.

To summarize, the previous literature indicates that online behaviors and activities could predict depression, its level, and suicide prediction. This work collects data from Bangladeshi university students and analyze these data to find out factors of depression.

### 3 Methodology

This research work has gone through several processes like questionnaire creation, data collection, preprocessing and analysis.

#### 3.1 Data collection

This study is carried out to collect data by online questionnaire system. There were some questions to answer and submit the google form. This questionnaire was distributed to university students. Questionnaire is set to know the gender, study year, CGPA and factors which are responsible for depression and suicidal ideation as responders' opinion. Responder can select multiple factors option from the following options and also can add others option as wish.

**Table 1.** Sample Dataset


Gender	Study Year	CGPA	Factors
Male	Second Year	3.00-3.50	Academic Pressure, Family Expectations
Female	First Year	<3.00	Uncertain Future
Female	Fourth Year	3.00-3.50	Loneliness, Financial Crisis, Academic Pressure
Male	Third Year	>3.50	Unemployment, Job Pressure

Table 1 shows the dataset sample. These factors are Loneliness, Hopelessness, Relationship Issues, Helplessness, Grade Problems, Parental Problems, Financial

Crisis, Academic Pressure, Social Comparison, Social Media Influence, Family Expectations, Lack of Sleep, Uncertain Future, Health Issues, Bullying, Substance Abuse, Unemployment, Job Pressure, Working Environment, Monotony and Sexual Frustration.

### 3.2 Model

In the depression and suicidal ideation factors identification stage, a factors list is created from the data which are collected from university students. A questionnaire was designed to collect data from university students. After completion of the questionnaire design, it deployed to the Bangladeshi university students. Figure 1 shows the system architecture.



**Fig. 1.** System Structure

When data collection was completed, the data cleaning and normalization process has held. Then data analyzed to identify the depression factors. From the data analysis approach, a factor list of depression found for Bangladeshi university students. This factor list help to detect depression among students.

Apriori association algorithm is applied to find out the factors list with related support, confidence and lift value. Support, confidence and lift can be defined by –

$$\text{Support } \{x\} = \text{Number of sets with } x / \text{Total number of factors sets}$$

$$\text{Confidence } \{x \rightarrow y\} = \text{Support } \{x,y\} / \text{Support } \{x\}$$

$$\text{Lift } \{x \rightarrow y\} = \text{Support } \{x,y\} / (\text{Support } \{x\} * \text{Support } \{y\})$$

This algorithm also makes a relationship between antecedent support and consequent support for various factors length.

## 4 Result and Discussion

The collected data using questionnaire is analyzed to get findings. The research has 348 participants, using Google Form circulated to Bangladesh university students. The following table 1 shows the profile background of respondents among university students in Bangladesh.

**Table 2.** Summary of Data

ITEMS	FREQUENCY	PERCENTAGE(%)
<b>Gender</b>		
Male	196	56.32%
Female	152	43.68%
<b>Study Year</b>		
First Year	88	25.29%
Second Year	64	18.39%
Third Year	136	39.08%
Fourth Year	60	17.24%
<b>CGPA</b>		
>3.5	104	29.89%
3.0-3.5	184	52.87%
<3.0	60	17.24%

#### 4.1 Percentage of Factors


In this work, there were 21 options and students can choose multiples. This work found that 72.41% of the surveyed student thinks academic pressure is a reason of depression and suicidal event. Second most 56.32% students think uncertain future is key reason.

**Table 3.** Percentage of various factors for depression and suicidal ideation

FACTORS	PERCENTAGE(%)
Loneliness	41.38%
Hopelessness	48.28%
Relationship Issues	31.03%
Helplessness	22.99%
Grade Problems	35.63%
Parental Problems	8.05%
Financial Crisis	42.53%
Academic Pressure	72.41%
Social Comparison	28.74%
Social Media Influence	17.24%
Family Expectations	47.13%
Lack of Sleep	26.44%
Uncertain Future	56.32%

<b>Health Issues</b>	12.64%
<b>Bullying</b>	20.69%
<b>Substance Abuse</b>	6.90%
<b>Unemployment</b>	37.93%
<b>Job Pressure</b>	27.59%
<b>Working Environment</b>	12.64%
<b>Monotony</b>	18.39%
<b>Sexual Frustration</b>	1.15%

According to the surveyed 48.28%, 47.13%, 42.53%, 41.38% and 37.93% students' opinion hopelessness, family expectations, financial crisis, loneliness and unemployment respectively are responsible for depression and suicidal ideation among university students. Figure 2 shows the graphical representation of table 3.



**Fig. 2.**Graphical representation for percentage of various factors

#### 4.2Gender Based Percentage of Factors


Percentage of factors for depression and suicidal ideation differs based on gender. Male students thinking and female students thinking varies in some points. In this work, there were 56.32% male and 43.68% female students are in survey who can choose multiple options from the factors. This study found that most of the students, 40.23% and 32.18% of the surveyed male and female student respectively think academic pressure is a primary reason of depression and suicidal event. Table 4 shows the gender-based percentage of various factors for depression and suicidal ideation.

**Table4.** Gender based percentage of various factors for depression and suicidal ideation

<b>FACTORS</b>	<b>MALE (%)</b>	<b>FEMALE (%)</b>
<b>Loneliness</b>	20.69%	20.69%
<b>Hopelessness</b>	21.84%	26.44%

<b>Relationship Issues</b>	17.24%	13.79%
<b>Helplessness</b>	13.79%	9.20%
<b>Grade Problems</b>	13.79%	21.84%
<b>Parental Problems</b>	5.75%	2.30%
<b>Financial Crisis</b>	28.74%	13.79%
<b>Academic Pressure</b>	40.23%	32.18%
<b>Social Comparison</b>	17.24%	11.49%
<b>Social Media Influence</b>	13.79%	3.45%
<b>Family Expectations</b>	28.74%	18.39%
<b>Lack of Sleep</b>	11.49%	14.94%
<b>Uncertain Future</b>	29.89%	26.44%
<b>Health Issues</b>	9.20%	3.45%
<b>Bullying</b>	9.20%	11.49%
<b>Substance Abuse</b>	4.60%	2.30%
<b>Unemployment</b>	25.29%	12.64%
<b>Job Pressure</b>	17.24%	10.34%
<b>Working Environment</b>	9.20%	3.45%
<b>Monotony</b>	10.34%	8.05%
<b>Sexual Frustration</b>	1.15%	0.00%

Second most 29.89% male students think uncertain future is key reason but 26.44% female students think hopelessness as a reason. 28.74% male students have opinion about financial crisis as a factor but only 13.79% female students think financial crisis as a factor. 25.29% male students have opinion about unemployment but only 12.64% female students have opinion about unemployment as a factor. Figure 3 shows the graphical representation of table 4.



**Fig. 3.**Graphical representation for gender percentage of various factors

#### 4.3Apriori Association Rule on Factors



Using the Apriori association rule, the support is computed for single items means single factors. Table 5 shows the single factors list with the support value.

**Table 5.** Support for single factors

<b>FACTORS</b>	<b>Support (Single Factor)</b>
<b>Loneliness</b>	0.021277
<b>Hopelessness</b>	0.021277
<b>Relationship Issues</b>	0.021277
<b>Helplessness</b>	0.021277
<b>Grade Problems</b>	0.063830
<b>Parental Problems</b>	0.063830
<b>Financial Crisis</b>	0.191489
<b>Academic Pressure</b>	0.255319
<b>Social Comparison</b>	0.234043
<b>Social Media Influence</b>	0.148936
<b>Family Expectations</b>	0.489362
<b>Lack of Sleep</b>	0.319149
<b>Uncertain Future</b>	0.638298
<b>Health Issues</b>	0.148936
<b>Bullying</b>	0.255319
<b>Substance Abuse</b>	0.063830
<b>Unemployment</b>	0.489362
<b>Job Pressure</b>	0.425532
<b>Working Environment</b>	0.170213
<b>Monotony</b>	0.255319
<b>Sexual Frustration</b>	0.936170

From the Apriori association algorithm, this work found some associations. The following associations are of frequent factors for higher support.

*Family Expectations -> Financial Crisis*

*Monotony -> Uncertain Future*

*Social Media Influence -> Unemployment*

*Uncertain Future -> Family Expectations*

*Unemployment -> Family Expectations*

*Job Pressure -> Unemployment*

*Job Pressure -> Uncertain Future*

*Financial Crisis -> Parental Problems*

*Financial Crisis -> Social Comparison*

*Monotony -> Bullying*

*Social Media Influence -> Bullying*

*Social Media Influence -> Family Expectations*

*Social Media Influence -> Financial Crisis*

*Unemployment -> Financial Crisis*


*Job Pressure -> Social Media Influence*

*Social Media Influence -> Monotony*

*Job Pressure -> Monotony*


### *Unemployment -> Monotony*

Figure 4 shows the scatter plot of antecedent support and consequent support for the association rule which have minimum support 0.005 and maximum length 2.




**Fig. 4.**Scatter plot of antecedent support and consequent support (Max Length 2)

Figure 5 shows the scatter plot of antecedent support and consequent support for the association rule which have minimum support 0.005 and maximum length 3.




**Fig. 5.**Scatter plot of antecedent support and consequent support(Max Length 3)

Figure 6 shows the scatter plot of support, confidence and lift for the association rule which have minimum support 0.005 and maximum length 2. Here X axis refers to the support, Y axis refers to the confidence and Size means the lift.



**Fig. 6.**Scatter plot of Support, Confidence and Lift (Max Length 2)

Figure 7 shows the scatter plot of support, confidence and lift for the association rule which have minimum support 0.005 and maximum length 3. Here X axis refers to the support, Y axis refers to the confidence and Size means the lift.



**Fig. 7.**Scatter plot of Support, Confidence and Lift (Max Length 3)

This work found the support list for single factors, association factors frequent list, scatter plot of antecedent support and consequent support for maximum length 2 and 3, also found the scatter plot of support, confidence and lift for maximum length 2 and 3.

#### 4Conclusion and Future Work

This study found that Academic Pressure, Uncertain Future, Hopelessness, Family Expectations, Financial Crisis, Loneliness, Unemployment, Grade Problems, Relationship Issues and Social Comparison are the key factors for depression and suicidal ideation among university student in Bangladesh. These factors percentage varies based on gender male and female. Academic Pressure, Uncertain Future, Financial Crisis, Family Expectations, Unemployment, Hopelessness, Loneliness and Relationship Issues are the key factors according to male students. On the other hand, Academic Pressure, Hopelessness, Uncertain Future, Grade Problems, Loneliness, Family Expectations, Lack of Sleep and Relationship Issues are the key factors according to female students. We can save many students lives by concerning and stopping suicide cases. Depression and suicidal incident are very much possible to stop it by precaution and taking proper steps on the factors. In future, we can work on student's mental health, depression to stop suicidal incidents.

## References

- [1] Sarokhani, D., Delpisheh, A., Veisani, Y., Sarokhani, M. T., Manesh, R. E., & Sayehmiri, K. (2013). Prevalence of depression among university students: a systematic review and meta-analysis study. *Depression research and treatment*, 2013.
- [2] Othieno, C. J., Okoth, R. O., Peltzer, K., Pengpid, S., & Malla, L. O. (2014). Depression among university students in Kenya: Prevalence and sociodemographic correlates. *Journal of affective disorders*, 165, 120-125.
- [3] Islam, M. A., Low, W. Y., Tong, W. T., Yuen, C. W., & Abdullah, A. (2018). Factors associated with depression among university students in Malaysia: a cross-sectional study. *KnE Life Sciences*, 415-427.
- [4] Orsal, O., Orsal, O., Unsal, A., & Ozalp, S. S. (2013). Evaluation of internet addiction and depression among university students. *Procedia-Social and Behavioral Sciences*, 82, 445-454.
- [5] Park, Y. R., & Jang, E. H. (2013). Impact of stress on depression among university students: testing for moderating effect of social support. *Korean Journal of Adult Nursing*, 25(5), 549-558.
- [6] Santos, H. G. B. D., Marcon, S. R., Espinosa, M. M., Baptista, M. N., & Paulo, P. M. C. D. (2017). Factors associated with suicidal ideation among university students. *Revistalatio-americana de enfermagem*, 25, e2878.
- [7] Arria, A. M., O'Grady, K. E., Caldeira, K. M., Vincent, K. B., Wilcox, H. C., & Wish, E. D. (2009). Suicide ideation among college students: A multivariate analysis. *Archives of suicide research*, 13(3), 230-246.
- [8] Medrano, J. L. J., Lopez Rosales, F., & Gámez-Guadix, M. (2018). Assessing the links of sexting, cybervictimization, depression, and suicidal ideation among university students. *Archives of suicide research*, 22(1), 153-164.
- [9] Tasnim, R., Islam, M. S., Sujon, M. S. H., Sikder, M. T., & Potenza, M. N. (2020). Suicidal ideation among Bangladeshi university students early during the COVID-19 pandemic: Prevalence estimates and correlates. *Children and youth services review*, 119, 105703.
- [10] Lin, C. C. (2015). The relationships among gratitude, self-esteem, depression, and suicidal ideation among undergraduate students. *Scandinavian journal of psychology*, 56(6), 700-707.
- [11] Islam, M. A., Low, W. Y., Tong, W. T., Yuen, C. W., & Abdullah, A. (2018). Factors associated

with depression among university students in Malaysia: a cross-sectional study. *KnE Life Sciences*, 415-427.

[12] Peltzer, K., Pengpid, S., Olowu, S., & Olasupo, M. (2013). Depression and associated factors among university students in Western Nigeria. *Journal of psychology in Africa*, 23(3), 459-465.

[13] Rajesh Kumar, E., Rama Rao, K. V. S. N., Nayak, S. R., & Chandra, R. (2020). Suicidal ideation prediction in twitter data using machine learning techniques. *Journal of Interdisciplinary Mathematics*, 23(1), 117–125. doi:10.1080/09720502.2020.1721674

[14] Sharma, S., & Sharma, S. (2020). Analyzing the depression and suicidal tendencies of people affected by COVID-19's lockdown using sentiment analysis on social networking websites. *Journal of Statistics and Management Systems*, 24(1), 115–133. doi:10.1080/09720510.2020.1833453

[15] Jung, W., Kim, D., Nam, S., & Zhu, Y. (2021). Suicidality Detection on Social Media Using Metadata and Text Feature Extraction and Machine Learning. *Archives of Suicide Research*, 1–16. doi:10.1080/13811118.2021.1955783

[16] Roy, A., Nikolitch, K., McGinn, R. et al. A machine learning approach predicts future risk to suicidal ideation from social media data. *npj Digit. Med.* 3, 78 (2020). <https://doi.org/10.1038/s41746-020-0287-6>

[17] K. Katchapakirin, K. Wongpatikaseree, P. Yomaboot and Y. Kaewpitakkun, "Facebook Social Media for Depression Detection in the Thai Community," 2018 15th International Joint Conference on Computer Science and Software Engineering (JCSSE), 2018, pp. 1-6, doi: 10.1109/JCSSE.2018.8457362.