

Original Research Article

Inventory of Herbaceous Species and Bioaccumulation of Heavy Metals in their various Parts: Case in the Urban Ecosystem of Ngaoundere, Cameroon.

ABSTRACT

This study was carried out to inventory herbaceous species and to assess the bioaccumulation of heavy metals in these species. The ecosystem chosen was an area of the city of Ngaoundere (Cameroon) divided into three sites, namely a control site and two sites subject to heavy metal pollution. In order to compare the potential for bioaccumulation, two species were assessed. It shows that out of 19 herbaceous species present in the study area, five are very abundant in all three sites regardless of the degree of pollution. Concerning the bioaccumulation, the two herbaceous plants chosen do not absorb the same amounts of heavy metals. Indeed, the concentrations show that the concentration of cadmium, copper, iron, nickel and lead at all three sites is higher in *C. benghalensis* than in *A. conyzoides*. The concentration of zinc at the control site and the hospital site is higher in *C. benghalensis* while at the prison site this concentration is higher in *A. conyzoides*. The concentration of cadmium is higher in the leaves than in the other parts in both species. Copper is stored at the roots in both species. *A. conyzoides*, mainly accumulates Iron in the roots and *C. benghalensis*, accumulates a significant concentration of Iron throughout the plant. *A. conyzoides* and *C. benghalensis* store Nickel more in the roots than in the other parts. Regarding lead, the species studied store it more in the roots followed by the leaves and finally the stems. As for Zinc, *A. conyzoides* and *C. benghalensis* store it more in the roots than in the other parts. The bioaccumulation of these heavy metals in edible herbaceous plants could be a threat if these pollutants enter the human food chain.

Keywords: Inventory, herbaceous plants, heavy metals, bioaccumulation, Ngaoundere, Cameroon.

1. INTRODUCTION

One of the ecosystems that is under strong pressure in terms of pollution is the urban ecosystem [1]. Urban ecosystems in developing countries suffer more and the causes are not lacking. These include the lack of planning or anarchic development of cities; promiscuity; poverty, presence or proximity to industrial areas, intense road traffic with its corollary air pollution; discharge of wastewater with its corollary soil pollution etc. Ngaoundere, a medium-sized town located in the Adamawa region of Cameroon is no exception to this observation. Recent studies have revealed the pollution of water [2] and soil [3] in this city. One consequence is the transfer of these pollutants into plants via soil, water or air. Thus, pollution can influence the presence or absence of certain plant species. Pollution can also influence the distribution of these pollutants in the differences of these plants once absorbed [4,5]. Heavy metals contamination of soils, following local atmospheric fallout (industrial and urban) and various inputs (sludge from wastewater treatment plants, composts, fertilizers, etc.) can explain the high concentrations of heavy metals such as cadmium (Cd), copper (Cu), zinc (Zn) and lead (Pb) in certain plant species [4]. Their transfer to plants may pose a risk to human health through contamination of the human food chain [6].

Plants are exposed to heavy metals in two ways: through the air parts and through the roots. Heavy metals can be deposited on the surface of leaves and roots or penetrate into the plant. Heavy metals can enter through aerial parts (leaves, stems and fruits), from airborne particles, gaseous compounds or compounds dissolved in rainwater or irrigation water. They can penetrate through the roots from the ground. Once removed by the plant, heavy metals may be trapped and not circulating in the plant, or transported from the place of absorption to another plant organ [4].

The main objective of this study is on the one hand to inventory herbaceous species and on the other hand to evaluate the bioaccumulation of heavy metals in herbaceous plants of an urban ecosystem that constitutes the area around the flow channel of a river called *Soumsoum* in the city of Ngaoundere in Cameroon.

2. MATERIEL ET METHODES

2.1. Study Area

Figure 1 shows the geographical location of the study area and the three main sites where the plant samples were collected. There are three sites S1; S2 and S3. S1 is the control area with the points P13 to P17, S2 is the prison's site with the locations P1 to P6 and S3 is the hospital's site with the points P7 to 12. Plants from the site of central prison and the Ngaoundere regional hospital were compared with the plants of control site. The control site is located upstream of the two above-mentioned sites and does not suffer from any pollution. Three replicates were performed and averages were obtained.




Figure 1: Localisation of the study area.

2.2. Experimental set-up and herbaceous inventory

Floristic surveys of herbaceous species along the river were carried out at 03 sites on both sides of the river (Control, Prison and Hospital). The herbaceous inventory was carried out using the minimum area method. It is the smallest area that contains all the herbaceous species. By a plotted representation, it is the area at which the curve becomes horizontal [7–9]. This made it possible to quantify the floristic composition of the plant cover on the basis of the centesimal frequency or relative abundance of the herbaceous species recorded.

2.3. Collection of plant samples

Sampling was carried out in such a way as to preserve information, using sharp ceramic tools (scissors) for the stems and ordinary steel-based materials for the roots. Leaves were removed by hand or with scissors. After harvesting, the samples were placed in a polyethylene bag. Between collection and transfer to the laboratory, the samples were placed in a cool, dark container, a cool box.

Quantity relative to the number of individuals of a given species per unit area or volume relative to the total number of individuals of all species combined.

The plant samples were dissolved and then assayed for heavy metals using a flame atomic spectrophotometer. All analyses were carried out three times by flame atomic spectrophotometer (Aurora Instruments Ltd-AI 1200).

2.4. Statistical analysis

The data were analyzed using STATGRAPHICS plus 5.0. Significance tests were performed using ANOVA and Duncan's test at the 5% level.

3. RESULTS AND DISCUSSION

3.1. Abundance of herbaceous species.

A total of 19 herbaceous species divided into 19 genera and 12 families were recorded in the three study areas. Table 1 shows that five (05) herbaceous species, namely *Ageratum conyzoides*, *Amaranthus spinosus*, *Bidens pilosa*, *Commelinabenghalensis* and *Galinsoga parviflora* are the most abundant in the three study zones. The very high abundance at all three sites can be explained by the fact that these are invasive species. [10–12]. These species can grow in both healthy and polluted soils. While the control site has 12 species, the prison and hospital sites have 16 and 14 species respectively. The low number of species on the control site compared with the other two sites can be explained by the pollution of the water. [2]andsoils[3]from the latter two sites.

Table 1: Abundance of herbaceous plants species.

Herbaceous species	Familie	Sites		
		Control	Prison	Hospital
<i>Aundo donax</i>	Poaceae	+	+	0
<i>Pergulariatomentosa</i>	Apocynaceae	0	+	0
<i>Zeamays</i>	Poaceae	+	+	+
<i>Tithoniadiversifolia</i>	Asteraceae	0	++	+
<i>Musa paradisiaca</i>	Musaceae	++	0	+
<i>Euphorbiahirta</i>	Euphorbiaceae	0	+	+
<i>Ricinuscommunis</i>	Euphorbiaceae	+	+	+
<i>Ageratum conyzoides</i>	Asteraceae	+++	+++	+++
<i>Amaranthus spinosus</i>	Amaranthaceae	+++	+++	+++
<i>Bidens pilosa</i>	Asteraceae	+++	+++	+++
<i>Hemerocallisfulva</i>	Xanthorrhoeaceae	0	+	0
<i>Galeopsis tetrahit</i>	Lamiaceae	+	+	0
<i>Sparmanniaafricana</i>	Malvaceae	0	+	+
<i>Panicum capildare</i>	Poaceae	+	+	+
<i>Vaccinium myrtillus</i>	Ericaceae	0	0	+
<i>Centellaasiatica</i>	Apiaceae	0	0	+
<i>Commellinabenghalensis</i>	Commelinaceae	+++	+++	+++
<i>Galinsogaparviflora</i>	Asteraceae	+++	+++	+++
<i>Eurybiadivaricata</i>	Asteraceae	+	+	0

Absence of the specie: 0; abundance; +; Medium abundance: ++; High abundance:+++.

3.2. Description of the most abundant species.

Ageratum conyzoides

A. conyzoides belong to the Asteraceae family (Figure 2) is an annual herb that grows on ferruginous soils or on rubbish tips. It is an annual tropical herbaceous species [13,14]. It is well established in tropical rainforests in India [15]. In Cameroon, it is found in most regions in fallow fields and around dwellings [16]. Its stem is cylindrical, often reddish and hairy, and twiggy from the base. The lower leaves are opposite, hairy and oval-oblong, while the others are almost rhombic[17].



Figure 2: *Ageratum conyzoides*.

Amaranthus spinosus

A. spinosus (Figure 3) a plant in the *Amarantaceae* family known as spiny brenna. Amaranths are classified as one of the most troublesome weeds in agricultural production systems [18]; thanks to their aggressive growth, amaranths have the ability to compete with crops for water, nutrients and light, causing severe reductions in crop yield and quality [19].



Figure 3: *Amaranthus spinosus*.

Bidens pilosa

B. pilosa (Figure 3) is a weed of crops in a wide variety of latitudes and on various continents [20]. It is native to the tropical forests of South America, Africa, the Caribbean and the Philippines [14]. *B. pilosa* is an annual herb in the *Asteraceae* family that is widespread throughout the world, particularly in tropical and subtropical regions[21].



Figure 4 :*Bidens pilosa*

Commelinabenghalensis

Also known as big waterweed, *C. benghalensis* (Figure) is a plant in the *Commelinaceae* family that grows in wet soils during the rainy season[22]. It is a perennial herb in tropical regions[23], but grows as an annual grass in temperate regions such as Georgia in the United States of America, where it is classified as one of the worst weeds affecting crop production in many African and Asian countries[24]. *C. benghalensis* has been found to be an indicator of pollution[25].

Figure 5: *Commelinabenghalensis*.



Galinsoga parviflora

G. parviflora (Figure 4) is a medium-sized (20-60 cm) annual herbaceous plant belonging to the *Asteraceae* family[26,27]. It has an erect, rigid stem that is strongly branched and covered with stiff, shaggy hairs. Its leaves are green to apple-green, heavily pubescent, triangular, with embedded veins and clearly toothed margins[28].



Figure 6: *Galinsoga parviflora*

3.3. Bioaccumulation of heavy metals.

Table 2 shows the total concentrations of each pollutant studied in both plant species. The highest concentrations are obtained at the hospital site, followed by the prison site and finally the control site. The quantitative distribution of heavy metal contents is as follows regardless of the site and plant species: Fe>Cu>Ni>Zn>Pb>Cd. Compared to the maximum allowable values for plants by FAO/WHO, the concentrations are in decaf, except for cadmium and lead. Authors (Alowat and Steinnes, 1999; Greger, 1999; Felix et al.; 1999) reported comparable concentrations of cadmium in species such as *Populus tremula*, *Salix aurita* and *Thlaspi caerulescens*.

Table 2: Total concentrations (mg.kg⁻¹) in plant.

Plant specie	Sites/Norm	Total heavy metal concentration (mg/kg)					
		Cd	Cu	Fe	Ni	Pb	Zn
<i>A. conyzoides</i>	Control	0.28±0.1	10.3±2.4	330±25	7.23±1.3	5.3±0.4	30.3±1.2
	Prison	2.15±0.3	20.2±3.2	175±14	15.32±2.4	15±1.8	125.7±1.5
	Hospital	2.72±0.2	30.2±5.3	222±30	21.76±3.2	13.1±2.2	157.6±1.7
<i>C. benghalensis</i>	Control	0.31±0.1	14.5±4.2	380±35	8.3±0.8	3.5±0.6	33.5±1.3
	Prison	2.17±0.3	28.8±2.5	165±15	16.35±3.5	16.2±3.4	190.2±1.7
	Hospital	2.75±0.2	37.1±3.1	242±21	22.9±5.1	13.3±3.1	110.12±1.8
	FAO/WHO	0.2	73.3	425.5	67.9	0.3	99.4

Bioaccumulation du Cadmium

Figure 7 below shows the results of the determination of cadmium in the leaves, stems and roots of *A. conyzoides* and *C. benghalensis*; sampled at the sites of the control area, Hospital and Prison.




Figure 7 : Cadmium content (mg.kg⁻¹ dry matter) in the parts of *A. conyzoides* and *C. benghalensis* in the three sites.

In *A. conyzoides*, they evolve from 0.07 ± 0.01 to 1.62 ± 0.01 mg.kg⁻¹ in leaves, from 0.05 ± 0.00 to 0.81 ± 0.01 mg.kg⁻¹, in stems and from 0.02 ± 0.01 to 0.24 ± 0.02 mg.kg⁻¹ in roots. In *C. benghalensis*, the concentration of Cd varies from 0.09 ± 0.02 to 2.23 ± 0.03 mg.kg⁻¹ in leaves, from 0.07 ± 0.01 to 1.44 ± 0.02 mg.kg⁻¹ in stems and finally in roots from 0.04 ± 0.01 to 0.81 ± 0.02 mg.kg⁻¹. The highest concentration of Cd is observed in the leaves of *C. benghalensis* at the hospital site (2.23 ± 0.03 mg.kg⁻¹). Cadmium has high assimilability and mobility, it enters plants via the roots and is translocated in the leaves [29,30]. *A. conyzoides* and *C. benghalensis* mainly accumulate cadmium in leaves [31]. Both species studied have a strong accumulation power rather in leaves such as tobacco (*Nicotiana tabacum* L.). Other factors such as plant species, intense root activity, and soil characteristics suggest that accumulation in leaves may favour accumulation in leaves (Kabata-Pendias and Mukherjee 2007; Prasad and Freitas, 2003; Yang et al., 2017). The plants from the Hospital site accumulate more, followed by those from the Prison and finally from the control area. This is due to the higher amounts of cadmium in the hospital site, followed by the prison site and in the control site [3]. In *A. conyzoides*, and *C. benghalensis*, the analysis of variances shows that there are significant variations ($P < 0.05$) between cadmium concentration in different parts of plants at different sites.

Bioaccumulation of Copper

Figure 8 below shows the results of the determination of copper in leaves, stems and roots of *A. conyzoides* and *C. benghalensis*; sampled at the sites of the control area, Hospital and Prison




Figure 8: Copper content (mg.kg^{-1} dry matter) in the parts of *A. conyzoides* and *C. benghalensis* in the three sites.

In *A. conyzoides*, they range from 1.17 ± 0.03 to $8.03 \pm 0.05 \text{ mg.kg}^{-1}$ in leaves, from 1.12 ± 0.03 to $3.90 \pm 0.01 \text{ mg.kg}^{-1}$ in stems and from 8.15 ± 0.01 to $18.83 \pm 0.02 \text{ mg.kg}^{-1}$ in roots. In *C. benghalensis*, the Cu concentration varies from 1.26 ± 0.04 to $9.09 \pm 0.09 \text{ mg.kg}^{-1}$ in leaves, from 1.90 ± 0.09 to $5.90 \pm 0.03 \text{ mg.kg}^{-1}$ in stems and in roots from 12.05 ± 0.06 to $23.90 \pm 0.40 \text{ mg.kg}^{-1}$. The highest concentrations in roots, stems, and leaves are observed in *C. benghalensis* at the hospital site. *A. conyzoides* and *C. benghalensis* mainly accumulate Cu in roots where more than 50% of the total copper is retained in it compared to other parts of the plant. Similar observations have been reported for endemic species such as *Phleum pratense*; *Thymus kotschyanus*; *Achillea millefolium* and *Trifolium pratense* (Ghazaryan et al. 2019). Both species can be used as phytoextractors in the remediation of copper-contaminated soils. Cultivated species such as maize exhibit an ability to store copper more in the roots than in other parts [36]. The plants from the Hospital site accumulate more, followed by those from the Prison and finally from the control area. As for cadmium, the analysis of variances shows that there are significant variations ($P < 0.05$) between the copper concentration in the different parts at the different sites. Overall, these plants store more in the roots than in other parts of the plant.

Bioaccumulation of iron

La concentration en Fer dans les différentes parties de deux plantes dans les sites est présentée à la Figure 9. In *A. conyzoides*, they range 53.53 ± 0.96 to $93.21 \pm 9.31 \text{ mg.kg}^{-1}$ in leaves, from 37.21 ± 0.66 to $76.08 \pm 0.90 \text{ mg.kg}^{-1}$ in stems and finally in roots and 96.55 ± 0.48 to $183.01 \pm 0.48 \text{ mg.kg}^{-1}$. *A. conyzoides*, mainly accumulate iron in the roots. In *C. benghalensis*, the concentration of Iron varies from 63.23 ± 0.67 to $128.23 \pm 0.86 \text{ mg.kg}^{-1}$ in the leaf, from 52.23 ± 0.07 to $115.55 \pm 0.17 \text{ mg.kg}^{-1}$ in the stem and from 86.16 ± 0.74 to $196.15 \pm 2.53 \text{ mg.kg}^{-1}$ in the roots. Like *A. conyzoides*, *C.*

benghalensis accumulates a large amount of iron in the roots. The iron levels in the above-ground parts, i.e. in the leaves and stems, are substantially similar. Similar observations for *C. benghalensis* have been reported by other authors (Sekabira et al. 2011). This can make these two species potential plants in the phytostabilization of iron in soils polluted with this heavy metal. However, this ability is low compared to species such as *Alyssum bertoloni*, which can accumulate more than three the iron content in the roots compared to the leaves [38].




Figure 9: Iron content (mg.kg⁻¹ dry matter) in the parts of *A. conyzoides* and *C. benghaensis* in the three sites.

The highest concentration in roots, stems, and leaves is observed in *A. conyzoides* at the control site. The plants in the control zone accumulate more iron, followed by the Hospital and the end of the Prison site. As for cadmium, copper, analysis of variances shows that there are significant variations ($P < 0.05$) between the iron concentration in different parts at different sites.

Bioaccumulation of Nickel

Figure 9 shows the results of the determination of Nickel in the leaves, stems and roots of *A. conyzoides* and *C. benghalensis*; sampled at the sites of the control area, Hospital and Prison.




Figure 10: Nickel content (mg.kg^{-1} dry matter) in the parts of *A. conyzoides* and *C. benghalensis* in the three sites.

In *A. conyzoides*, they evolve from 1.31 ± 0.00 to $2.98 \pm 0.10 \text{ mg.kg}^{-1}$ in the leaf, from 1.23 ± 0.01 to $1.53 \pm 0.12 \text{ mg.kg}^{-1}$ in the stem and from 4.02 ± 0.10 to $18.31 \pm 0.00 \text{ mg.kg}^{-1}$ in the root. In *C. benghalensis*, the concentration of Ni varies from 1.83 ± 0.01 to $4.28 \pm 0.19 \text{ mg.kg}^{-1}$ in the leaves, from 1.05 ± 0.17 to $1.87 \pm 0.00 \text{ mg.kg}^{-1}$ in stems and finally in roots and 6.06 ± 0.08 to $20.23 \pm 0.21 \text{ mg.kg}^{-1}$. The highest concentration of Ni in leaves and stems is observed in *C. benghalensis* at the hospital site. *A. conyzoides* and *C. benghalensis* mainly accumulate nickel in the roots as reported for plants (Kiran et al., 2022) and more specifically for *C. benghalensis* (Sekabira et al., 2011). Both species accumulate nickel concentrations in the roots more than 80% greater than in the above-ground parts. Both plant species are potential nickel hyperaccumulators in nickel-polluted soils. This observation has also been reported by some authors [40,41]. The plants from the Hospital site accumulate more, followed by those from the Prison and finally from the control area. As for cadmium, copper and iron, the analysis of variances shows that there are significant variations ($P < 0.05$) between the copper concentration in the different parts at the different sites.

Bioaccumulation of Lead

The distribution of lead in the different parts of two plant species is shown in Figure 11. In *A. conyzoides*, they vary from 1.87 ± 0.16 to $5.28 \pm 0.18 \text{ mg.kg}^{-1}$ in the leaves, from 1.22 ± 0.02 to $3.87 \pm 0.13 \text{ mg.kg}^{-1}$ in the stems and finally in the roots and 2.08 ± 0.05 to $8.08 \pm 0.00 \text{ mg.kg}^{-1}$. In *C. benghalensis*, the concentration of Pb varies from 1.15 ± 0.01 to $4.86 \pm 0.15 \text{ mg.kg}^{-1}$ at the leaf level, from 1.06 ± 0.01 to $2.86 \pm 0.01 \text{ mg.kg}^{-1}$ at the stem level and from 2.52 ± 0.1 to $9.95 \pm 0.04 \text{ mg.kg}^{-1}$.




Figure 11: Lead content (mg.kg^{-1} matter) in the parts of *A. conyzoides* and *C. benghalensis* in the three sites.

A. conyzoides and *C. benghalensis* accumulate lead more in the roots, followed by the leaves and finally in the stems. Plants generally follow this sequence (Kiran et al. 2022; Kumar et al., 2017). This sequence has been observed for *A. conyzoides* in polluted soils in Nigeria [43] and for *C. benghalensis* [37]. The highest concentration of lead is observed in roots of *C. benghalensis* at the Prison site. The lowest concentration was measured in *C. benghalensis* leaves at the control site. The plants of the Prison site accumulate more, followed by those of the Hospital and finally that of the control area. Analysis of variance shows that site, species and plant part have a significant effect ($P < 0.05$) on the variation in Pb levels. The small difference in the concentration of lead in the leaves and roots allows both species to be used as both phytoextractors and phytostabilizers.

Bioaccumulation of Zinc

Figure 12 shows the results of the determination of Zinc in the leaves, stems and roots of *A. conyzoides* and *C. benghalensis*; sampled at the sites of the control area, Hospital and Prison.




Figure 12: Zinc content (mg.kg⁻¹dry matter) in the parts of *A. conyzoides* and *C. benghalensis* in the three sites.

In *A. conyzoides*, they evolve from 10.87 ± 0.19 to 113.86 ± 0.01 mg.kg⁻¹ at the leaf level, from 8.47 ± 0.52 to 93.83 ± 0.23 mg.kg⁻¹ at the stem level and from 17.27 ± 0.12 to 161.83 ± 0.27 mg.kg⁻¹. In *C. benghalensis*, the Zn concentration varies from 12.15 ± 0.02 to 99.07 ± 0.09 mg.kg⁻¹ in the leaves, from 8.01 ± 0.0 to 85.42 ± 0.03 mg.kg⁻¹ in stems and finally in roots and 21.83 ± 0.03 to 147.56 ± 0.02 mg.kg⁻¹. The highest concentration of zinc is observed in roots of *C. benghalensis* at the hospital site. The plants from the Hospital site accumulate more, followed by those from the Prison and finally from the control area. The distribution of zinc in the different parts of two plants shows only a slight difference in terms of proportions. Both plants accumulate almost equal amounts of zinc in their leaves and roots. This was observed for *A. conyzoides* (Mkumbo S et al. 2012).

CONCLUSION

The evaluation of the level of toxicity of heavy metals (Fe, Cu, Ni, Zn, Cd and Pb) in the different parts of these species in comparison with the control area shows that plants from the Prison and Hospital areas have the highest concentrations. In all sites and in both species, Cd is the most concentrated element in the aerial part while Fe, Cu, Zn, Ni and Pb in the roots. The highest concentrations are obtained in *C. benghalensis*. The plants species of the Prison and Hospital sites are characterized by extremely high concentrations compared to FAO/WHO standards.

These results show the ability of these two species to serve as accumulators of heavy metals.

References :

1. Breuste, J.; Pauleit, S.; Haase, D.; Sauerwein, M. Urban Ecosystems. Function, Management and Development.; 2021; ISBN 9783662632789.

2. Boukar, H.; Hareinda, N.L.; Dalatou, T.; Adamou, I.; Nestor, N.M.J.J. Physico-Chemical Characterization and Assessment of the Risk of Pollution: A Case of Wastewater Generated by the Regional Hospital and the Central Prison of the City of Ngaoundere, Cameroon. *Int. J. Environ. Clim. Chang.* 2023, 13, 921–932, doi:10.9734/ijecc/2023/v13i92314.
3. Hareinda, N.L.; Boukar, H.; Robert, A.; Dalatou, T.; Adamou, I. Pollution Assessment of Two Urban Soils in the City of Ngaoundere, Cameroon. *J. Geogr. Environ. Earth Sci. Int.* 2023, 27, 83–92, doi:10.9734/jgeesi/2023/v27i9707.
4. Singh, P. Effect of Soil Polluted by Heavy Metals: Effect on Plants, Bioremediation and Adoptive Evolution in Plants. In *Plant Responses to Soil Pollution*; 2020; pp. 89–102 ISBN 978-981-15-4963-2.
5. Tremel-Schaub, A.; Feix, I. Contamination Des Sols - Transferts Des Sols Vers Les Plantes; 2005; ISBN 286883793X.
6. Singh, S.P.; Singh, M.K. Soil Pollution and Human Health BT - Plant Responses to Soil Pollution. In *Plant Responses to Soil Pollution*; Singh, P., Singh, S.K., Prasad, S.M., Eds.; Springer Singapore: Singapore, 2020; pp. 205–220 ISBN 978-981-15-4964-9.
7. Pradesh, A.; Phukan, M.M.; Boruah, T. Determination Of Minimum Quadrat Size For The Herbaceous Vegetation : A Case Study Of Dulpang Reserve Forest ., *Nat. Volatiles Essent. Oils* 2021, 8, 8355–8359.
8. Ramalakshmana, J.; Padal, S.B.; Pradesh, A.; Pradesh, A. Determination of Minimum Quadrat Size for Herbaceous Species in Andhra. 2022, 9515, 281–284, doi:10.36347/sajb.2022.v10i11.003.
9. Tchobsala Influence Des Coupes de Bois Sur La Dynamique de La Végétation Naturelle de La Zone Peri-Urbaine de Ngaoundéré (Adamaoua)., Université de Yaoundé, 2011.
10. Noba, K.; Bassene, C.; Ngom, A.; Gueye, M.; Camara, A.; Kane, M.; Ndoye, F.; Dieng, B.; Rmballo, R.; Ba, N.; et al. Invasive Plants of West Africa : Concepts , Overviews and Sustainable Management.; 2017; Vol. 2;.
11. Holou, R.A.Y.; Achigan-Dako, E.G.; Sinsin, B. Ecology and Management of Invasive Plants in Africa. In *Invasive Plant Ecology*; Jose, S., Singh, H.P., Batish, D.R., Ravinder, K.K., Eds.; Taylor & Francis Group, LLC, 2013; pp. 161–174 ISBN 9781439881279.
12. Djotan, A.K.G.; Aoudji, A.K.N.; Gbaguidi, G.C.R.; Akouehou, G.S.; Ganglo, J.C. Vulnérabilité Des Aires Protégées Du Bénin à l'invasion de *Ageratum Conyzoides* L. (Asteraceae) En Rapport Avec Les ChangementsClimatiques. *Eur. Sci. Journal, ESJ* 2018, 14, doi:10.19044/esj.2018.v14n33p313.
13. Okunade, A.L. *Ageratum Conyzoides* L. (Asteraceae). *Fitoterapia* 2002, 73, 1–16.
14. Carriere, M. Impact Des Systèmesd'élevagePastoraux Sur l'environnement En Afrique et En Asie Tropicale et SubtropicaleAride et Sub-Aride. *Cirad-Emvt* 1996, 70 p.
15. Swamy, P.S.; Sundarapandian, S.M.; Chandrasekar, P.; Chandrasekaran, S. Plant Species Diversity and Tree Population Structure of a Humid Tropical Forest in Tamil Nadu, India. *Biodivers. Conserv.* 2000, 9, 1643–1669, doi:10.1023/A:1026511812878.

16. Menut, C.; Lamaty, G.; Zollo, P.H.A.; Kuiate, J.R.; Bessière, J.M. Aromatic Plants of Tropical Central Africa. Part X Chemical Composition of the Essential Oils of *Ageratum Houstonianum* Mill. and *Ageratum Conyzoides* L. from Cameroon. *Flavour Fragr. J.* 1993, 8, 1–4, doi:<https://doi.org/10.1002/ffj.2730080102>.
17. Santos, R.F.; Nunes, B.M.; Sá, R.D.; Soares, L.A.L.; Randau, K.P. Morpho-Anatomical Study of *Ageratum Conyzoides*. *Rev. Bras. Farmacogn.* 2016, 26, 679–687, doi:<https://doi.org/10.1016/j.bjp.2016.07.002>.
18. Sellers, B.A.; Smeda, R.J.; Johnson, W.G.; Kendig, J.A.; Ellersieck, M.R. Comparative Growth of Six *Amaranthus* Species in Missouri. *Weed Sci.* 2003, 51, 329–333, doi:10.1614/0043-1745(2003)051[0329:cgosas]2.0.co;2.
19. Pigweed, R.; Echinochloa, B. Weed Science Society of America Retroflexus) and Barnyardgrass (*Echinochloa Crus-Galli*) Redroot Pigweed (*Amaranthus* Interference in Potatoes (*Solanum Tuberosum*) 1. 2015, 38, 338–343.
20. Bartolome, A.P.; Villaseñor, I.M.; Yang, W. *Bidens Pilosa* L. (Asteraceae): Botanical Properties , Traditional Uses , Phytochemistry , and Pharmacology. *Evidence-Based Complement. Altern. Med.* 2013, 2013, doi:10.1155/2013/340215.
21. Nezomba, H.; Mtambanengwe, F.; Tittonell, P.; Mapfumo, P. Practical Assessment of Soil Degradation on Smallholder Farmers' Fields in Zimbabwe: Integrating Local Knowledge and Scientific Diagnostic Indicators. *CATENA* 2017, 156, 216–227, doi:<https://doi.org/10.1016/j.catena.2017.04.014>.
22. Bàrberi, P. Chapter Four - Ecological Weed Management in Sub-Saharan Africa: Prospects and Implications on Other Agroecosystem Services. In; Sparks, D.L.B.T.-A. in A., Ed.; Academic Press, 2019; Vol. 156, pp. 219–264 ISBN 0065-2113.
23. Fibrich, B.; Lall, N. Chapter 11 - *Commelina Benghalensis*. In; Lall, N.B.T.-U.M.P. from S.-S.A., Ed.; Academic Press, 2020; pp. 77–85 ISBN 978-0-12-816814-1.
24. Holm, L.G.; Plucknett, Pancho, J. V.; Herberger, J. V The World's Worst Weeds : Distribution and Biology.; University Press of Hawaii: Honolulu :, 1977;
25. Noukeu, N.A.; Priso, R.J.; Dibong, S.D.; Ndongo, D.; Kono, L.; Essono, D. Floristic Diversity of Receiving Environments Polluted by Effluent from Agri-Food Industries. *Heliyon* 2019, 5, e02747, doi:<https://doi.org/10.1016/j.heliyon.2019.e02747>.
26. Riemens, M.M et Van den weide, R.Y. Biology and Control of *Galinsoga Parviflora*, Overview of a Literature Survey. *Plant Res. Int. B.V.*, wageningen 2008, 32.
27. Warwick, S.I.; Sweet, R.D. The Biology of Canadian Weeds 58. *Galinsoga Parviflora* and c. *Quadriradiata* (= *G. Ciliata*). *Can. J. Plant Sci.* 1983, 709, 695–709.
28. Ivany, J.A.; Sweet, R.D. Germination, Growth, Development, and Control of *Galinsoga*. *Weed Sci.* 1973, 21, 41–45, doi:DOI: 10.1017/S0043174500031647.
29. Dong, J.; Mao, W.H.; Zhang, G.P.; WU, F.B.; Cai, Y. Root Excretion and Plant Tolerance to Cadmium Toxicity - a Review. *Plant, Soil Environ.* 2007, 53, 193–200, doi:10.17221/2205-PSE.
30. Singh, P.; Siddiqui, H.; Sami, F.; Arif, Y.; Bajguz, A.; Hayat, S. Cadmium: A Threatening

- Agent for Plants BT - Plant Responses to Soil Pollution. In *Plant Responses to Soil Pollution*; Singh, P., Singh, S.K., Prasad, S.M., Eds.; Springer Singapore: Singapore, 2020; pp. 59–88 ISBN 978-981-15-4964-9.
31. Haider, F.U.; Liqun, C.; Coulter, J.A.; Cheema, S.A.; Wu, J.; Zhang, R.; Wenjun, M.; Farooq, M. Cadmium Toxicity in Plants: Impacts and Remediation Strategies. *Ecotoxicol. Environ. Saf.* 2021, 211, 111887, doi:<https://doi.org/10.1016/j.ecoenv.2020.111887>.
 32. Trace Elements of Group 8 (Previously Part of Group VIII) BT - Trace Elements from Soil to Human. In; Kabata-Pendias, A., Mukherjee, A.B., Eds.; Springer Berlin Heidelberg: Berlin, Heidelberg, 2007; pp. 209–224 ISBN 978-3-540-32714-1.
 33. Yang, Y.; Ge, Y.; Zeng, H.; Zhou, X.; Peng, L.; Zeng, Q. Phytoextraction of Cadmium-Contaminated Soil and Potential of Regenerated Tobacco Biomass for Recovery of Cadmium. *Sci. Rep.* 2017, 7, 7210, doi:[10.1038/s41598-017-05834-8](https://doi.org/10.1038/s41598-017-05834-8).
 34. Prasad, M.N.V.; Freitas, H.M. de O. Metal Hyperaccumulation in Plants - Biodiversity Prospecting for Phytoremediation Technology. *Electron. J. Biotechnol.* Vol 6, No 3 2003.
 35. Ghazaryan, K.; Movsesyan, H.; Ghazaryan, N.; Amanda, B. Copper Phytoremediation Potential of Wild Plant Species Growing in the Mine Polluted Areas of Armenia *. *Environ. Pollut.* 2019, 249, 491–501, doi:[10.1016/j.envpol.2019.03.070](https://doi.org/10.1016/j.envpol.2019.03.070).
 36. Brun, L.A.; Maillet, J.; Hinsinger, P.; Pépin, M. Evaluation of Copper Availability to Plants in Copper-Contaminated Vineyard Soils. *Environ. Pollut.* 2001, 111, 293–302, doi:[https://doi.org/10.1016/S0269-7491\(00\)00067-1](https://doi.org/10.1016/S0269-7491(00)00067-1).
 37. Sekabira, K.; Oryem-Origa, H.; Mutumba, G.; Kakudidi, E.; Basamba, T.A. Heavy Metal Phytoremediation by *Commelina Benghalensis* (L) and *Cynodon Dactylon* (L) Growing in Urban Stream Sediments. *Int. J. Plant Physiol. Biochem.* 2011, 3, 133–142.
 38. Brooks, R. Plants That Hyperaccumulate Heavy Metals. In *Plants and the Chemical Elements. Biochemistry, uptake, Tolerance and Toerance*; 2008; pp. 87–105 ISBN 9783527615919.
 39. Kiran; Bharti, R.; Sharma, R. Effect of Heavy Metals: An Overview. *Mater. Today Proc.* 2022, 51, 880–885, doi:<https://doi.org/10.1016/j.matpr.2021.06.278>.
 40. Hassan, M.U.; Chattha, M.U.; Khan, I.; Chattha, M.B.; Aamer, M.; Nawaz, M.; Ali, A.; Khan, M.A.U.; Khan, T.A. Nickel Toxicity in Plants: Reasons, Toxic Effects, Tolerance Mechanisms, and Remediation Possibilities—a Review. *Environ. Sci. Pollut. Res.* 2019, 26, 12673–12688, doi:[10.1007/s11356-019-04892-x](https://doi.org/10.1007/s11356-019-04892-x).
 41. Shahzad, B.; Tanveer, M.; Rehman, A.; Cheema, S.A.; Fahad, S.; Rehman, S.; Sharma, A. Nickel; Whether Toxic or Essential for Plants and Environment - A Review. *Plant Physiol. Biochem.* 2018, 132, 641–651, doi:<https://doi.org/10.1016/j.plaphy.2018.10.014>.
 42. Kumar, B.; Smita, K.; Cumbal Flores, L. Plant Mediated Detoxification of Mercury and Lead. *Arab. J. Chem.* 2017, 10, S2335–S2342, doi:<https://doi.org/10.1016/j.arabjc.2013.08.010>.

43. Ogbonna, C.; Okezie, I.; Okezie, E.; Uzoma, U.; Ugbogu, E.A. Impacts of Quarry Mining Activities on Herbaceous Plant *Ageratum Conyzoides* L. in Ugwuele-Uturu, Abia State, Nigeria. 2020.
44. Mkumbo S; Mwegoha W; Renman G Assessment of the Phytoremediation Potential for Pb, Zn and Cu of Indigenous Plants Growing in a Gold Mining Area in Tanzania. *Int. J. Environ. Sci.* 2012, 2, 2425–2434, doi:10.6088/ijes.00202030123.

UNDER PEER REVIEW