

Common Moths And Their Ecological Importance In Bisra Range, Rourkela Forest Division, (Odisha; India)

ABSTRACT

Rourkela Forest Division presents a sub-tropical climatic condition with high humidity and rainfall after the onset of the monsoon. Lepidopterans are a group of insects that are potent pollinators and protectors of different terrestrial ecosystems. Among all the lepidopterans, moths are cosmopolitan and very important. Moths are mostly nocturnal pollinators and play a vital role in the pollination of different plant species and their seed production. Besides that, they play the important role of both prey and predator in a food chain. They are less known and have always been given less importance, unlike bees and butterflies, because very few studies have been done about moths and their importance in ecosystems. A field survey was done for the documentation of some common moths' diversity in Bisra Range, Rourkela Forest Division, Odisha, India, during 2022–2023. About 20 species of common moths were reported from the study area. This study depicts the ecological importance of moths, how they are inevitable members of the food chain in different types of terrestrial ecosystems, and their interactions with other plant and animal species in the division. *Macrobrosia gigas* and *Phocodermis velutina* larvae are abundant and major defoliators of *Shorea robusta* during the rainy season; *Lymantria mathura* is a major defoliator of *Mangifera indica*, *Shorea robusta*, *Terminalia*, *Mitragyna*, and *Eugenia* spp.; *Xyleutes strix* and *Xyleutes persona* are stem borers; and *Phocodermis velutina* shows a parasitic fungus.

Keywords: Moths, diversity, India, defoliators, borer, parasitic

INTRODUCTION

Among the invertebrates, the phylum Arthropoda is one of the most diversified and efficient phyla in every type of ecosystem, except the oceanic benthic zone (Abrol, 2019; Jamal, 2021). Pollinators are the soul of every ecosystem; they pollinate different plant species, and the majority of pollination occurs by bees and wasps (56%), followed by birds (12%), moths and butterflies (11%), flies (10%), wind (8%), and beetles (3%) (Wychkyus, 2019). Adult moths mainly feed on the nectar and juice of extra-ripe fruit (Sondhi and Sondhi 2016; Lees and Zilli 2019). Moths are nocturnal insects and are mainly responsible for pollinating night-blooming flowers (Singh et al. 2017). Moths are indicators of climate change, primary consumers, potent pollinators, mostly nocturnal, and crucial members of the nutrient cycling

process (Kitching et al., 2000), so they are considered ecologically important insects. Moths are nocturnal insects that belong to the order Lepidoptera. Moths can be easily identified by their activities, antennae structure, and wings. Unlike butterflies, moths have saw-edged antennae and drab-coloured scales on their bodies. Moths are prey for different animals like bats, lizards, amphibians, and some insectivorous birds. There are approximately 160,000 species of moths found worldwide, among them about 12,000 species found in India (Chandra and Nema, 2007), with at least a few thousand undescribed species. Moths are divided into five families: Arctiidae, Noctuidae, Geometridae, Saturniidae, and Sphingidae. Mostly moths are infamous as dull-coloured insects and the 'poor cuisine' of butterflies. Contrary to the fact that some of the moth species have bright body colours with spectacular patterns on their wings, It is well known that invertebrates play the most important role in any type of ecosystem (Cardinal et al., 2006; Bashir, 2019; Shakeel, 2019).

STUDY AREA

Rourkela Forest Division is one of the three forest divisions in Sundargarh District. Other divisions are the Sundargarh Forest Division and the Bonai Forest Division (Pradhan et al., 2023). The division has Reserved Forests, Proposed Reserved Forests, Demarcated Protected Forests, Village Forests, Protected Forests, and DLC Forests. The total forest area was computed to be 1100.43 sq. km, which is about 36.73% of the geographical area of the division. It is also known as the Steel City of Odisha," and it is situated in the northern district of Sundargarh, Odisha, India

METHODOLOGY

A field survey has been conducted in the monsoon months of June to July 2023, i.e., for 2 months. In the present study, preliminary data were collected during a field survey in Rourkela Forest Division, Odisha, India. A total of two surveys were done at 12 different sites, during both the day and night. Each site is approximately 2–3 km away from the next site. During the field survey, a total of 87 moths were photographed through light traps, then released into their wild habitat. Moth specimens were identified by the authors using their morphological characters and with the help of available literature (Sondhi and Sondhi 2016).

RESULTS AND DISCUSSION

The survey revealed a total of 20 common moth species, out of which 18 species are in adult moth form (Plate 1) and the remaining 02 are in caterpillar form (Plate 2), belonging to 17 genera and 10 different families from study areas (Table 1). Among the reported 20 common moth species, 5 are considered serious pests and ecologically important, and 1 moth species shows prey-predator interaction with a parasitic fungus. The common moth diversity of the study areas indirectly portrays the ecological and biodiversity status of the study areas. From the reported 20 species of common moths, 2 species belong to the family Cossidae, 7 species to Erebidae, 1 species to Eupterotidae, 3 species to Geometridae, 2 species to Sphingidae, and 1 species to Limacodidae, Lasiocampidae, Notodontidae, Saturnidae, and Uraniidae, respectively. The study areas are dominated by the family Erebidae, followed by Geometridae, Cossidae, and Sphingidae (Figure 1). *Macrobrochis gigas* and *Phocoderma velutina* larvae are abundant and major defoliators of *Shorea robusta* during the rainy season; *Lymantria mathura* is a major defoliator of *Mangifera indica*, *Shorea robusta*, *Terminalia*, *Mitragyna*, and *Eugenia* spp.; *Xyleutes strix* and *Xyleutes persona* are stem borers; and *Phocoderma velutina* shows a parasitic fungus. It was observed that *Phocoderma velutina* and *Macrobrochis gigas* are very common in the study areas, and their caterpillars are abundant during the monsoon season. The present study revealed that the study area is home to one of the ecologically important fungi, *Cordyceps* spp. Which is a parasitic fungus that grows on different wasps and caterpillars of *Phocoderma velutina*. The predator fungus growth is facilitated by the high humidity, rainfall, and diversified species composition of the study area.

Table 1. Checklist of common moths recorded in the study areas

Family	Scientific name	Species descriptor
Cossidae	<i>Xyleutes strix</i>	Guillou, 1841
	<i>Xyleutes persona</i>	Linnaeus, 1758
Erebidae	<i>Thyas coronata</i>	Fabricius, 1775
	<i>Lymantria semicincta</i>	Walker, 1855
	<i>Lymantria mathura</i>	Moore, 1866
	<i>Lymantria marginata</i>	Walker, 1855
	<i>Eudocima materna</i>	Linnaeus, 1767
	<i>Macrobrochis gigas</i>	Walker, 1854
	<i>Mocis undata</i>	Fabricius, 1775

Eupterotidae	<i>Eupterote hibisci</i>	Fabricius, 1775
Geometridae	<i>Eumelea ludovicta</i>	Guenee, 1858
	<i>Hyperythra lutea</i>	Stoll, 1781
	<i>Scopula ornata</i>	Scopoli, 1763
Limacodidae	<i>Phocoderma velutina</i>	Koller, 1844
Sphingidae	<i>Ambulyx moorei</i>	Moore, 1858
	<i>Theretra silhetensis</i>	Walker, 1856
Saturnidae	<i>Loepa ktinka</i>	Westwood, 1848
Lasiocampidae	<i>Trabala vishnou</i>	Lefebvre, 1827
Notodontidae	<i>Phalera grotei</i>	Moore, 1859
Uraniidae	<i>Acropteris striataria</i>	Clerck, 1764




Figure 1: Moth species diversity in study areas



Plate 1: Some common moths in study areas, 1) *Loepa ktinka* 2) *Lymantria semicincta* 3) *Thyas coronata* 4) *Lymantria marginata* 5) *Theretra silhetensis* 6) *Ambulyx moorei* 7) *Phalera grotei* 8) *Eudocima materna*

Other researchers also been studied on moth's diversity and reported their diversity on different parts of the country. In 2018, Jena et al. reported 30 different species of moths from Gupteswar proposed reserve forest of Eastern Ghats, Koraput, Odisha, India. In 2022 Sanath et al. reported 31 moth species of 12 different families from Barsuan Range, Bonai Forest Division, Odisha, India. In 2019, Singh et al. reported 486 species of moths from different districts of Punjab. Alex et al. also reported moths of 503 different species from Kavvai river basin of Kerala. In 2021 Pawar et al. reported 45 moth species from Panvel, Navi Mumbai, Maharashtra, India. In 2021, Singh et al. reported 17 moth species from Lalwan Community Reserve of Punjab.

ECOLOGICAL IMPORTANCE OF MOTHS

Moths are very important members of different terrestrial ecosystems. Most of these are nocturnal, some are diurnal and crepuscular, and they mostly pollinate night-blooming flowers. It is well known that moths play an important role in different types of food chains as herbivores, creating connections between primary producers and consumers at different trophic levels of the food chain, food web, and food pyramid. In the present study, it was observed that the larvae of *Phocoderma velutina* (Koller, 1844) were infested by *Cordyceps* spp. (a parasitic fungus). *Cordyceps* sp., contains one of the polysaccharides, which is bioactive, exhibits a wide range of bioactivities, and is used as a medicinal herb (Jdrejko et al., 2021). The infestation and parasitic growth of *Cordyceps* spp. inside the body of *P. velutina*, which controls the brain and derives nutrients and then finally leads to larval death,

is a unique parasitic and prey-predator relationship. It shows that besides linking primary producers with different consumers as an herbivore, moths are also linking parasitic organisms with other organisms (Plate 2).




Plate 2: a. Caterpillar of *Phocoderma velutina*, b. Cordyceps infested caterpillar of *P. velutina*.

CONCLUSION

Rourkela Forest Division reveals biodiversity and different ecosystems, almost like Myurbhanj district in Odisha, India. The study areas are dominated by moths of the families Erebiidae, Geometridae, Cossidae, and Sphingidae. During the monsoon season, a large number of caterpillars of *Phocoderma velutina* of the family Limacodidae and *Macrobrochis gigas* of the family Erebiidae feed on their host plants like *Mangifera indica*, *Artocarpus heterophyllus*, and *Shorea robusta*. The study area offers dry deciduous and moist green forests of Sal and mixed-type forest ecosystems, along with their associated plant and animal species. The peculiarity of the study area is that its humid weather promotes the growth of *Cordyceps* sp., which is a parasitic fungus species. The present study highlights the diversified terrestrial ecosystems with some economically and ecologically important species and brings more attention to the conservation of the habitat of the Royal Bengal Tiger.

ACKNOWLEDGEMENT

Authors are thankful to the field staffs and local communities of study areas.

REFERENCES

Alex CJ, Soumya KC, Sajeew TV. A report on the moth (Lepidoptera: Heterocera) diversity of Kavvai River basin in Kerala, India. *Journal of Threatened Taxa*. 2021; 13(2):17753–17779.

- Cardinale, B. J., D. S. Srivastava, J. E. Duffy, J. P. Wright, A. L. Downing, M. Sankaran, and C. Jouseau. 2006. Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature* 443:989–992.
- Chandra, K. and Nema, D.K, “Fauna of Madhya Pradesh (including Chattisgarh)” part- 1. State Fauna series 2007,15:347. Published by Director, Zoological Survey of India. Kolkata.
- Jamal, Z.A.Abou-Shaara, H.F., Qamer, S., Alotaibi, M.A., Khan, K.A., Khan, M.F., Bashir, M.A., Hannan, A., AL-Kahtani, S.N., Taha, E.K.A. and Anjum, S.I., 2021. Future expansion of small hive beetles, *Aethina tumida*, towards North Africa and South Europe based on temperature factors using maximum entropy algorithm. *Journal of King Saud University-Science* 33 (1), 101242. <https://doi.org/10.1016/j.jksus.2020.101242>.
- Jędrejko, K. J., Lazur, J., & Muszyńska, B. (2021). *Cordyceps militaris*: An overview of its chemical constituents in relation to biological activity. In *Foods* (Vol. 10, Issue 11). MDPI. <https://doi.org/10.3390/foods10112634>.
- Jena SK, Singh AP, De K. Diversity of moths (Insecta: Lepidoptera) in the Gupteswar Proposed Reserve Forest of the Eastern Ghat Hill, Koraput, Odisha, India: Preliminary Study. 2018;11(3):11- 17.
- Kitching RL, Orr AG, Thalib L, Mitchell H, Hopkins MS, Graham AW. Moth assemblages as indicators of environmental quality in remnants of upland Australian rain forest. *Journal of Applied Ecology*. 2000; 37:284- 297.
- Kumar N., S., Pradhan, I., Mishra, A. K., & Kumar, S. (2022). Moths Diversity in Barsuan Range, Bonai Forest Division, Odisha, India and their Ecological Importance. *Asian Journal of Biology*, 1–5. <https://doi.org/10.9734/ajob/2022/v15i330238>.
- Lees DC, Zilli A. Moths: Their biology, diversity and evolution. Natural History Museum, London.2019, 208.
- Pawar PR, Supnekar SP, Meshram LN, Pawar NB, Rokade AG. Checklist of moth fauna (Lepidoptera: Heterocera) from Panvel, Navi Mumbai, Maharashtra, West Coast of India. *Uttar Pradesh Journal of Zoology*. 2021;42(13): 11-21.
- Singh AP, Chandra A. Uniyal VP, Adhikari BS. Catalogue of selected insect groups of Lalwan Community Reserve and Ranjit Sagar Conservation Reserve, Punjab, India. *Journal of Threatened Taxa*. 2021;13(3):18020-18029.
- Singh N, Ahmad J, Joshi R. Diversity of Moths (Lepidoptera) with New Faunistic Records from North East Jharkhand India. *Record of Zoological Survey. India*. 2017; 117:326-340.
- Singh N, Pathania PC, Das A, Majumdar A, Sood R. Insecta: Lepidoptera Heterocera (Moths): Fauna of Punjab, State Fauna Series, 23:307-355. (Published by Director, Zool. Surv. of India, Kolkata). 2019.
- Sondhi Y, Sondhi S. A partial checklist of moths (Lepidoptera) of Dehradun, Mussoorie and Devalsari in Garhwal, Uttarakhand, India. *Journal of Threatened Taxa*. 2016; 8: 8756–8776.