

Original Research Article

~~“EFFECTS OF APPLICATION OF DIFFERENT DOSES OF VERMICOMPOST ON THE PHYSICAL PROPERTIES AND PHYSICO-CHEMICAL PROPERTIES OF NON GM COTTON CULTIVARS UNDER RAINFED AGRICULTURAL SYSTEM IN VIDARBHA REGION OF INDIA”.~~

ABSTRACT:

~~The present experiment was conducted at Research Farm Centre for Organic Agriculture Research and Training (COART), Department of Agronomy and department of SSAC, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola during Kharif 2021-22 To assess~~ the effects of application of different doses of vermicompost as a source of organic manure on the physical properties and physico-chemical properties of Non GM cotton cultivars under rainfed agricultural system. The experiment was laid out in Research..... FRBD with eight treatments, each replicated thrice. After organic cotton experimentation, significantly highest reduction in bulk density of soil (1.24 Mg m^{-3}), higher improvement in water holding capacity (58.22 %) and highest (5.67 g kg^{-1}) organic carbon content was recorded in the treatment T_4 where 60 kg N ha^{-1} was supplied through vermicompost to Hirsutum cultivar, Chetna J1 variety. Lowest value of pH (7.57) and EC (0.25 ds m^{-1}) were registered in treatment T_2 where 60 kg N ha^{-1} was supplied through vermicompost to Arboreum cultivar, AVC 14 variety.

Comment [U1]: Lacking an introduction frase

Formatted: Highlight

The abstract is incomplete and not well organized.

Keywords: Vermicompost, Organic cotton, Rainfed Agricultural system

1. INTRODUCTION:

Cotton (*Gossypium spp.* author) is one of the most important fibre and cash crop of India. It plays a key role in Indian agriculture and industrial economy. It is a backbone of textile industries. It is an important cash crop globally known as ‘King of Fiber’. Cotton seed contain 15-20% oil and used as vegetable oil in soap industries. The leftover cake, a by-product of cotton mill is very important feed for livestock. Due to its importance in agriculture and industrial economics it is designated as “White Gold”.

Cotton is one of the most chemically intensive crops in the world. According to the U.S. Department of Agriculture, 84 million pounds of pesticides were applied to the nation's 14.4 million acres of cotton in the year 2000 and more than two billion pounds of fertilizers were spread on those same fields. Seven of the 15 pesticides commonly used on cotton in the United States are listed as "possible," "likely," "probable" or "known" human carcinogens by the Environmental Protection Agency. The intensive farming practices in cotton have shown a range of problematic side-effects on natural resources and farm households (Kooistra&Termorshui-zen,2006). As an attempt to improve the ecological and socio-economics sustainability of cotton production, organic cotton initiatives were started not only in India, but also in other Asian, African and South American countries. Vermicompost is organic fertilizer, it is an excellent source of nutrients for crops and helps to convert organic waste into rich humus. It is derived by composting organic waste by using various species of earthworms. Vermicompost enhance soil fertility physically, chemically, biologically. Vermicompost treated soil has better aeration, porosity, bulk density and water retention. Earthworms create vermicompost, which is a rich source of micro and macronutrients, vitamins, growth hormones, and enzymes. The nutrients present in it are readily water soluble for the uptake of micro flora. (Bhavalkar, 1991). Most of the soils are poor in nitrogen and organic matter. Particularly the soils of the cotton growing areas are low in organic carbon, nitrogen and available phosphorus (Rattan *et al.*1999). Increase in production and productivity can be achieved only through enhanced soil fertility which can be sustained if the nutrients removed from soil are replenished by way of addition. Supplying the entire quantity of nutrients required for cotton through fertilizer may not be possible. It is essential to develop strategies for utilizing all available organic resources and to develop crop management system for sustainable crop production.

Comment [U2]: The introduction need more specific information about some studies already done. And it was just three citation and they are from about 17 years ago. In the last years we have plenty of studying in this subjects.

2. MATERIALS AND METHODS:

Soil:The present investigation was conducted at Research Farm Centre for Organic Agriculture Research and Training (COART), Department of Agronomy and department of SSAC, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola during *Kharif* 2021-22. The topography of the experimental plot was fairly uniform. Soil of experimental field was classified as Vertisols, particularly montmorillonitic, hyperthermic a family of Typic Haplustert. It has smectite clay minerals with swell- shrink properties. The soil

of the experimental field was clay loam in texture and moderately alkaline in reaction. Akola is situated in sub tropical region between 22° 30' 42" N latitude and 77° 02' E longitudes. The altitude of the place is 304.42 m above 30° 3' mean sea level. The climate of Akola is semi arid and characterized by three distinct seasons viz., hot and dry summer from March to May, warm humid rainy season from June to October and mild cold winter from November to February. The experimental rainfed cotton crop was sown on 24th June 2020 and harvested in December 4th. The total rainfall received during the crop growth period was 1084 mm in 54 rainy days.

Vermicompost application: 3 doses of vermicompost have been given to the crop at different rates i.e. 40 kg N/ha and 60 kg N/ha to different cultivars of cotton.

It is important to describe the vermicompost composition.

Analytical procedures: The clod coating method was used to calculate bulk density. Keen Reckzonski boxes were used to measure the soil's ability to hold water. Following a 30-minute equilibration period during which the soil was periodically stirred with water, the pH of the soil was measured in a suspension of soil and water (1:2.5 soil: water) using a glass electrode pH meter. After equilibrating the soil with water and leaving the sample undisturbed until the supernatant was recovered and measured using a conductivity meter, the electrical conductivity was assessed in soil suspension (1:2.5 soil:water). The wet oxidation method was used to determine the soil's organic carbon content of soil.

How was the data analysed? Statistical process?

3. RESULTS:

Influence of varying levels of nitrogen through vermicompost on Physical properties of the soil like bulk density and water holding capacity:

3.1) Bulk density:

~~Data on soil bulk density are shown in Table 2 for distinct organic treatments.~~ The data range from 1.24 to 1.32 Mg m⁻³, however the influence of vermicompost on soil bulk density was not statistically significant (Table 2). It was discovered that adding more organic manures, such as vermicompost, causes the bulk density to gradually decrease. Lowest bulk density 1.24 Mg m⁻³ was recorded in the treatment T₄ received 60 kg N/ha through vermicompost to hirsutum cultivar, Chetna J1 variety. The highest (1.32 Mg m⁻³) bulk density was recorded in the treatment T₇ where 40 kg N/ha was applied through vermicompost to hirsutum cultivar, GHV 8023 variety Table ?:-

Formatted: Normal, Space After: 0 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers, Tab stops: Not at 0.95"

Formatted: Font: Calibri, 11 pt, Bold

Table-1. Initial soil properties[Improve this title](#)

S.No.	Particulars	Value	Method adopted
1	Bulk density (Mg m^{-3})	1.26	Clod coating method Author
2	Water holding capacity (%)	50.3	Keen box method Author
3	pH (1:2.5)	7.6	Glass electrode pH meter
4	Electrical conductivity (ds m^{-1})	0.3	Electrical conductivity bridge Author
5	Organic Carbon (g kg^{-1})	4.9	Walkley and Black rapid titration method Author

Table 2. Effect of varying levels of nitrogen through vermicompost on bulk density

Treatment	Bulk density(Mg m^{-3})
T ₁ - AV1N1- Arboreum cultivar + 40 kg N ha ⁻¹	1.29
T ₂ - AV1N2- Arboreum cultivar + 60 kg N ha ⁻¹	1.25
T ₃ - HV1N1- Hirsutum cultivar + 40 kg N ha ⁻¹	1.27
T ₄ -HV1N2- Hirsutum cultivar + 60 kg N ha ⁻¹	1.24
T ₅ -AV2N1- Arboreum cultivar + 40 kg N ha ⁻¹	1.30
T ₆ -AV2N2- Arboreum cultivar + 60 kg N ha ⁻¹	1.26
T ₇ -HV2N1- Hirsutum cultivar + 40 kg N ha ⁻¹	1.32
T ₈ -HV2N2- Hirsutum cultivar + 60 kg N ha ⁻¹	1.26
SE (m) ±	0.008
CD at 5%	NS
Initial value	1.26

NOTE: 1. HV1-Chetna J1
 2. HV2-GHV 8023
 3. AV1-AVC 14
 4. AV2-GAV 8011

Table 3. Effect of varying levels of nitrogen through vermicompost on Water holding capacity:

Treatment	Water holding capacity (%)
T ₁ -AV1N1- Arboreum cultivar + 40 kg N ha ⁻¹	50.93

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted: Font: Not Bold, Highlight

Formatted: Justified, Indent: Left: 0",
 Hanging: 0.75", Space Before: 3 pt, After:
 3 pt, Tab stops: 1", Left

T ₂ -AV1N2- Arboreum cultivar + 60 kg N ha ⁻¹	56.37
T ₃ -HV1N1- Hirsutum cultivar + 40 kg N ha ⁻¹	51.77
T ₄ -HV1N2- Hirsutum cultivar + 60 kg N ha ⁻¹	58.22
T ₅ -AV2N1- Arboreum cultivar + 40 kg N ha ⁻¹	48.81
T ₆ -AV2N2- Arboreum cultivar + 60 kg N ha ⁻¹	54.11
T ₇ -HV2N1- Hirsutum cultivar + 40 kg N ha ⁻¹	47.89
T ₈ -HV2N2- Hirsutum cultivar + 60 kg N ha ⁻¹	52.60
SE (m) ±	1.151
CD at 5%	NS
Initial value	50.3

3.2) Water Holding Capacity:

Influence of vermicompost under different organic treatments on water holding capacity of soils was non significant ~~it was presented in~~ (Table 3) and it was ranged from 47.89% to 58.22 %. Lowest water holding capacity was observed in hirsutum cultivar, GHV 8023 variety T₇ where 40 kg N/ha was applied through vermicompost and highest water holding capacity was recorded in hirsutum cultivar Chetna J1 variety which is supplemented with 60kg N/ha.

Though there was improvement in the physical properties of the soil by the application of vermicompost, its effects are not seen immediately. It helps to boost the soil physical properties in the long run.

Table 3. Effect of varying levels of nitrogen through vermicompost on Water holding capacity:

<u>Treatment</u>	<u>Water holding capacity (%)</u>
T ₁ - AV1N1- Arboreum cultivar + 40 kg N ha ⁻¹	50.93
T ₂ - AV1N2- Arboreum cultivar + 60 kg N ha ⁻¹	56.37
T ₃ - HV1N1- Hirsutum cultivar + 40 kg N ha ⁻¹	51.77
T ₄ -HV1N2- Hirsutum cultivar + 60 kg N ha ⁻¹	58.22
T ₅ -AV2N1- Arboreum cultivar + 40 kg N ha ⁻¹	48.81
T ₆ -AV2N2- Arboreum cultivar + 60 kg N ha ⁻¹	54.11
T ₇ -HV2N1- Hirsutum cultivar + 40 kg N ha ⁻¹	47.89
T ₈ -HV2N2- Hirsutum cultivar + 60 kg N ha ⁻¹	52.60

<u>SE (m) ±</u>	<u>1.151</u>
<u>CD at 5%</u>	<u>NS</u>
<u>Initial value</u>	<u>50.3</u>

Influence of varying levels of nitrogen through vermicompost on Soil pH, Electrical conductivity and Organic carbon content:

3.3) Soil pH

Table 4 displays the findings relating the pH of the soil at cotton harvest. The pH range of the test soil was 7.57 to 7.68. The T₇ recorded a pH value as its highest reading (7.68) where 40kg N ha⁻¹ given through vermicompost to Hirsutum cultivar, GHV 8023 variety. however lowest value of pH 7.57 was recorded in treatment T₂ where 60 kg N/ha was supplied to Arboreum cultivar, AVC 14 variety.

Table 4. Effect of varying levels of nitrogen through vermicompost on pH and Electrical conductivity

Formatted: Highlight

Treatment	pH (1:2.5)	EC (dS m⁻¹)
T ₁ - AV1N1- Arboreum cultivar + 40 kg N ha ⁻¹	7.64	0.26
T ₂ - AV1N2- Arboreum cultivar + 60 kg N ha ⁻¹	7.57	0.25
T ₃ - HV1N1- Hirsutum cultivar + 40 kg N ha ⁻¹	7.66	0.3
T ₄ -HV1N2- Hirsutum cultivar + 60 kg N ha ⁻¹	7.58	0.26
T ₅ -AV2N1- Arboreum cultivar + 40 kg N ha ⁻¹	7.67	0.3
T ₆ -AV2N2- Arboreum cultivar + 60 kg N ha ⁻¹	7.58	0.27
T ₇ -HV2N1- Hirsutum cultivar + 40 kg N ha ⁻¹	7.68	0.31
T ₈ -HV2N2- Hirsutum cultivar + 60 kg N ha ⁻¹	7.60	0.28
SE (m)±	0.004	0.005
CD at 5%	0.012	NS
Initial value	7.6	0.3

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

3.4) Electrical Conductivity:

The results regarding electrical conductivity of soil after harvest of cotton was presented in Table 4. The values of electrical conductivity of soil were statistically non significant under various organic treatments.

The Electrical conductivity of the experimental soil ranged from (0.25 to 0.31) dSm⁻¹. The highest Electrical conductivity value (0.31dSm⁻¹) was recorded in the treatment T₇ received 40kg N ha⁻¹ through vermicompost to Hirsutum cultivar, GHV 8023 variety. Electrical conductivity was found to be decreased as the doses of organic nutrient sources increases. The lowest value of Electrical conductivity (0.25 dSm⁻¹) was observed in the treatment T₂ received Vermicompost @60kg N ha⁻¹ toArboreum, AVC 14 variety.

3.5) Organic carbon:

The results pertaining to organic carbon in soil at harvest of cotton is presented in Table 5. The results indicated the impact of vermicompost on build up of soil organic carbon content. Significantly superior organic carbon (5.67 g kg⁻¹) build up in soil was noticed in treatment T₄ where 60kg N ha⁻¹ supplied through vermicompost to Hirsutam cultivar, Chetna J1 Variety. However the lowest organic carbon (5.17 g kg⁻¹) was registered with treatment T₁ where 40kg N ha⁻¹ was applied to Arboreum cultivar, AVC 14 variety.

Table 5. Effect of varying levels of nitrogen through vermicompost on organic carbon:

Treatment	Organic carbon (g kg ⁻¹)
T ₁ - AV1N1- Arboreum cultivar + 40 kg N ha ⁻¹	5.17
T ₂ - AV1N2- Arboreum cultivar + 60 kg N ha ⁻¹	5.41
T ₃ - HV1N1- Hirsutum cultivar + 40 kg N ha ⁻¹	5.29
T ₄ -HV1N2- Hirsutum cultivar + 60 kg N ha ⁻¹	5.67
T ₅ -AV2N1- Arboreum cultivar + 40 kg N ha ⁻¹	5.26
T ₆ -AV2N2- Arboreum cultivar + 60 kg N ha ⁻¹	5.40
T ₇ -HV2N1- Hirsutum cultivar + 40 kg N ha ⁻¹	5.20
T ₈ -HV2N2- Hirsutum cultivar + 60 kg N ha ⁻¹	5.60
SE (m)±	0.013
CD at 5%	0.039
Initial value	4.9

ALL THE TABLE (From 2 to 4) YOU CAN ORGANIZE THE DATA FOR JUST ONE.

4. DISCUSSION:

Influence of varying levels of nitrogen through vermicompost on Physical properties of the soil like bulk density and water holding capacity:

4.1) Bulk density

It is clear from the experiment that adding organic manures helps to lower soil weight by maintaining the right soil:moisture:air ratio, which ultimately results in a decrease in the bulk density of soils. The increase in organic manure application rates is directly correlated with the improvement in soil bulk density. Similar results were observed by Marathe and Bharambe (2005). Continuous application of FYM @ 10 t ha may lead to increased aggregation of 3 -1 1 soil particles followed by vermicompost @ 5 t ha-1 over control. In contrast, Laharia et al. (2013) found that applying 100% RDN by vermicompost + jeevamrut resulted in the soil having the lowest bulk density. Treatments using thoroughly decomposed organic materials, such as vermicompost, often shown a considerable reduction in bulk density of soil. Katkar et al. (2007) found a significant drop in bulk density with the application of FYM @ 10 t ha-1 above control. Bellakki and Badanur (1997), Surekha and Rao (2009), and Vora et al. (2015) reported similar findings reporting a significant decrease in bulk density of Vertisols with the addition of sunhemp and FYM above control.

Comment [U3]: Looking for recent publications

[Need more discussion](#)

4.2) Water Holding Capacity:

Vermicompost enhance soil structure and increase its water holding capacity. This is because vermicompost contains humus and organic matter, which improve soil aggregation, allowing it to retain more water. Additionally, the presence of beneficial microorganisms in vermicompost enhances soil porosity, allowing water to infiltrate and be stored more effectively. As a result, the application of vermicompost can lead to better water retention in the soil, benefiting plant growth and overall soil health. [Author](#)

According to Jadhav et al. (1993), the soil's ability to store more water was improved in the three plots that got vermicompost at a rate of 5 tonnes per hectare. Similar to this, Mastiholi (1994) reported that applying 2 tonnes of vermicompost per hectare to the soil increased its moisture content compared to applying 1 tonnes. According to Patil (1998), FYM incorporation at 2.5 tonnes per ha and vermicompost application at 1.0 tonnes per ha improved the soil moisture content compared to simply fertilizer treatment in the vertisols of Bijapur. Bhatia and Shukla (1982) found that the use of FYM, either alone or in conjunction with fertilizers, greatly enhanced the water holding capacity and moisture retention at field capacity.

Comment [U4]: Try to find a recent references

[And what? More discussion](#)

Influence of varying levels of nitrogen through vermicompost on pH, Electrical conductivity and organic carbon content:

4.3) Soil pH

pH was found to be decreased as the doses of organic nutrient increases. Various treatments have showed significant influence of H^+ concentration in soil and it was reduced towards neutrality as increment in doses of vermicompost. The pH decrease upon application of vermicompost is likely due to the presence of organic acids in the compost. Vermicompost contains various organic compounds, including humic and fulvic acids, which are acidic in nature. When vermicompost is added to the soil, these organic acids gradually get released into the soil, causing a slight decrease in pH.

Moreover, as the organic matter in the vermicompost decomposes, it undergoes microbial activity, and this process can also produce some acidic byproducts. As a result, the overall effect of vermicompost application is a slight reduction in soil pH. While there is a pH decrease upon initial application, over time, the buffering capacity of the soil may help stabilize the pH, resulting in a more balanced soil pH in the long run. The similar results were reported by Katkaret *et al.* (2007) revealed that pH was decreased with application of FYM (10 t ha⁻¹). The findings are in line with observations recorded by Pawar *et al.* (2007), Halemaniet *al.* (2004) and Oja *et al.* (2014).

4.4) Electrical Conductivity

The electrical conductivity (EC) of the soil can decrease upon the application of vermicompost due to its low salt content. Vermicompost is a type of organic matter that has undergone decomposition through earthworms. Unlike some synthetic fertilizers or mineral-rich composts, vermicompost typically has a lower concentration of soluble salts, which are the primary contributors to the soil's electrical conductivity. When vermicompost is added to the soil, it introduces more organic matter and nutrients, which can enhance soil structure and support microbial activity. As a result, the organic matter can help reduce the leaching of salts from the soil, leading to a decrease in electrical conductivity. Similar results were reported by Halemaniet *al.* (2004), noted that organics alone and their combination had no significant influence on electrical conductivity of soil. Results are also in ~~conformity~~ conformity with findings of Badanure *et al.* (1990), Nehra & Hooda (2002) and Katkaret *et al.* (2002). Similar results were reported by Halemaniet *al.* (2004), noted that organics alone and their combination had no significant influence on electrical conductivity

of soil. Results are also in ~~conformity~~conformity with findings of Badanure *et al.* (1990), Nehra&Hooda (2002) and Katkare *et al.* (2002).

4.5) Organic carbon

Utilization of organic inputs like vermicompost, as source of plant nutrient to cotton reported significant improvement in the organic carbon content in soil. The increase in organic carbon in soil might be due to direct addition of organic matter through vermicompost and increasing root biomass of plant. Organic matter in vermicompost consists of partially decomposed plant and animal materials, which are rich in carbon-based compounds. When vermicompost is added to the soil, its organic matter is incorporated into the soil matrix. This organic matter serves as a food source for soil microbes, promoting microbial activity and decomposition. As the microorganisms break down the organic matter, they release carbon compounds into the soil, thus increasing the soil's organic carbon content. Higher organic carbon levels in the soil offer several benefits, such as improved soil structure, water holding capacity, nutrient retention, and enhanced microbial diversity. Additionally, organic carbon is crucial for the formation and stability of soil aggregates, which contributes to better soil aeration and root growth. Tadesse *et al.* (2013) reported that, organic carbon content of soil increased due to higher (15 t ha⁻¹) FYM application. Similar results were also reported by Badole *et al.* (2000), Sonune *et al.* (2003) and Oja *et al.* (2014).

Conclusion:

Based on the observations taken, it is concluded that the utilization of vermicompost as a nutrient source significantly improves physical and physico- chemical properties of soil. Application of 60 kg N ha⁻¹ through vermicompost to Non GM cultivars of cotton is found to be beneficial and increased physical and physico- chemical properties of soil than application of 40 kg N ha⁻¹ through vermicompost to Non GM cultivars of cotton. Hirsutum Cultivar, Chetana J1 variety responded well when compared to arboretum cultivars for organically grown Cotton under rainfed agricultural system.

References:

Improve the references with recent one.

Badanur, V. P., C. M. Poleshi and B. K. Naik, 1990. Effect of organic matter on crop yield and physical and chemical properties of Vertisol. *J. Indian Soc. Soil Sci.* 38 (3): 426-429.

Comment [U5]: It is important to answer the aims in the conclusion and making clear this.

- Badole, S. B. and S. D. More, 2000. Organic carbon status as influenced by organic and inorganic nutrient sources in Vertisol. *J. Maharashtra Agric. Univ.* 25 (2): 220-222.
- Bellakki, M. S. and V. P. Badanur, 1997. Long term effect of integrated nutrient management on properties of a Vertisol under dryland agriculture. *J. Indian Soc. Soil Sci.* 45(3) : 438-442.
- Bhatia, K.S., and K.K. Shukla, 1982, Effect of continuous application of fertilizers and manure on some physical properties of eroded alluvial soil. *Journal of the Indian Society of Soil Science*, 30: 33-36.
- Bhavalakar, U., 1991. Vermiculture biotechnology for LEISA - seminar on low external input sustainable agriculture. Amsterdam, Netherlands, pp: 16.
- Halemani, H. L., S.S. Hallikeri, S. S. Nooli, R. A. Nandagavi and H. S. Harish Kumar, 2004. Effect of organics on cotton productivity and physico-chemical properties of soil. In: International symposium on "Strategies for sustainable cotton production-A global vision". 2. Crop production, 23-25., VAS, Dharwad: 174-180.
- Jadhav, S. B., M.B. Jadhav, V.A. Joshi and P. B. Jagatap, 1993, Organic farming in the light of reduction in use of chemical fertilizers. Proceedings of 43rd Annual Deccan Sugar Technology Association, Pune, Part I, pp. SA53-SA65.
- Katkar, R.N., P.D. Bhalerao, P.U. Ghatol and B. R. Patil, 2007. Influence of organics and inorganics on soil fertility, seed cotton yield and fibre quality in Vertisol. *PKV Res. J.*, 31(2): 216-222.
- Kooistra, K.J., A.J. Termorshuizen, and R. Pyburn, 2006. *The sustainability of cotton: Consequences for man and environment* (No. 223). Science Shop Wageningen UR.
- Laharia, G. S., D. U. Patil and P. R. Damre, 2013. Effect of organic sources on soil fertility, nutrient uptake and yield of soybean. *Crop Res.* 45 (1, 2 & 3):155-159.
- Marathe, R. A. and P. R. Bharambe, 2005. Physical characteristics of Vertisols as influenced by INM system under sweet orange. *J. Indian Soc. Soil Sci.* 41 (1): 52-57.
- Mastiholi, A. B., 1994, Response of rabi sorghum (*Sorghum bicolor* L.) to biofertilizer and In situ moisture conservation practices in deep black soil. *M. Sc. (Agri.) Thesis*, University of Agricultural Sciences, Dharwad.
- Nehra, A.S. and I. S. Hooda, 2002. Influence of integrated use of organic manures and inorganic fertilizers on wheat yield and soil properties. *Res. on Crops*, 3(1):11-16.
- Ojha R. B., S. C. Shah, K. R. Pande and D. D. Dhakal, 2014. Residual Effects of Farm Yard Manure on Soil Properties in Spring Season, Chitwan, Nepal. *International J. of Sci. Res. in Agric. Sci.*, 1(8), pp. 165-171.
- Patil, S. L., 1998, Response of rabi sorghum [*Sorghum bicolor* (L.) Moench] to tillage, moisture conservation practices, organics and nitrogen in Vertisols of semi-arid tropics. *Ph. D. Thesis*, University of Agricultural Sciences, Dharwad.

- Pawar, R. B. and C. V. Patil, 2007. Effect of vermicompost and fertilizer levels on soil properties, yield and uptake of nutrients by maize. *J. Maharashtra agric. Univ.*, 32(1): 11-14.
- Rattan, R.K., S. Neelan and S.P. Datta, 1999. Micronutrient depletion in Indian soils: extent, causes and remedies. *Fertil. News.*, 44(2):35-50.
- Sonune, B. A., K. B. Tayade, V.V. Gabhane and R. B. Puranik, 2003. Long- term effect of manuring and fertilization on fertility and crop productivity of Vertisols under sorghum- wheat sequence. *Crop Res.* 25(3):460-467.
- Surekha. K and K. V. Rao, 2009. Direct and residual effects of organic sources on rice productivity and soil quality of Vertisol. *J. Indian Soc. Soil. Sci.*, 57 (1): 53-57.
- Tadesse, T., N.Dechassa, W. Bayu and S. Gebeyehu, 2013. Effects of Farmyard Manure and Inorganic Fertilizer Application on Soil Physico-Chemical Properties and Nutrient Balance in Rain-Fed Lowland Rice Ecosystem. *Am.J. of Plant Sci.*, Vol 4, 309-316.
- Vora, V. D., K.D.Rakholiya, K.V.Rupapara, G.S. Sutaria and K.N. Akbari, 2015. Effect of Integrated Nutrient Management on Bt Cotton and Post Harvest Soil Fertility under Dry Farming Agric. *Asian J. of Agric. Res.*, 9 (6): 350-356.