

The influence of Slope Gradient on Termite Assemblages and some Soil Physico-chemical Properties in Koudi Secondary Forest Patches of the Central Region of Cameroon

ABSTRACT

Soil is the Cameroon's primary economic driver and an indispensable natural resource for agriculture. Various factors including physical, chemical and biological directly affect agricultural production by enhancing soil fertility. The influence of relief on termites' diversity was extensively studied. However, the effect of the slope gradient on termite diversity and functional groups remains less studied. This study We evaluated termite assemblages in relation to some edaphic parameters, with emphasis on gradual changes in soil slope. The study was conducted in Koudi, in the semi-deciduous forest zone of Southern Cameroon. The study We aimed to evaluate (1) the effect of slope gradient on termite diversity and some soil physico-chemical factors; (2) the influence of some physico-chemical soil properties on termites. To achieve these, three slope gradients: gentle slope: A [0 - 15°]; moderate slope: B [15.1°- 31°]; and steeper slope: C [x > 31°] were chosen. The experimental design was composed of 66 rectangular quadrants of 5m x 2m in order to determine the presence or absence of termites. Two soil samples were taken per slope gradient for the determination of physico-chemical parameters. Termite community was characterized using species richness, Shannon diversity, Pielou Evenness and Simpson dominance indexes. Data obtained were subjected to Spearman correlation test and ANOVA followed by the Tukey HSD test at a threshold p = 0.05. Our findings suggested thirty-five (35) termite species belonging to eighteen (18) genera, six (6) subfamilies that were sampled and identified. Termites' diversity and species richness decreased as slope gradient increased; slope gradient displayed a significant detrimental correlation with sand (%) and total organic (g/kg) content, with a positive significant correlation with clay (%) content, according to Spearman's test. It appeared that variations of slope at soil surface have been identified as one of the factors controlling nutrients and termite diversity.

Key words: Slope, Termites, Physico-chemical parameters, Semi-deciduous Forest

Introduction

Soil is the nation's primary economic driver and a valuable natural resource for agriculture. Both Physico-chemical and biological factors influence soil fertility, directly affecting agricultural production. Topography through slope aspect is one of the soil-forming factors that influences soil characteristics and controls soil erosion processes by redistributing soil particles and soil organic matter (OM) [1,2]. Soil loss is typically expected to increase with increasing slope gradient due to increased surface runoff velocity and decreased infiltration rate [3,4]. Several studies showed differences in some soil characteristics may be related to slope gradient [5]. With differences in solar radiation

Comment [IL1]: Abstract needs to be recasted to capture the missed methods and some key results

Comment [IL2]: This contradict the region stated in the title. Please clarify.

Comment [IL3]: Methods for the determination of these parameters were not captured in this abstract.

Comment [IL4]: Values of these parameters were not stated.

Comment [IL5]: See above comment. The procedure for determining soil textural classes was not captured

between northern and southern slopes, this variation in slope gradient at some scales can result in distinct climates and raise temperatures and evapotranspiration in one hemisphere relative to another [6,7]. Consequently, topographic characteristics significantly affect plant functional traits and diversity. It is possible that the observed changes in plant communities in response to slope also had cascading impacts on nearby wildlife [8].The accompanying fauna may have been affected in a cascading pattern by the observed changes in plant communities in response to slope [8].

One of the organisms that alter the soil qualities is the termite (Order: Blattaria, Infraorder: Isoptera). According to [9]there are 3106 extant or fossil termite species in the world, divided into 12 families, with more than 75% of all known species belonging to the Termitidae family[10,11].The humid tropical region has the highestdiversitydue to its physical complexity and small-scaleheterogeneity[12]. Termites play a crucial ecological role by influencing pedogenesis and the physico-chemical properties and functions of soils. They are directly or indirectlyinvolved, in a variety of soil processes, such as organic matter dynamics, aeration, water transport, and water holding capacity [13]. They are soil engineersas they participatein organic matter breakdown, concentration, storage and redistribution of mineral and organic elements. Physicochemical changes in the soil can affect termite assemblages. Numerous factors can influencesoilqualities. Based on functional or taxonomic groups, previous studies have investigated the distribution of termites at different spatial scales [14, 15]. However, determining the spatial scale is crucial for assessingthe impact of environmental factors on termite communities; spatial scale has been shown to assess biotic processes, such as dispersal, intraand interspecific interactions [16]. In tropical forests, spatial heterogeneity affects the distribution of soil-level nutrients and environmental attributes that influencespecies activity. However, it is still unclear how environmental factors influence the spatial distribution of termite communities.

Topography is likelyto be one of the most significant factors. Depending on the slope's gradient, drainage and runoff encourage erosion andalterthe physico-chemical characteristics of the soil by moving soil nutrients from steep slopes to gentle slopes [1].As a result, nutrients are unevenly distributed in the soil depending on the landform.The relief of Cameroon's semi-deciduous forests is complex, with moderate to steeply sloping hills alternating with undulating land interspersed with plains having relatively flat topography [1, 17].This has an impact on the diversity and distribution of nutrients in the soil. Fewstudies have beenconducted to evaluate the effect of slope gradient on termite diversity. So, the present study weaimed toevaluate--the influence of slope gradient on termite assemblages and soil physicochemical properties in Koudi forest patches of central region of Cameroon

- (1) ~~The effect of slope gradient on termite diversity;~~
- (2) ~~The effect of slope gradient on some soil physico-chemical properties;~~
- (3) ~~The relationship between some soil physico-chemical properties and termites.~~

- ##Material and Methods
- ##1 Study Site

Formatted: Indent: Left: 0", First line: 0.49", Space After: 8 pt, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

The study was conducted in a semi-deciduous forest zone with a bimodal rainfall regime in Cameroon at the Koudi locality (3°95'22.06"N; 11°26'40.9 East; altitude: 681 ± 3 m) in the Centre Region, Department of the Lekié and Lobo District from 14 April to 31 May, 2021 (Figure 1). This area is characterized by the Guinean equatorial climate type with alternating four seasons, including two rainy seasons (mid-March to mid-June; mid-September to mid-December) and two dry seasons (mid-June to mid-August; mid-December to mid-March) [17,18]. The average annual rainfall is 1800 mm, with a difference of about 280 mm between the driest and the wettest months. The average annual temperature is 23.6°C, with a thermal variation of 2.1°C [17]. The Centre Region is part of the Southern Plateau of Cameroon. The average altitude of this large plateau is 600-700 m; the predominant vegetation is dense tropical forest, which gradually turns into savannah in the northern part; this is a point of severe ecosystem degradation. The vegetation in the study area is of the mosaic type, with remnants of primary forest on the hilltops and secondary forest and fallow predominating on the plains. The vegetation is characterized by the richness of the epiphytic flora caused by the extremely high humidity of these environments, being the most distinctive physiognomic link. Species such as *Allanblackia gabonensis*, *Cola verticillata*, *Beilschmiedia obscura*, *Leonardoxa africana*, *Garcinia polyantha*, and *Myrianthus serratus* predominated others [19]. The main cash and consumption crops are *Theobroma cacao*, *Manihot esculenta*, *Arachis hypogaea*, *Zea mays*. Soils are clayey in consistency and reddish in colour with an irregular slope [18]. A secondary forest plot over 40 years old was selected in this forest.

2 Methods

Experimental Set-up and Sampling

The experimental set-up consisted of a 5 hectare forest plot in which 66 non-contiguous 5 m x 2 m quadrats were randomly delimited. These quadrats were at least 10 m apart to cover the entire topographic profile of the study area. The slope was measured using a digital inclinometer that previously calibrated on a horizontal plane; termite sampling was carried out in each quadrat according to [20] approach. Subsequently, 12 soil samples (20 cm x 20 cm x 10 cm) were collected using an entomological trowel, placed on a tray, and thoroughly studied for 30 minutes by two collectors. Termite activity and the presence or absence of both young and old epigeal nests were documented. Soldiers and workers from the various termite castes were collected and preserved in tubes containing 80% alcohol. These samples were sent to the Entomology laboratory of the IRAD (Institut de Recherche Agricole pour le Développement) for identification using the keys of [21], [10] and [22].

Two soil samples were taken per slope gradient using 100 cm³ cylindrical rings which had previously been weighed for the determination of physico-chemical parameters of the soil. The soil (0–10 cm) was collected after litter removal. Chemical and textural analyses were then carried out. Soil samples were taken from each slope gradient to determine soil particle size and certain chemical parameters. Soil samples were then sent to the soil laboratory of Dschang's University, where all samples were crushed and sieved to a size of 2 mm and 0.5 mm after air drying. Bulk density, organic carbon, total nitrogen, assimilable phosphorus, water pH, exchangeable cations, carbon/nitrogen ratio, and particle size were measured.

Comment [IL6]: See above comment on region and provide clarification.

Bulk density: Fresh 100 cm³ soil samples were weighed, oven-dried at 105°C, and weighed again to estimate water content and bulk density[citation needed].

Determination of Soil Texture

Soil texture:The Bouyoucos hydrometer method was used to analyzethe soil textural fractions (sand, silt, and clay), in which 15 g of air-dried soil sieved to 2 mm were weighed into 500 ml beakers and subjected to organic matter removal treatment with H₂O₂, followed by dispersion of the soils with sodium hexametaphosphate [23].The resulting mixtures were agitated on a mechanical shaker for 3 hours. The suspensions were then transferred to sedimentation apparatusand built up to a capacity of 1000 ml with distilled deionized water in sedimentation measuring cylinders. To suspend the particles, the mixtures were extensively stirred with a mechanical rotator. A hydrometer was then used to take readings after 40 seconds (first reading, R1) and two hours (second reading, R2) respectively[24].

Determination of Soil Physico-chemical Parameters

Soil pH was determined in a water suspension with a soil/ water ratio of 2:5. Exchangeable bases (Ca²⁺, Mg²⁺, K⁺ and Na⁺) were extracted using 50 ml of ammonium acetate (1.0 M NH₄OAc) solution buffered at pH 7. Using complexometric titration and a flame photometer, the concentrations of magnesium (Mg²⁺), calcium (Ca²⁺), potassium (K⁺), and sodium (Na⁺) were determined in the extract, respectively. After that, phosphorus was detected using a Malachite green colourimetric method. Organic carbon was determined by chromic acid digestion and spectrophotometry. Total Nitrogen was determined using the Kjeldahl method for digestion as well as ammonium electrode determination [25, 26].

II- 2 Data Analysis

Assessment of sampling effort and termites community diversity

To assess the sampling effort, the diversity obtained from non-parametric biodiversity estimators using rare species in the community was compared to the theoretical diversity of the site to predict the number of missing species [27]were used to compare the obtained diversity to the theoretical diversity of the site. Chao1 is a non-parametric species richness estimator based on abundanceand correlated with other diversity measures. Jackknife1 is non-parametric estimator of species richness based on abundance and incidence. Bootstrap is a simulation of standard deviation by randomly generating samples to estimate the potential number of observed and theoretical termite species in each gradient slope.We also calculated the Simpson's dominance index (D), as well as the Pielou evenness index (J) and the Shannon-Wiener diversity index (H') per slope gradient.Species rarefaction curves were plotted using species accumulation as a function of sample number.All these analyses were performed using Primer (Plymouth Routines In Multivariate Ecological Research) software version 6.0 and Past 4.02.

The numbers of termites in the different quadrats were subjected to a logarithmic transformation of the Log(N+1) type to normalize them, then subjected to a simple analysis of variance (ANOVA) followed by the Tukey HSD test for pairwise comparison.

Formatted: Indent: First line: 0"

Comment [IL7]: For any heading, please capitalise each word.

Relationship between soil property and diversity parameters:A Canonical Correspondence Analysis (CCA) was applied according to[28]to analyze the variation of slope gradient versustermites’ variables and soil physicochemical parameters. Non-metric multidimensional scaling analysis (NMDS) was applied to show similarities in the composition of termiteassemblages among slope gradients using the Bray–Curtis similarityindex.

Spearman's correlation was used to find the relationship between environmental parameters, species richness and termite abundance. All statistical analyses and tests were carried out using Past 4.0.2 and STATISTICA 6.0 software, and the results were appreciated at a 5% confidence interval.

III- Results

1Evaluation of sampling effort

A total of 1404 termites from 18 genders and 35 species were gathered from four different slope gradients. Apicotermitinae (nine species), Cubitermitinae (seven species), Microcerotermitinae (one species), Macrotermitinae (two species), Nasutitermitinae (one species), and Termitinae (15 species) all belonged to the same family, Termitidae.In terms ofthe functional groups, soil feeding (94.80%), wood-feeding (1.64%) and fungus growers (3.56%) constituted the total sampledpopulation.

Comment [IL8]: Please use table to present these results.

In terms of sampling effort, independent of slope, estimated species richness was substantially higher than measured ones using the Chao, Jackknife1, and bootstrap estimators. While the Chao1 estimate for the species was 38.63, the Jackknife estimate was 42.87, and the bootstrap estimate was 38.80. Chao1 estimations were more conservative, with 10.37 percent more species estimated than observed (Figure 2). It should be noticed that these three curves continue to increase, implying that certain species were not sampled.

2Effect of slope gradient on termite’s community diversity

The diversity of termites may indicate the effect of the slope on the ground. As shown in Figure 3 and Table 1, Anova followed by Tukey's tests revealed that the number of species decreasedas the level of the slope gradient increased, but not substantially (2ddl; F (2, 63) = 1,885; p=0, 16). Furthermore, the Shannon-wiener index (H) decreased as the level of the slope gradients increased. On the contrary, Simpson's dominance index (D) gradually increased. All of the species studied were divided into three functional groups: fungus growers (Gr II_r), wood-feeding (Gr II), and soil-feeding (Gr III & IV). The results collected reveal that fungus growers (2ddl; F (2, 63) = 0.07; p=0.93), wood-feeding (2ddl; F (2, 63) = 0.015; p=0.85), and soil-feeding termites (2ddl; F (2, 63) = 0.063; p=0.93), did not differ between slope gradient.

Comment [IL9]: This table was not presented in this manuscript. Please provide it.

3 Effect of slope gradient on soil properties and termite distribution

Table 2 depicts the effect of slope gradient on the physical and chemical properties of the soil. According to the ANOVA, the pH-water and the values of C/N ratios decreased somewhat, although the water content increased slightly but not considerably (p>0.05). Concerning the exchangeable cations, a higher magnesium concentration was observed in the steeper slopes (C) than in the gentle and moderate slopes (A and B). Potassium (K) seemed to be influenced by the slope gradient with a lower concentration in the steeper slope (C). In the same way, exchangeable sodium (Na) and calcium (Ca) showed a difference in slope gradient with lower concentrations in the steeper slope (C). All these differences were not statistically different. In terms of soil texture, sand and silt content decreased while clay content increased with increasing slope gradient, with no significant difference at each level of slope gradient (p > 0.05); however, statistical differences in bulk density were observed, with moderate slope (B)being higher than gentle (A) and steeper (C) slopes (2ddl; F(2,33)=1.37; p=0.009).

CCA analysis was used to investigate the impact of the slope aspect on termite assemblages and soil physico-chemical parameters (Figure 4). According to this analysis, CCA1 and CCA2 accounted for 100 per cent of the total variance. Certain soil variables, such as Mg and clay concentration, appear to be associated with increasing slope, whereas sandy, silt and compounds such as Ca, K, Na, and C/N are associated with gentle (A) and moderate slopes (B). Regarding termites and soil variables, we found that termite abundance, species richness, fungus growers (FG), wood-feeding (WF), and soil feeding (SF) were unrelated to soil variables; however, the figure shows that some species, such as *Angulitermes frontalis*, *Profastigitermes putnami*, *Isognathotermes henghi*, Ap5 and WF (wood feeding) are positively associated with Mg. pH of water, bulk density, K, Na, P, C/N, and silt fraction are all positively associated with *Apilitermes longiceps*, *Protermes prorepens*, *Proboscitermes tubuliferus*, *furculitermes winifredae*, and Ap4.

Considering slope gradient, regardless of soil physico-chemical variables, NMDS analysis revealed clustering of some species based on slope gradient. Termite species that preferred gentle slope (A) were clearly distinguished from those that preferred moderate slope (B) and steeper slope (C) (Figure 5).

4. Correlation Analysis between slope gradient and Physico-Chemical Characteristics of Soil

As demonstrated in Table 3, the slope gradient level affected the soil's physical and chemical features. The clay fraction had a significant positive correlation with the slope gradient (r=0.88; p<0.05), but the sand (r= -0.88; p<0.05) and silt (r=0.44; p > 0.05) fractions had a negative correlation. In addition to this, we found significant and positive correlations of water content with clay fraction (r = 0.90; p<0.05) and a positive relationship of bulk density with slope gradient (r = 0.60; p > 0.05). However, there was a positive correlation between pH-water and bulk density (r=0.9; p <0.05).

Concerning soil organic carbon content, a negative correlation was found between slope gradient and organic carbon content (r= -0.82; p<0.05). Soil organic carbon content was positively correlated with sand (r= 0.71; P > 0.05) and silt (r= 0.36 P > 0.05) fractions and negatively correlated with clay fraction (r= -0.71; P > 0.05). The same trend was observed for total nitrogen content, which exhibited negative

Comment [IL10]: This table was not presented in this manuscripts. Please provide it

Comment [IL11]: See above comments on tables.

correlations with slope gradient ($r = -0.71$; $P > 0.05$), clay ($r = -0.48$; $p > 0.05$) fraction and positive correlations with silt ($r = 0.14$; $p > 0.05$) and sand ($r = 0.48$; $p > 0.05$) fractions.

The result for available P showed that it was influenced by both slope gradient ($r = -0.46$; $p > 0.05$) and soil particle size through sand ($r = 0.67$; $p > 0.05$), silt ($r = 0.49$; $p > 0.05$) and clay ($r = -0.67$; $p > 0.05$) fractions. It was also positively correlated with soil nitrogen content ($r = 0.81$; $p < 0.05$) and C/N ratio ($r = 0.89$; $p < 0.05$). The cation exchangeable capacity (CEC) was not significantly ($r = 0.08$; $p > 0.05$) affected by the slope gradient. It was positively correlated with clay ($r = 0.42$; $p > 0.05$), while it was inversely correlated with sand ($r = -0.42$; $p > 0.05$) and silt ($r = -0.77$; $p > 0.05$) fractions. Concerning Exchangeable cations, Exchangeable calcium ($r = -0.57$; $p > 0.05$), potassium ($r = -0.41$; $p > 0.05$), sodium ($r = -0.35$; $p > 0.05$) were negatively correlated with slope gradient, but Magnesium was positively correlated ($r = 0.42$; $p > 0.05$).

5. **Correlation Analysis between termite’s diversity and some Physical and Chemical Characteristics of Soil**

As can be seen under different slope gradients seemed to have a negative impact on species richness ($r = -0.24$; $p > 0.05$) and diversity ($r = -0.5$, $p > 0.05$); the physical and chemical characteristics of the soil had a greater effect on termite attributes. The correlation was significant between species richness and pH-water ($r = 0.85$; $p < 0.05$); it was the same trend with potassium ($r = 0.43$, $p > 0.05$) sodium ($r = 0.35$, $p > 0.05$) and C/N ($r = 0.16$, $p > 0.05$). Concerning the functional groups, CEC showed positive correlation with soil-feeding ($r = 0.77$, $p > 0.05$) as opposed to wood feeders ($r = -0.39$, $p > 0.05$) and fungus growers ($r = -0.13$, $p > 0.05$). Taking into account soil particles size, the results revealed that sand ($r = 0.33$, $p > 0.05$), silt ($r = 0.3$, $p > 0.05$) fractions seemed to favour species richness contrary to clay fraction ($r = -0.33$, $p > 0.05$).

Comment [IL12]: Please capitalize each word to ensure uniformity.

IV. **Discussion**

The present study revealed that the slope gradient has an inverse effect on the sandy and clayey fractions of the soil, with the sandy fraction being higher on gentle slopes and the clayey fraction being higher on steeper slopes. The results for the sandy fraction are due to the geomorphology of the soil, which influences pedogenesis through its influence on erosion and leaching, as well as through the intermediary of the climate. The steeper the slope, the greater the runoff, which causes the movement of large particles, and the more unstable sands are evacuated from the upper slopes to the lower slopes, confirming previous findings [29]. On the other hand, the results for the clay fraction confirm those of [30] who obtained the lowest values on gentle slopes while the highest were observed on higher slopes. The prevalence of clayey texture on the steeper slopes may be related to the recent history of cultivation and logging.

The slope gradient has a significant negative correlation with total organic carbon. This is due to erosion, a geomorphological process that transports particles from steep slopes to gentler slopes. The lowest organic carbon content was found on higher slopes, while the highest was found on lower slopes. Studies reveal an increase in organic carbon content on lower slopes; this could be explained by the soil materials being moved downwards with run-off water from the upper slope and accumulating at the bottom slope position. [31, 32, 33, and 34].

It was observed that soil water content and clay increased as slope gradient increased, this resulted to the higher water-holding capacities of clay fraction with water [35]. Our findings seem to contradict some studies that stated that soil water content declines as the

Comment [IL13]: NMDS analysis was not captured in the discussion.

Comment [IL14]: Where did you determine this parameter ? see your methods

slope gradient increases. According to these studies, the higher soil water content in moderate slope can be explained by long-term buildup of soil moisture[36, 37].

The pH-water data correlated negatively with slope gradient; indeed, the pH-water is lowest on steeply sloping soils (C), and highest on gently sloping soils (A and B)(Table 2). This could be attributable to the loss of base-forming cations due to leaching, as well as runoff caused by faster erosion, which reduces soil pH and consequently raises soil acidity [38]. In addition to this pH-water increased as slope gradient increased. This is likely due to the relatively greater amount of organic matter in moderate and lower slope. The release of hydrogen ions that were associated with organic anions during decomposition contributed to the lower pH [39]. Consequently, bulk density was positively correlated to pH-water.This may be explainedby the soil compaction decreases porosity and reduction of organic matterin steeper slope [40].Conversely, bulk density was negatively correlated to soil organic matter, confirming some researches thatrevealed the importance of organic carbon in soil aggregation,which decreases bulk density [41].

In general, diversity represents the structure of a community; this variable reflects changes in species richness as well as habitat quality [42, 43].We evaluated sampling efficiency using species accumulation curves within 66 quadrats of 5m X 2m. The value of the estimator of total observed species (Chao1, Jackknife 1 and Bootstrap), were all higher than 35 which was the total number of observed species in our study area. This indicates that species richness could be improved with additional sampling effort. Among the termite species sampled, soil-feeding species were the most abundant (94.80% of the total sample). Termites of this functional group are commonly found in habitats such as tropical rain forests and savannah woodland with higher concentrations of organic matter in the soil. It appears that the soil substrate of these areas is suitable for the development of termites of this functional group[44].Furthermore,the dominance index ranges from 0.07 to 0.085, which is regarded as a low value. This index's purpose is to determine whether any species or groups of species are distinctive in a certain environment[45]. The index seeks to determine the significance of species in relation to the entire community by taking into account the ethological differencesbetween taxonomic groups whose interactions may be synergistic or competitive [46]. The dominance index values in this study demonstrated that the slope gradient does not distinguish any particular functional group or species of termites from the others.

Concerning the influence of some physico-chemical properties of soil on termites' assemblages, our results showed that slope gradient strongly affected termite species richness and diversity. However, there was no difference in termite species abundance. Some termite species were associated with some specific slope gradient (Figure 5). This indicates that the slope gradient appears to be one of the termite assemblages distributing factors at the soil level. Some studies suggest that a larger slope inclination leads to nutrients leaching towards the softer slope. This phenomenon governs the geographical distribution of elements and nutrients at ground level, disrupting the living conditions of the least resistant species and resulting in a decline in diversity [47].

Some soil physico-chemical variables affected termite attributes. pH-water was the most influential soil variable for termite species richness. Some studies demonstrated that, alkaline environment seem to favours some functional groups such as fungus-growing species; therefore, our results showed positive correlations between pH-water and all the termite functional groups, which confirmed the researches of [48, 49 ,50]. This is the consequence of the lack of organic matter in tropicalsoilsaccording to [51]. Others soil chemical variables such as potassium, sodium, and C/N ratio showed positive correlations, with termite species richness; in addition, soil feeding termites correlated positively with CEC. These

Comment [IL15]: See above comments on tables. All the mentioned tables were not presented. Please provide all you mentioned.

Comment [IL16]: This was not captured in your methods and results. Please cross check.

results seemed to contradict some findings that showed no correlation between some soil properties and termites assemblages attributes [52], even though sodium enhanced litter termite abundance in some areas [53].

In our study, differences in the distribution of texture, water, carbon content, and other chemical elements between slope gradients resulted in differences within termite communities, with certain termites such as *Procubitermes niapuensis*, *Isognathotermes sulcifrons*, and *Adaiphrotermes* sp. associated with gentle slopes and *Isognathotermes*, *heghi pericapritermes* sp2. Being associated with steeper slopes, diversity was higher on moderate slopes than on steep slopes. These were confirmed by the CCA analysis supplemented by the NMDS ordination, which considered soil particle size chemistry and soil topography to explain the termite species distribution method according to soil slope. This research revealed changes in species richness composition along slope gradients, with a declining trend in the number of termite species with increasing slopes.

Conclusion

This studyWe evaluated the influence of the slope gradient on termite diversity in relation to certain edaphic factors. The results showed a decline in termite diversity. Additionally to this, the increase in soil slope had a negative impact on soil organic carbon,sandy andsilty fractions, but not on the clay fraction. Our findings demonstrated that slope gradient is one of the factors controlling the soil's nutrient availability via leaching and runoff. The significant correlation between pH-water and termite species richness suggests that these abiotic factors had an impact on termite assemblages. In addition, NMDS analysis revealed the clustering of some termite species along specific slope gradients. However, these slope variations had no effect on termite abundance. this encourages us to conduct more detailed research. Based on these results, it is we can concluded that the slope gradient influences the spatial distribution of nutrients and termites at soil level.

References

1. Farmanullah Khan, Zubair Hayat, Waqar Ahmad, Muhammad Ramzan, Zahir Shah, Muhammad Sharif, Ishaq Ahmad Mian, and Muhammad Hanif. Effect of slope position on physicochemical properties of eroded soil. *Soil Environ.*2013. 32(1):22-28.

2. Ziadat FM,Taimeh AY. Effect of rainfall intensity, slope, land use and antecedent soil moisture on soil erosion in an arid environment. *Land Degradation & Development*2013.24(6), 582–590. doi:10.1002/ldr.2239

3. Luk SH, Cai Q,and Wang GP. Effects of surface crusting and slope gradient on soil and water losses in the Hilly loess region, North China. *Catena Supplement*1993.24:29–45.

4. Zhang KL and Hosoyamada K. Influence of slope gradient on interrill erosion of Shirasu soil. *Soil Phys. Cond. Plant Growth, Jpn.*1996.73: 37-44.

5. GesslerPE, ChadwickOA, Chamran F, Althouse Land Holmes K. Modeling soil-landscape and ecosystem properties using terrain attributes. *Soil Sci. Soc. Am. J.*2000;64, 2046–2056.

6. FinkelM, FragmanO and Nevo, E.Biodiversity and interslope divergence of vascular plants caused by sharp microclimatic differences at “evolution canyon II”, lower Nahal Keziv, upper Galilee. *Israel. Isr. J Plant Sci.*2001; 49:285–296. doi: 10.1560/H8TG-HL9L-2D2P-TAGJ

7. Gutiérrez-Jurado HA, Vivoni ER, Cikoski C, Harrison JBJ, Bras RL and Istanbuloglu, E. On the observed ecohydrologic dynamics of a semiarid basin with aspect-delimited ecosystems. *Water Resour. Res.*2013; 49:8263–8284. doi: 10.1002/2013WR014364

8. Pywell PF, Bullock JM, Walker KJ, Coulson SJ, Gregory SJandStevenson MJ. Facilitating grassland diversification using the hemiparasitic plant *Rhinanthus minor*. *J. Appl. Ecol.*2004; 41:880-887.

9. Kumari, Rina, Minu Sachdev, Ram Prasad, Amar P. Garg, Shweta Sharma, Pham Huong Giang, Ajit Varma. Microbiology of Termite Hill (Mound) and Soil. Intestinal Microorganisms of Termites and Other Invertebrates.2006;pp 351-372. Helmut König Ajit Varma (Eds.).492p.

10. Engel MS, Grimaldi DA& Krishna K. Termites (Isoptera): Their phylogeny, classification, and rise to ecological dominance. *American MuseumNovitates.*2009; 3650: 1-27. doi : 10.1206/651.1

11. Krishna K., Grimaldi DA, Krishna K, Engel MS,. Treatise on the Isoptera of the World. *Bulletin of the American Museum of Natural History*, 2013; 377 (7): pp. 2704.

12. Deblauwe Isra, Dibog Luc, Missoup Alain D, Dupain Jef, Elsacker Van Linda, Dekoninck Wouter, Bonte Dries and Hendrickx Frederik.Spatial scales affecting termite diversity in tropical lowland rainforest: A case study in southeast Cameroon.African Journal of Ecology. 2007; 46(1):5 – 18. DOI: 10.1111/j.1365-2028.2007.00790.x

13. Jouquet P, Guilleux N, shanbhang R, Subramanian S. Influences of soil types on the properties of termite mound nests in southern India *Applied Soil Ecology*.2015;96:282-287

14. Eggleton P, Tayasu, I, Feeding groups, lifetypes and the global ecology of termites. *Ecological Research*.2001; 16, 941–960.

15. Dambros CS, Morais JW, Azevedo RA, GotelliNJ. Isolation by distance, not rivers, control the distribution of termite species in the Amazonian rain forest. *Ecography*2017; 40:1242–1250.

16. Gao YF, Gong YJ, Cao LJ, Chen JC, Gao YL, Mirab-balou M, Chen M, Hoffman AA, Wei SJ. Geographical and interspecific variation in susceptibility of three common thrips species to the insecticide, spinetoram *J Pest Sci*.2020; <https://doi.org/10.1007/s10340-019-01128-2>

17. Becker Fabian. Soil Erosion Risk Assessment in the Upper Mefou Subcatchment, Southern Cameroon Plateau. *M. Sc. Geographie (Terrestrische Systeme)*.2015; Freie Universität Berlin.150 p.

18. Tchindjang Mesmin, Miaro IIIIudovic, Mboringong Fideline, EtogaGilles; VoundiandEric Emmanuel Pierre, NgomJonathan. Environmental Impacts of the Oil Palm Cultivation in Cameroon. *Preprints*.2021; DOI: 10.5772/intechopen.97862

19. Achoundong, G. Les forêts sommitales au Cameroun. Végétation et flore des collines de Yaoundé. *Bois & Forêts des tropiques*.1996 ; 247(247) :37–52. <https://doi.org/10.19182/bft1996.247.a19890>

20. Jones David T, Eggleton Paul. Sampling termite assemblages in tropical forests: testing a rapid biodiversity assessment protocol. *Journal of Applied Ecology*.2000; 37(1), 191–203. doi:10.1046/j.1365-2664.2000.00464.x

21. Bouillon A,Mathot, G. Quel est ce termite africain ? 1965 ;*Zooleo*. N° 1. (eds.). Université de Léopoldville, 115p.

22. Hellemans S, Deligne J, Roisin Y & Josens, G. Phylogeny and revision of the “Cubitermes complex” termites (Termitidae : Cubitermitinae). *Systematic Entomology*. 2021. 46(1), 224–238. doi:10.1111/syen.12458

23. Pauwels JM, van Ranst E, Verloo M, Mvondo ZE A. Manuel de laboratoire de pédologie. Méthodes d’analyses de sols et de plantes, équipements, gestion de stocks de verrerie et de produits chimiques.1992 ;*AGCD, Publications Agricoles* 265p.

24. Tellen Valentine Asong, Yerima, Bernard PK. Effects of land use change on soil physicochemical properties in selected areas in the North West region of Cameroon. *Environmental Systems Research*.2018;7(1), 3–29. Doi: 10.1186/s40068-018-0106-0

25. Tabatabai MA and Bremner JM.Assay of Urease Activity in Soils. American Journal of Soil Science Society.1972;41:350-352.

26. Bremner JM. and MulvaneyCS. “Total nitrogen”, In: A.L. Page, R.H. Miller and D.R. Keeny, (Eds.), Methods of Soil Analysis, American Society of Agronomy and Soil Science Society of America, Madison.1982; pp. 1119-1123.<https://doi.org/10.2134/agronmonogr9.2.2ed.c31>

27. Moreno CEandHalffter G. Assessing the Completeness of Bat Biodiversity Inventories Using Species Accumulation Curves. *Journal of Applied Ecology*. 2000; 37, 149-158. <https://doi.org/10.1046/j.1365-2664.2000.00483.x>

28. Legendre P, Borcard D, Peres-Neto PR. Analyzing beta diversity: partitioning the spatial variation of community composition data. *Ecological Monographs*.2005; 75(4):435-450 .DOI:10.1890/05-0549

29. Mulugeta A. Effect of Slope Gradient on Selected Soil Physicochemical Properties of Dawja Watershed in Enebse Sar Midir District, Amhara National Regional State. *American Journal of Scientific and Industrial Research*.(2015);6(4): 74-81.

30. Fanuel L, Kibebew L, Tekalign M, Erik K, Heluf G. Physiographic characteristics of agricultural lands and farmer’s soil fertility management practices in Wolaita zone, Southern Ethiopia. *Environmental Systems Research*.2016;5(1): 1-15. DOI 10.1186/s40068-016-0076-z

31. Abebe N, Endalkachew K. Physicochemical characterization of Nitisol in Southwestern Ethiopia and its fertilizer recommendation using NuMaSS. *Global Advanced Research Journal of Agricultural Science*.2012;1(4): 66-73

32. Amuyou UA. and Kotingo KE. Toposequence analysis of soil properties of an agricultural field in the Obudu Mountain slopes, cross river state- Nigeria. *European journal of physical and agricultural sciences*.2015;3(1):2056-5879

33. Wubie, Mesfin Anteneh; Assen, Mohammed 2019. Effects of land cover changes and slope gradient on soil quality in the Gumara watershed, Lake Tana basin of North–West Ethiopia. *Modeling Earth Systems and Environment*, 6:85–97. Doi: 10.1007/s40808-019-00660-5

34. Kedir Mohammed.Influence of Slope Gradient on Coffee Soil Fertility in Goma Woreda of Oromia Region. *Advances in Oceanography & Marine Biology*.2021; 2(4):1-6 DOI: 10.33552/AOMB.2021.02.000544

35. Astrid M. Lambooy. Relationship between cation exchange capacity, clay content and water retention of Highveld soils, *South African Journal of Plant and Soil*.1984;1(2): 33-38, DOI: 10.1080/02571862.1984.10634106

36. Takahashi K,Murayama Y. Effects of topographic and edaphic conditions on alpine plant species distribution along a slope gradient on Mount Norikura, central Japan. *Ecol. Res.* 2014; 29, 823–833.

37. Zhao C,Zhang HL,Li JL, Tian J, Wu WZ, Jin X, Zhang XF, Jiang YW, Wang XL,He CS.Analysis of the relationships between the spatial variations of soil moisture and the environmental factors in the upstream of the Heihe River watershed. J. Lanzhou Univ. *Nat Sci*.2014;50:338–347.

38. Nega E, Heluf G. Effect of Land Use Changes and Soil Depth on Soil Organic Matter, Total Nitrogen and Available Phosphorus Contents of Soils in Senbat Watershed, Western Ethiopia. *Journal of Agricultural and Biological Science*.2013; 8(3):206-2012.

39. LiYuan, Li Zhou, Cui Song, Zhang, Qingping. Trade-off between soil pH, bulk density and other soil physical properties under global no-tillage agriculture. *Geoderma*.2020;361, 114099–114108. doi:10.1016/j.geoderma.2019.114099

40. Prashant Deo Singh, Ashok Kumar, DhyaniBP, Satendra Kumar, ShahiUP, Adesh Singh, Anand Singh. Relationship between compaction levels (bulk density) and chemical properties of different textured soil. *Int. J. Chem. Stud.*2020; 8(5):179-183. DOI: 10.22271/chemi.2020.v8.i5c.10294
41. Achalu C, Heluf G, Kibebew K, Abi T. Status of selected physicochemical properties of soils under different land use systems of Western Oromia, Ethiopia.*J. Bio. Env. Sci.*2012;2(3), 57-7
42. Guo Q, Berry WL. Species richness and biomass: Dissection of the hump-shaped relationships. *Ecology.*1998;79, 2555–2559
43. Mittelbach G,Steiner C,Scheiner S Gross, K Reynolds H,Waide R,Willig, M,Dodson,S,Gough L.What is the observed relationship between species richness and productivity? *Ecology.*2001;82:2381–2396.
44. Jones DT, EggletonP. Global Biogeography of Termites: A Compilation of Sources. In *Biology of Termites: A Modern Synthesis*; Springer Science and Business Media LLC: Berlin, Germany.2011;pp. 477–498.
45. Lohbeck M, Bongers F, Martinez- Ramos M and Poorter L. The importance of biodiversityand dominance for multiple ecosystem functions in a human- modified tropical landscape*Ecology.*2016;97:2772– 2779
46. Heip CHR, Herman PMJ, Soetaert K. Indices of diversity and evenness. *Océanis (Paris).*1998;24(4): 61-87
47. Kopec M, Zarzycki J, Gondek, K. Species diversity of sub montane grasslands: Effects of topographic and soil factors. *Pol. J. Ecol.* 2010;58 : 285–295
48. BOYER Ph.Action des termites constructeurs sur l'évolution des sols tropicaux. *Annales Sci. Nat., Zoologie(Paris).* 128 Sér. T. 1973; 15 : 329-498
49. Kanvaly D, Souleymane K, Daouda A and Linsenmair KE.Termite diversity and abundance across fire-induced habitat variability tropical moist savanna (Lamto, Central Côte d'Ivoire).*Journal of Tropical Ecology.* 2010; 26(3):323-334
50. Kanyi Nahashon Chege, Karuri Hannah, Nyasani Johnson O, Mwangi Benson. Land use effects on termite assemblages in Kenya, Heliyon.2021; 7(12) <https://doi.org/10.1016/j.heliyon.2021.e08588>
51. Watson JP. The use of mounds of the termite *Macrotermes falciger* (Gerstäcker) as a soil amendment. *Journal of Soil Science.*1977; 28(4): 664-627
52. Bourguignon T, Drouet T, Šobotník J, Hanus R, Roisin Y. Influence of Soil Properties on Soldierless Termite Distribution. *PLoS One.*2015; 13; 10(8):e0135341. doi: 10.1371/journal.pone.0135341. Erratum in: PLoS One. 2015;10(11):e0143776
53. Kaspari M, Yanoviak SP, Dudley R, Yuan M, Clay NA Sodium shortage as a constraint on the carbon cycle in an inland tropical rainforest. *Proc Natl Acad Sci USA.*2009;106:19405–19409




Figure 1: Location of the study area




Figure 2: Rarefaction Curves of the global species richness observed in three sloped gradients, according to the nonparametric estimators




Figure 3: Rarefaction curves of species richness observed on different gradients slope A= [0°; 15°] ; B =[15.1°; 31°] and C= > 31° [.




Figure 4: Canonical Correspondence Analysis (CCA). A bi-plot ordination of the 35 termite's species and functional groups according to environmental variables and slope gradient.




Figure 5: NMS analysis of the different slope gradients based on the Composition of termites assemblages (FG: fungus growers, SF: soil Feeding; WF: wood feeding)