

Original Research Article

Effect of GA₃ and Potassium on Growth and Yield in Red Radish (*Raphanus sativus* L.) under subtropical conditions of Jammu

ABSTRACT

An experiment entitled, **Effect of GA₃ and Potassium on growth and yield in Red Radish (*Raphanus sativus* L.) under subtropical conditions of Jammu** was carried out at Vegetable Experimental Farm, Division of Vegetable Science, SKUAST Jammu during *Rabi* season of 2022-2023. The experiment was laid out under factorial RBD design with three replications comprising of sixteen treatments. The treatments consisted of four doses of Potassium i.e., 40, 50, 60 and 70 kg/ha and four concentrations viz. 0, 100, 200 and 300 ppm of GA₃.

The growth and yield parameters maximum viz. shoot length (34.15 cm), root length (18.73 cm), number of leaves/root (14.30), root weight (220.49 g), shoot weight (119.47 g), root-shoot ratio (2.04), fresh root shoot yield (112.42 q/ha) were recorded with 200 ppm GA₃ application. Similarly, highest shoot length (29.06 cm), root length (17.53 cm), number of leaves/root (11.18), root weight (171.67 g), shoot weight (104.84 g), root-shoot ratio (1.87), fresh root shoot yield (92.31 q/ha), were recorded with 50 kg/ha K₂O application.

However, highest root length (19.20 cm), number of leaves/root (14.87), root weight (226.14 g), shoot weight (123.54 g), root-shoot ratio (2.10), fresh root shoot yield (115.90 q/ha) were recorded with treatment combination G3K2 (200 ppm GA₃ + 50 K₂O/ha). Based on the investigation, it can be concluded that the application of 200 ppm GA₃ at three stages in combination with 50 kg potassium/ha resulted in the highest growth parameters and yield.

Keywords: Growth regulator, Potassium, Radish, Root shoot ratio

INTRODUCTION

Radish (*Raphanus sativus* L.) is one of the most significant winter vegetable crop grown in both tropical and temperate region. It is popular choice for cultivation as it is easy to grow, fits well in crop rotation, and is rapidly maturing cash and catch crop. It is highly cross-pollinated crop that belongs to the family Cruciferae and finds its origin in Europe and Asia. The most popular edible part of radish is the tuberous root, which is consumed either cooked or raw, and tops can be used as leafy vegetable. It helps in the treatment of gastrodynia, gall bladder issues, sleep disorders, persistent diarrhea, neuralgic headaches, and urinary symptoms (Sandhu, 1993). It is good source of Vitamin C and minerals (i.e., calcium, potassium, and phosphorus) (Zohary *et al.*, 2012). 100 g fresh edible portion of radish contains 0.7% protein, 3.4-6.8% carbohydrates and 0.2% fat and 50 I.U. of vitamin A. Its tender roots provide about 50 mg of calcium, 22 mg of phosphorus and 0.5 mg of iron per 100 g of edible portion.

In India, radish occupied an area of 209 thousand hectares with an annual production of 3347 thousand metric tonnes (NHB, 2021). It is cultivated in almost all states of India. The major radish producing states include West Bengal, Haryana, Punjab, Bihar and Assam. In Jammu & Kashmir, it is grown over an area of 3.20 thousand hectares with an annual production of 116.07 thousand metric tonnes (NHB, 2021).

In the recent past, due to significant increase in area under radish crop, the demand for quality seed has increased manifold. State Government and Government of India have laid out a lot of emphasis to improve availability of quality seed, which has not only increased production and productivity of this crop, however, it also provided new opportunities for export of quality seed. The nation's total production of vegetable seeds is insufficient to fulfil the nation's rising demand. Currently, the demand for quality seeds is met to the extent of 20% only. Farmers themselves meet 75% seed demand through own saved

Comment [1]: English of your manuscript is extremely poor. Please consider taking help of a native English speaker or hire editing service.

Comment [2]: Correct spacing and grammatical issues in entire manuscript.

Comment [3]: Major-Abstract-
In this part, there is no need to mention too many details such as experiment time and place, the specific values measured, etc. The authors should mainly summarize the main findings and conclusions.

Comment [4]: Introduction-1.
The content of the three paragraphs needs to be adjusted, the authors can briefly propose the objective of this study.

seeds (Poonia *et al.*, 2013). India continues to import vegetable seeds from foreign nations, the two main ones being cabbage and pea.

The physiological efficiency of plants, particularly photosynthetic capacity and efficient assimilate partitioning, can be improved by plant growth regulators (Solaimalai *et al.*, 2001). By promoting the transfer of photo assimilates, vegetable crop productivity can be raised. In particular, it has been noted that GA₃ has amazing effects on promoting blooming and raising seed production in vegetable crops. Gibberellins are crucial for many aspects of plant growth and development, including seed germination. Gibberellic acid stimulates the cells of growing seeds to produce mRNA molecules, which controls the activity of hydrolytic enzymes. The use of GA₃ controls a number of physiological processes and causes metabolic changes that have an impact on the quantity and quality of the intended product. An increase in seed quality parameters due to greater accumulation of food reserves by the application of GA₃ has been reported (Ghanome, 2011). The use of potassium fertilizer is an effective way to increase radish productivity. Potassium regulates the opening and closing of stomata. It activates enzymes that are required for generation of ATP. Starch and protein synthesis are facilitated by potassium in plants. It is also required for transport of sugar to the storage organs (Chhetri *et al.*, 2019). That the growth and yield of red radish can successfully be produced under Jammu conditions however, the quantity of the growth and yield is less. Therefore, to increase the quantity and quality of growth and yield through application of growth regulators and nutrients, the present study has been designed.

MATERIALS AND METHODS

The present investigation was carried out on **Effect of GA₃ and Potassium on the growth and yield in Red Radish (*Raphanus sativus* L.) under subtropical conditions of Jammu** was carried out at Vegetable Experimental Farm, Division of Vegetable Science, SKUAST Jammu during Rabi season of 2022-2023. The investigation was conducted at Chatha which is located at an elevation of 332 m above mean sea level having geographical bearing of 32.3° North latitude and 74.53° East longitude. The experiment was laid out under factorial RBD design with three replications comprising of sixteen treatments. The treatments consisted of four doses of Potassium i.e., 40, 50, 60 and 70 kg/ha and four concentrations viz. 0, 100, 200 and 300 ppm of GA₃.

Comment [5]: Source of this data?

Table 1: Details of treatment

GA ₃ application (04)	Potassium doses (04)
G ₁ : 0 ppm	K ₁ : 40 kg/ha
G ₂ : 100 ppm	K ₂ : 50 kg/ha
G ₃ : 200 ppm	K ₃ : 60 kg/ha
G ₄ : 300 ppm	K ₄ : 70 kg/ha

The detail of various cultural operations carried out during the crop growing have been given below:

Land preparation for mother block

The field was ploughed twice followed by planking to attain good tilth. Field layout and preparation of ridges was done manually with the help of rope and spade. Well rotten FYM @ 20 tonnes per hectare was applied in the field before sowing. Recommended dose of N, P @ 60, 30 kg per hectare and four dose of K treatment @ 40,50,60,70 kg per hectare was applied as a basal dose at the time of sowing. The seeds were sown on ridges by dibbling methods in 5 November 2022. The GA₃ were sprayed at 30 Days after sowing in mother block. Proper aftercare operations in respect of irrigation, plant protection was undertaken till the roots were ready for the transplanting.

Preparation of field for transplanting of stecklings

The field was ploughed twice followed by levelling and planking to attain good tilth. Field layout and preparation of pits was done manually with help of rope and spade. Well rotten FYM @ 20 tonnes per hectare was applied in the field before planting. Recommended dose of N, P @ 60, 30 kg per hectare and four dose of K treatment @ 40,50,60,70 kg per hectare was applied as a basal dose at the time of planting. The steckling were prepared by cutting 1/3rd of the tops and 1/3 of the roots were transplanted as per the treatment. Other intercultural operations were carried out in accordance with the packaging of practices of radish crop seed production from time to time.

Preparation of spray solution

GA₃ stock solutions of 100 ppm was prepared by dissolving 1 g of pure GA₃ in small quantity of alcohol and then making the final volume up to 1 liter by adding distilled water.

The solutions were prepared fresh at the time of each spray and dilutions of required concentrations were made using stock solution. The GA₃ sprayed at 30 Days after sowing and 15 Days after transplanting of steckling, at the time of siliquae formation.

The growth and yield parameters viz. shoot length, root length, number of leaves/root, root weight, shoot weight, root-shoot ratio, fresh root shoot yield.

Statistical analysis

The data recorded on various parameters of the present study after proper tabulation were put to statistical analysis as per the randomized complete block method prescribed by (Panse and Sukhatme, 1989) to draw the inferences of the study. The significance of treatment effects was tested through variance ratio and the significance of difference between any two means was judged with the critical difference (C.D) at 5 per cent level of significance was worked out. Data for the parameters were put to statistical analysis by using OPSTAT software developed by O.P. Sheoran *et al.* (1998) of CCS HAU, Hissar.

RESULTS

Shoot length (cm)

It was evident from table 1 that the Growth regulators (GA₃) and potassium have significantly influenced the shoot length. Maximum shoot length 34.15 cm was in G₃ which was at par with G₄. Lowest shoot length 18.73 cm was recorded in G₁ which was significantly lower than all of the treatments. Similarly, The Potassium levels had significant effect on shoot length. Maximum shoot length was obtained K₂ 29.06 cm which had at par results with K₃ and K₄ and Minimum shoot length was recorded in K₁ 26.70 cm was recorded in K₁ which was significantly lower than all of the treatments.

However, the interaction between GA₃ and Potassium with respect to shoot length was found to be non-significant.

Root length (cm)

It was evident from table 2 that the Growth regulators (GA₃) and potassium have significantly influenced the root length of red radish. Maximum root length (18.73 cm) was obtained in G₃ which

Comment [6]: Better to use t/ha format. Change all other units too.

Comment [7]: Provide some figures there is no figure at all

exhibited at par results with G₄. Lowest root length 14.78 cm was recorded in G₁ which was significantly lower than all of the treatments. Similarly, The Potassium levels had significant effect on root length. Maximum root length was obtained K₂ dose (40kg/ha) 17.53 cm which had at par results with K₃ and K₄. Minimum root length was recorded in K₁ 16.38 cm was recorded in K₁ which was significantly lower than all of the treatments.

Interaction studies depicts that a combination of G₃K₂ maximum root length (19.20 cm) which was at par with the treatment combinations of G₃K₃, G₃K₄, G₄K₃ and G₄K₄. Lowest root length was recorded in a treatment combination of G₁K₁ (13.19cm) which was significantly lower than all of the treatment combinations.

Number of leaves per root

It was evident from Table 3 that the number of leaves per root of red radish was significantly influenced by the Growth regulators (GA₃) and potassium. G₃ had the maximum number of leaves per root 14.30, which was at par with G₄. The Lowest number of leaves per root 6.07, was recorded in G₁, which was significantly lower than all of the treatments. Similarly, the number of leaves per root was significantly affected by the Potassium levels. K₂ had the Maximum number of leaves per root 11.18, which had at par results with K₃ and K₄. The Minimum number of leaves per root 10.41, was recorded in K₁, which was significantly lower than all of the treatments.

The interaction studies depicted that the treatment combination of G₃K₂ had the maximum number of leaves per root 14.87 cm, which was at par with the treatment combinations of G₃K₃, G₃K₄, G₄K₃, and G₄K₄. The lowest number of leaves per root was recorded in the treatment combination of G₁K₁ 5.30 cm, which was significantly lower than all of the treatment combinations.

Root weight (g)

It was evident from table 4 that the Growth regulators (GA₃) and potassium have significantly influenced the root weight of red radish. Maximum root weight (220.49 g) was in G₃ which was at par with G₄. Lowest root weight (112.33 g) was recorded in G₁ which was significantly lower than all of the treatments. Similarly, The Potassium levels had significant effect on root weight. Maximum root weight was obtained K₂ (171.67g) which had at par results with K₃ and K₄ and Minimum root weight was recorded in K₁ (164.55g) was recorded in K₁ which was significantly lower than all of the treatments.

Interaction studies depicts that a combination of G₃K₂ maximum root weight (226.14 g) which was at par with the treatment combinations of G₃K₃, G₃K₄, G₄K₃ and G₄K₄. Lowest root weight and shoot weight was recorded in a treatment combination of G₁K₁ (111.17g) which was significantly lower than all of the treatment combinations.

Shoot weight (g)

It was evident from table 5 that the Growth regulators (GA₃) and potassium have significantly influenced the shoot weight of red radish. Maximum shoot weight (119.47 g) was obtained in G₃ which exhibited at par results with G₄. Lowest shoot weight (78.20 g) was recorded in G₁ which was significantly lower than all of the treatments. Similarly, The Potassium levels had significant effect on shoot weight. Maximum shoot weight was obtained K₂ dose (104.84g) which had at par results with K₃ and K₄. Minimum shoot weight was recorded in K₁ (99.36g) was recorded in K₁ which was significantly lower than all of the treatments.

Interaction studies depicts that a combination of G₃K₂ maximum shoot weight (123.54g) which was at par with the treatment combinations of G₃K₃, G₃K₄, G₄K₃ and G₄K₄. Lowest shoot weight was recorded in a treatment combination of G₁K₁ (75.90 g) which was significantly lower than all of the treatment combinations.

Root-Shoot ratio

The root shoot ratio of red radish was significantly influenced by the Growth regulators (GA_3) and potassium as evident from table 6. The maximum root shoot ratio of 2.04 was observed in G_3 , which was at par with G_4 . The lowest root shoot ratio of 1.38 was recorded in G_1 , which was significantly lower than all of the treatments. Similarly, the root shoot ratio was significantly affected by the Potassium levels. The maximum root shoot ratio of 1.87 was obtained in K_2 , which had at par results with K_3 and K_4 . The minimum root shoot ratio of 1.75 was recorded in K_1 , which was significantly lower than all of the treatments.

It was observed from the interaction studies that the treatment combination of G_3K_2 (2.10) was at par with the treatment combinations of G_3K_3 , G_3K_4 , G_4K_3 , and G_4K_4 . The lowest root shoot ratio of 1.28 was recorded in the treatment combination of G_1K_1 , which was significantly lower than all of the treatment combinations.

Fresh root shoot yield (q/ha)

It was evident from table 7 that the Growth regulators (GA_3) and potassium have significantly influenced the fresh root shoot yield of red radish. Maximum fresh root shoot yield 98.75 q/ha was in G_3 which was at par with G_4 . Lowest fresh root shoot yield 72.96 q/ha was recorded in G_1 which was significantly lower than all of the treatments. Similarly, The Potassium levels had significant effect on fresh root shoot yield. Maximum fresh root shoot yield was obtained K_2 92.06 q/ha which had at par results with K_3 and K_4 and Minimum fresh root shoot yield was recorded in K_1 84.69 q/ha was recorded in K_1 which was significantly lower than all of the treatments.

Interaction studies depicts that a combination of G_3K_2 (103.58 q/ha) fresh root shoot yield which was at par with the treatment combinations of G_3K_3 , G_3K_4 , G_4K_3 , and G_4K_4 . Lowest fresh root shoot yield was recorded in a treatment combination of G_1K_1 (71.83 q/ha) which was significantly lower than all of the treatment combinations.

DISCUSSIONS

It was observed that GA_3 and potassium had a significant effect on the growth traits of red radish. When 200 ppm GA_3 was sprayed and 50 kg potassium per hectare was applied, the shoot length was recorded as 34.15 cm and 29.06 cm, respectively. However, the interaction studies 200 ppm GA_3 + 50 kg potassium per hectare did not show a significant effect on shoot length. Similarly, the root length was significantly affected by both GA_3 and potassium. The maximum root length recorded was 18.73 cm and 17.53 cm when 200 ppm GA_3 and 50 kg potassium per hectare were applied, respectively. The interaction had a statistically significant effect, resulting in a root length of 19.20 cm when the combination of 200 ppm GA_3 and 50 kg potassium per hectare was applied. The number of leaves per root was significantly influenced by both GA_3 and potassium. The highest number of leaves per root (14.30 and 11.18) was recorded when 200 ppm GA_3 and 50 kg potassium per hectare were applied, respectively. The interaction also had a significant effect on the number of leaves per root, with a recording of 14.87 when the treatment combination of 200 ppm GA_3 and 50 kg potassium per hectare was applied.

The increase in shoot length, root length, and number of leaves per root observed in response to GA_3 application may be attributed to the stimulation of cell division, cell size, and metabolic activities in the plant. The enhanced growth observed at optimum levels could be attributed to the pivotal role of potassium in activating enzymes, regulating stomata, and facilitating photosynthesis. These findings are supported by similar results reported by Sharma (2003), Mehta & Patnayak (2005), Jatav *et al.* (2007), Abou EL-Nasr (2008), Kiran (2016), and Shweta *et al.* (2018).

The root and shoot yield traits of red radish were significantly influenced by GA_3 and potassium. The maximum root weight (220.49 g), shoot weight (119.47 g), root-shoot ratio (2.09), and fresh root-

shoot yield (98.75 q/ha) were recorded when 200 ppm GA₃ was sprayed. Similarly, the maximum root weight (171.67 g), shoot weight (109.89 g), root-shoot ratio (1.87), and fresh root-shoot yield (92.06 q/ha) were recorded when 50 kg potassium per hectare was applied. The interaction effect between GA₃ and potassium was found to be significant for all yield traits. The maximum root weight (226.14 g), shoot weight (123.54 g), root-shoot ratio (2.10), and fresh root-shoot yield (103.58 q/ha) were recorded when the treatment combination of 200 ppm GA₃ and 50 kg potassium per hectare was applied.

The enhancement in yield attributes, such as root weight, shoot weight, root-shoot ratio, and fresh root-shoot yield, can be attributed to the stimulation of cell division, cell size, and expansion resulting from the application of GA₃. Similarly, increased yield attributes observed with an optimum level of potassium can be attributed to its role in carbohydrate synthesis and translocation, cell extension, and improved fertilizer use efficiency. These findings align with the results reported by Sharma *et al.* (2003), Anjaiah & Padmaja (2006), Patel (2007), Bawkar *et al.* (2011), Khan *et al.* (2021), Shweta *et al.* (2018), Nagaich and Mishra (2019).

CONCLUSIONS

Based on the investigation, it can be concluded that the application of 200 ppm GA₃ at three stages (15 days after seed sowing, 30 days after steckling planting and at the time of siliqua formation) in combination with 50 kg potassium/ha resulted in the highest growth parameters and yield. This treatment combination significantly improved the highest growth parameters and yield.

REFERENCES

1. Anjaiah, T. and Padmaja, G. 2006. Effect of potassium and farm yard manure on yield and quality of carrot. *Journal of Research ANGRAU.*, **34**(2): 91-93.
2. Sandhu, M. K. 1993. Root crops. In: Vegetable crops (Eds Bose and Som), *Naya Prakash Kolkata*, India, pp: 470-488.
3. Zohary, D., Hopf, M., & Weiss, E. 2012. Domestication of plants in the old world 4th edi. Oxford. *Oxford University Press*.
4. NHB. 2021. Horticulture Statistics At a Glance-2021. National Horticulture Board, Government of India. Gurgaon, India. www.nhb.gov.in.
5. Poonia, T. C. 2013. History of seed production and its key issues. *International Journal of Food, Agriculture and Veterinary Science*, **3**(1): 148-154.
6. Solaimalai A, Sivakumar C, Anbumani S, Suresh T, Arumugam K. 2001. Role of plant growth regulators in rice production- a review. *Agriculture Review*, **22**: 33-40.
7. Ghanome, A. A., Saleh, S. A., Read, G. S., Fawzey, Z. F. and El-Abd, S. O. 2011. Enhancement of carrot seed yield and quality by root dipping in gibberellic acid. *Journal of Applied Science Research*, **7**(12): 1821-1827.
8. Chetri, R.K., Panta, R., Shrestha, R. K. 2019. Effect of spacing and Potassium application on growth and yield of radish *Raphanus sativus* cv. mino early. *The International Journal of Biological Research*, **2**: 176-180.
9. Panse, V. G. and Sukhatme, P. V. 1989. *Statistical Methods for Agriculture workers* ICAR, New Delhi.
10. Patel, R., Tomar, B. S., Khan, R., and Yadav, K. 2017. Response of Radish (*Raphanus sativus* L.) to different concentration of Gibberellic acid. *International Journal of Chemical Studies*, **5**(6): 458-459.
11. Sharma, A., Sharma, R. P., Sonia, S. and Shartna J. 2003. Influence of integrated use of nitrogen, phosphorus, potassium and farm yard manure on the yield attributing traits and marketable yield of carrot (*Daucus carota*) under the high hills dry temperate conditions of north- western Himalayas.

Comment [8]: Many references are old update with new and recent references. May of the references are missing from the list. Please cross-check all references and provide those are missing.

Indian Journal of Agriculture Sciences,73(9): 500-503

12. Mehta, S.K. and Patnayak, C. P. 2005. Studies the effect of growth hormone (GA₃) on growth yield attributes of radish (*Raphanus sativus* L.) var. Pusa Deshi. *Journal of Advancements in Plant Science*,24(2): 121-126.
13. Shweta., Mal, D., Thakur, V., Dutta, 2018. Effect of different plant growth regulators (GA₃ and NAA) on growth and yield parameters of radish (*Raphanus sativus* L.) Var. Pusa Reshmi. *Journal of Pharmacognosy and Phytochemistry*,7(3): 3434-3437.
14. Nagaich, K.N., Mishra, P. 2019. Response of gibberellic acid on growth and yield of radish (*Raphanus sativus* L.) Cv. Japanese white. *Journal of Pharmacognosy and Phytochemistry*,8(2): 1521-1523.
15. Khan, B. P., Ara, N., Ali, S., Baigh, K.K., Wahab, A. 2021. Effect of seed priming on radish with different at various soaking durations. *Pakistan Journal of Agriculture Research*,34(1): 77-84.
16. Kiran, A., Jilani, M. S., Waseem, K. and Sohail, M. 2016. Effect of organic manures use of inorganic fertilizers on growth and yield of radish of radish (*Raphanus sativus* L.). *Pakistan Journal of Agriculture Research*,29(4): 363-372.
17. Jatav, B. S. 2007. Effect of plant growth regulators on growth and yield of Radish (*Raphanus sativus* L.) Var. Japanese white. *M.Sc. (Ag) Thesis*, RVSKVV, College of Agriculture, Gwalior.
18. Bawkar, S. O., Wagh, A. P., Dod, V. N., Das, B. 2011. Effect of spacing and seed soaking with GA₃ on growth, yield and quality of carrot. *The Asian Journal of Horticulture*,6(2): 373-375.

Table 1 Effect of GA₃ and potassium on Shoot length (cm) in red radish (SJRR-01)

Shoot length (cm)					
Treatments	K ₁	K ₂	K ₃	K ₄	Mean
G ₁	17.54	18.66	18.29	20.41	18.73
G ₂	25.14	26.66	26.62	26.71	26.28
G ₃	31.92	34.92	34.84	34.93	34.15
G ₄	32.18	35.98	34.26	33.99	34.10
Mean	26.70	29.06	28.50	29.01	
	SEm±			CD (0.05)	
G	0.62			1.79	
K	0.62			1.79	
(G×K)	1.24			NS	

Table 2 Effect of GA₃ and potassium on Root length (cm) in red radish (SJRR-01)

Root length (cm)					
Treatments	K ₁	K ₂	K ₃	K ₄	Mean
G ₁	13.19	15.31	15.33	15.30	14.78
G ₂	16.10	17.17	17.10	17.07	16.86
G ₃	18.23	19.20	18.80	18.70	18.73
G ₄	18.00	18.43	18.67	18.93	18.51
Mean	16.38	17.53	17.48	17.50	
	SEm±			CD (0.05)	
G	0.13			0.37	
K	0.13			0.37	

(G×K) 0.26 0.74
NOTE: G₁-0 ppm, G₂-100 ppm, G₃- 200 ppm, G₄-300 ppm, K₁-40 kg/ha, K₂-50 kg/ha, K₃-60 kg/ha, K₄-70 kg/ha

Table3 Effect of GA₃ and potassium on Number of leaves per root in red radish (SJRR-01)

Number of leaves per root					
Treatments	K ₁	K ₂	K ₃	K ₄	Mean
G ₁	5.30	6.20	6.00	6.77	6.07
G ₂	8.82	9.47	9.20	9.23	9.18
G ₃	13.77	14.87	14.30	14.27	14.30
G ₄	13.77	14.20	14.83	14.27	14.27
Mean	10.41	11.18	11.08	11.13	
	SEm±			CD (0.05)	
G	0.11			0.31	
K	0.11			0.31	
(G×K)	0.21			0.62	

NOTE: G₁-0 ppm, G₂-100 ppm, G₃- 200 ppm, G₄-300 ppm, K₁-40 kg/ha, K₂-50 kg/ha, K₃-60 kg/ha, K₄-70 kg/ha

Table 4 Effect of GA₃ and potassium on Root weight (g) in red radish (SJRR-01)

Root weight (g)					
Treatments	K ₁	K ₂	K ₃	K ₄	Mean
G ₁	111.17	113.67	113.12	111.34	112.33
G ₂	124.67	127.75	126.44	126.66	126.38
G ₃	204.84	226.14	225.74	225.22	220.49
G ₄	217.52	219.10	219.68	219.50	218.95
Mean	164.55	171.67	171.24	170.68	
	SEm±			CD (0.05)	
G	1.16			3.36	
K	1.16			3.36	
(G×K)	2.33			6.72	

Table 5 Effect of GA₃ and potassium on Shoot weight (g) in red radish (SJRR-01)

Shoot weight (g)					
Treatments	K ₁	K ₂	K ₃	K ₄	Mean
G ₁	75.90	79.81	78.43	78.65	78.20

G₂	92.88	96.96	95.45	94.52	94.95
G₃	110.90	123.54	121.03	122.39	119.47
G₄	117.78	119.04	120.54	119.81	119.29
Mean	99.36	104.84	103.86	103.84	
	SEm±			CD (0.05)	
G	0.74			2.13	
K	0.74			2.13	
(G×K)	1.48			4.27	

NOTE: G₁-0 ppm, G₂-100 ppm, G₃- 200 ppm, G₄-300 ppm, K₁-40 kg/ha, K₂-50 kg/ha, K₃-60 kg/ha, K₄-70 kg/ha

Table 6 Effect of GA₃ and potassium on Root-Shoot ratio (weight bases) in red radish (SJRR-01)

Root-shoot ratio					
Treatments	K₁	K₂	K₃	K₄	Mean
G₁	1.28	1.40	1.42	1.43	1.38
G₂	1.76	1.98	1.84	1.91	1.87
G₃	1.96	2.10	2.05	2.05	2.04
G₄	1.99	2.00	2.09	2.02	2.02
Mean	1.75	1.87	1.85	1.85	
	SEm±			CD (0.05)	
G	0.01			0.04	
K	0.01			0.04	
(G×K)	0.03			0.09	

Table 7 Effect of GA₃ and potassium on Fresh root shoot yield (q/ha) in red radish (SJRR-01)

Fresh root shoot yield (q/ha)					
Treatments	K₁	K₂	K₃	K₄	Mean
G₁	71.83	75.26	72.50	72.27	72.96
G₂	84.95	93.38	92.20	91.96	90.62
G₃	87.03	103.58	102.19	102.19	98.75
G₄	94.95	96.01	100.20	100.29	97.86
Mean	84.69	92.06	91.77	91.68	
	SEm±			CD (0.05)	
G	1.15			3.31	
K	1.15			3.31	
(G×K)	2.29			6.62	

NOTE: G₁-0 ppm, G₂-100 ppm, G₃- 200 ppm, G₄-300 ppm, K₁-40 kg/ha, K₂-50 kg/ha, K₃-60 kg/ha, K₄-70 kg/ha