

IMPLICATIONS OF (DAY AND NIGHT) TEMPERATURE VARIATIONS IN RICE PRODUCTION

(Impact of temperature variations on rice production)

Abstract

Climatic variables **effect**(**Affect**) growth and development of the crop differently at different growth stages. Rate of development of a crop depends on the climate. Temperature is the primary factor which influences the growth and productivity of crops. Higher or lower temperature than the optimum leads to improper development of a crop and it affects the phenological stages of rice. Extreme temperature may lead to early maturity of the crop and lower temperature lead to prolonged maturity of the crop which in both the cases leads to decrease in yield. Moderate increase in temperature will not have severe implications, but increase in average temperature by 2⁰ C will be likely to have strong negative effects on rice crop production. In several studies, it has been reported that, high night temperature decreases the yield potential of rice. Whereas, respiratory losses have been shown to increase as a result of higher day temperature, leaf conductance and net assimilation rates during the day were reported to be higher.

Key words: Rice, temperature, phenology, spikelet, grain yield.

Introduction

Rice plays an important role in food security of many countries as it is the staple food for more than half of the world's population. It is the staple food for over 65 per cent population in India. (Kumar *et al.*, 2018). Globally rice is cultivated in an area of 167.13 million hectares with a production of 782 million tonnes and with a productivity of 4,679 kg ha⁻¹ (Anon., 2019). In India rice is grown in an area of 44.15 m ha with 116.4 million tonnes of production and 2,638 kg ha⁻¹ productivity (Anon., 2020). 90 per cent of the world rice comes from Asia of which 85 per cent of production is used for human consumption and it deserves a special status among cereals as world's most important wetland crop (Rajeshkumar *et al.*, 2019). Regarding to human nutrition and caloric intake, it is considered to be the most significant cereal crop supplying over one-fifth of the calories consumed by humans globally (Smith and Bruce). Though for many years used as a model plant, however, in the last decades, the unprecedented increase in temperature extremities exposed a wide series of variances related with heat stress.

In different regions of the world, its harsh influences on different crops have been noticeably apparent (Amanullah *et al.*, 2017).

Currently, rice production is facing multiple challenges such as water stress, insect pest infestation, disease attack, which delay its planting and as a result barricade its sustainable production. Future crop yields will be influenced by complex interactions between the effects of increases in atmospheric concentrations of CO₂ (Baker *et al.*, 1990) and trace gases such as ozone (Maggs *et al.*, 1998) as well as the effects of temperature increases brought about by climate change (Rosenzweig and Parry, 1994). Global mean surface air temperature increased by nearly 0.5°C in the 20th century and is projected to further increase by 1.5 to 4.5°C in this century (Houghton *et al.*, 1995). In the past century, daily minimum night time temperature increased at a faster rate than daily maximum temperature in association with a steady increase in atmospheric greenhouse gas concentrations (Karl *et al.*, 1991; Easterling *et al.*, 1997). Experimental evidences have repeatedly repressed that a short period of high temperature (owing to climatic fluctuations) had greater negative impacts on grain than continuous mild stress (Reidsma *et al.*, 2010). A 25-year weather data report from International Rice Research Institute, Philippines has indicated greater increase in night time temperature (1.13°C) over day time temperature (0.35°C) (Peng *et al.*, 2005). World rice production must increase by 1 per cent annually to meet the growing demand for food that will result from population growth and economic development (Rosegrant *et al.*, 1995). Most of this increase must come from greater yields on existing cropland to avoid environmental degradation, destruction of natural ecosystems, and loss of biodiversity (Cassman, 1999; Tilman *et al.*, 2002). Achieving greater yields depends on increasing total crop biomass, because there is little scope to further increase the proportion of that biomass allocated to grain (Evans and Fischer, 1999). Total crop biomass is determined mainly by crop photosynthesis and respiration losses, both of which are sensitive to temperature (Yoshida, 1981).

Growth and grain yield of rice depends on the combined effect of genetic characteristics of the variety and environmental conditions of the area where the variety is grown (De Datta, 1970). Among all the environment factors, temperature plays a major role in rice yield. The optimum temperature for the normal development of rice plant ranges from 27°C to 32°C (Yin *et al.*, 1996). High temperature affects almost all the growth stages of rice which includes from emergence to ripening and harvesting (Yoshida, 1978; Yoshida, 1981, Satake & Yoshida 1978; Morita *et al.*, 2004). However, flowering (anthesis and fertilization) and booting

(microsporogenesis) are considered to be the most susceptible stages to temperature in rice (Satake & Yoshida, 1978; Farrell *et al.*, 2006). Temperatures higher than the optimum induces floret sterility and thus decreases rice yield (Nakagawa *et al.*, 2003). Spikelet sterility is greatly increased at temperatures higher than 35°C (Satake & Yoshida, 1978; Matsui *et al.*, 1997) which is the critical maximum temperature in rice. Night temperature of less than 19°C is the critical low temperature for inducing grain sterility in rice (Abeyasiriwardena *et al.*, 2002). The unimpeded growth of greenhouse gas emissions is raising the earth's temperature. The consequences include melting glaciers, more precipitation, more and more extreme weather events, and shifting seasons. The accelerating pace of climate change, combined with global population and income growth, threatens food security everywhere. Agriculture is extremely vulnerable to climate change. Higher temperatures eventually reduce yield of desirable crops while encouraging weed and pest proliferation. Adaptation forecasts that by 2050 rice prices will increase between 32 and 37 per cent as a result of climate change. They also show that rice productivity will reduce by 14 per cent in South Asia, 10 per cent in East Asia and the Pacific, and 15 per cent in Sub-Saharan Africa (Nelson *et al.*, 2009). The increased global temperature leads to rainfall deviations which can produce even more devastating results in crop growth (Ranatunge *et al.*, 2003; Dore, 2005; Lobell *et al.*, 2007; Auffhammer *et al.*, 2012).

Rice phenology is strongly influenced by the environmental conditions in a certain range, an increase in temperature leads to decrease in vegetative phase. Low thermal stress occurs in seedling and heading-flowering stages, whereas high thermal stress during the end of growing season causes the sterility of seeds. On the other hand, the risk of cold stress during the rice growth leads to reduction in yield (Shimono, 2011). With increasing temperature, maturity stage will decline (Baker *et al.*, 1990). By increasing in seasonal temperatures, it will cause increased risk of drought and the speed of photosynthesis will be limited (Tubiello *et al.*, 2007). Roberts (1943) suggested that the temperature during the night rather than the day largely determines the response of plants to temperature. Climate models foresee that a relatively greater increase in night time temperatures than daytime temperatures will occur because of less radiant heat loss due to increased cloudiness (Alward *et al.*, 1999). Over the past century the increases in global daily minimum temperatures were more than those twice that of daily maximum temperatures (Easterling *et al.*, 1997). Evidence of historical yields of rice (Peng *et al.*, 2004) and wheat (Lobell *et al.*, 2005) shows that cereal yield was strongly correlated with minimum nighttime temperatures rather than daytime maximum temperatures (Chen *et al.*, 2013).

Temperature is considered to be the most critical yield determinant of rice and high temperature in the lower elevation of the tropics and lower temperature in the temperate regions adversely affect the crop (Yoshida, 1981). Moderate warming in the coming decades, will possibly imply a net negative impact on yield of rice (Welch *et al.*, 2010) and the productivity may decrease by 4.17 per cent (Upadhyay, 2012). The rise in atmospheric temperature causes detrimental effects on growth, yield, and quality of the rice crop by affecting its phenology, physiology, and yield components (Singh, 2001 and Sheehy *et al.*, 2005). It can affect the crop at all stages of development, particularly during flowering when it causes spikelet sterility. It increases plant respiration, affects photosynthesis and shortens the grain filling period, all of which leads to lower productivity (Peng *et al.*, 2004). On the other hand, relatively low temperature and high solar radiation during the reproductive stage had remarkable effects on increasing spikelet number and hence grain yield (Yoshida and Parao, 1976). Rise in the night temperature is another reason for decline in historical rice yield in addition to the observed changes in the climatic conditions (Kumar *et al.*, 2004) particularly during the ripening phase of the crop. Night-time warming at the end of the growing season had a greater impact on yield than the increase in the maximum temperature and changes in rainfall characteristics (Auffhammer *et al.*, 2012; Auffhammer *et al.*, 2006).

Although an increase in global temperature has been well documented, a greater increase in night-time compared with day-time temperatures has been highlighted recently (Sillmann *et al.*, 2013). This differential increase in day and night temperature will result in a reduced diurnal temperature range, which has been shown to affect crop growth and development (Yin *et al.*, 1996; Peng *et al.*, 2004; Bahuguna and Jagadish, 2015).

Impact of day and night temperature variations on rice phenology

In a study conducted at IRRI, Philippines, it was noticed that increase in early night temperature from 24 to 30 or to 35 °C advanced start of anthesis by 2 or 2.5 hours respectively in cultivar WAB56-104, but no effect was detected in the other two cultivars. Late high night temperature advanced the start of anthesis in cultivar N22 by 30 min at both 30 and 35 °C, while a delay and advance in flowering by 30 min at 30 and 35 °C, respectively, was recorded in WAB56-104, with no change in CG14. CG14 opened its first flower before 0530 hours consistently and because observations started at 0500 h and every 30 min 23 thereafter no high night temperature effect could be detected (Coast *et al.*, 2010). High day temperature damage leaf gas exchange properties during the vegetative stage and even a short period of heat stress can cause

significant increases in the abortion of floral buds and opened flowers during the reproductive stage (Guilioni *et al.*, 1997). Increased night temperatures significantly increased the rate of senescence and maturity was achieved in those plants exposed to high night temperature 15 days earlier than in the normal temperature chamber. The effects of high minimum temperatures increased the rate of senescence and decreased the ability of the plant to efficiently produce grain (Hatfield and Prueger, 2015). In all genotypes tested (Nagina 22 (N22; heat tolerant), IR64 (heat susceptible), heat tolerant IR64 near-isogenic line (HT NIL) (Ye *et al.*, 2012), and two hybrids, H2 (private company hybrid) and H5 (International Rice Research Institute hybrid breeding programme)), the maximum (C_m) and mean (C) grain-filling rates were higher with high night temperature than for the control, whereas the time taken to reach the maximum grain-filling rate (t_m) and total grain-filling duration (t_e) were shortened by HNT compared with the control (Shi *et al.*, 2017). Results of a study conducted at IRRI, Philippines indicated that night temperature promoted flowering more efficiently than day temperature in three cultivars only, whereas the opposite effect was found in most other cultivars (Yin *et al.*, 1996).

Impact of day and night temperature variations on growth parameters

Three cultivars of rice (BR 3, BR 6 and BR 8) were studied in Bangladesh, under natural day length with normal night temperature, short day (10 hours) and low night temperature (16⁰C) and long day (16 hours) and high night temperature (35⁰C). The growth parameters like CGR, RGR, NAR and LAI increased in BR 8 while only LAI increased in BR 3 and both RGR and LAI were found to be increased in BR 6 under long day with high night temperature. The plant height and panicle length were found to be reduced under long day with high night temperature in all cultivars (Alam *et al.*, 1985). In a study conducted in China, it was found that plant height, leaf area, and shoot dry weight were greater in high night temperature (HNT- 27⁰C) than in CK (22⁰C) (Chen *et al.*, 2013). In a study conducted in Germany, large variation in leaf area was observed, but it was not significant in relation to day temperature (DT), night temperature (NT), or their combined effects, represented by the temperature treatments (TR). The marginal means according to NT were in the range of 436.16 ± 55.93 cm² at 25⁰C, and 454.75 ± 61.96 cm² at 30⁰C. The magnitude between DT and NT was not significant, though the highest LA was in TR D30 C/N20 C. Rice plant height, tiller number, dry mass, and all of its constituents (leaf, sheath, and root mass) were unaffected by DT, NT or by TR (Johnson, 2018). With high day temperature, the shoot mass fraction increased, whereas the root mass fraction decreased.

Specific leaf area increased at high night temperatures and led, along with the high leaf mass fraction at high night humidities, to higher growth rates. The results showed the importance of considering relative air humidity when focusing on plant responses to temperature, and strongly suggest that under asymmetric day and night temperature increases in the future, biomass partitioning rather than biomass itself will be affected (Stuerz and Asch, 2019). Plant leaf growth was increased by high night temperature as well as leaf area ratio (LAR) and specific leaf area (SLA), while the effect on organ weight and total dry matter was less clear, and varied between plant functional groups. As a result, it was concluded that complexities and challenges remain when seeking general patterns of plant growth in response to high night temperature (Jing *et al.*, 2016). Zhang *et al.*, 2013 found that plant height of rice was significantly increased by high night temperature in all the four experiments except one season (2011 DS) which might be due to the reason that elongated rice stems retained more photosynthate than the ears did under high night temperature (HNT) (Cheng *et al.*, 2009). Panicle number was not significantly affected between high night temperature and low night temperature (LNT). Biomass in HNT decreased by 9.2 per cent, 5.3 per cent, 5.5 per cent and 4.1 per cent than LNT in the four consecutive seasons, because HNT received lower biomass due to its lower CGR. Rice biomass production was determined by the balance between net photosynthesis rate and night respiration (Sakai *et al.*, 2001). Total dry weight is significantly greater in the HNT grown plants than in the LNT grown plants, because plants in HNT had an increase of leaf area and tiller number (Kanno *et al.*, 2009).

Impact of day and night temperature variations on yield and yield parameters

Fertile tillers, fertile grains and 1000-grain weight of three cultivars of rice (BR 3, BR 6 and BR 8) were found to be increased under long day with high night temperature as compared to other treatments (Alam *et al.*, 1985). In an experiment conducted at Tamil Nadu, India, empirically it was observed from the differences between the two seasons (*Kharif* and Summer) that on an average 1.2°C increase in night temperature had reduced the head rice recovery by 24.9 per cent whereas 4.4 °C increase in day temperature is required to produce the same effect on head rice recovery as that of night temperature. Hence it was concluded that very small magnitude increase in night temperature affects the head rice recovery negatively (Bhaskaran and Sebastian, 2017). In a study conducted at Bangladesh, it was revealed that maximum temperature is statistically significant and negatively affect the yield of all three rice crops (Aus,

Aman and *Boro*). In contrast, minimum temperature is highly significant and have positive impact on the yield of *Boro* rice only (Chowdhury and Khan, 2015). High night temperatures (22/34°C, day/night) were more harmful to grain weight in rice than high day temperatures (34/22°C) and control conditions (22/22°C) at optimum temperature (Morita *et al.*, 2004). From the report of a study conducted at Jharkhand, India, it was found that minimum temperature during vegetative stage in variety Vandana and flowering stage in rice varieties BVD-109 and BVD111 had negative correlation with rice yield (Kumari *et al.*, 2017). In a field experiment carried out in China, from rice heading stage to maturity (6 August to 19 September 2006 and 7 August to 20 September 2007), mean air temperatures of daytime (from 07:00 to 19:00 h) and nighttime (from 19:00 to 07:00 h) under high air temperature (HAT) were approximately 2.0 and 0.3 °C higher than those under natural air temperature (NAT). HAT treatment was conducted from rice heading stage, hence significant difference was not observed in effective panicles per m² between NAT and HAT. When compared to NAT, significant decreases in number of filled grains per panicle (25.70 %), seed-setting rate (22.09 %), 1000-grain weight 5.17 (%) and grain yield (31.24 %) under HAT were observed (Liu *et al.*, 2013). The night temperature (NT) had no effect on number of productive tillers, main-stem panicle length or number of primary branches per panicle. The decreased plant grain yield under high night temperature was due to increased spikelet sterility and decreased grain length and width (Mohammed and Tarpley, 2010). Night temperatures greater than 29° C can increase spikelet sterility in rice with a subsequent reduction in seed-set and grain yield (Satake and Yoshida 1978; Ziska *et al.*, 1996; Jagadish *et al.*, 2007). From the results of an experiment conducted at IRRI, Philippines, it was observed that there was a strong negative linear relationship between above ground total biomass (grain and straw) at maturity and minimum temperature over a very narrow range of minimum temperature (<2°C) in the dry season. Biomass production decreased by ≈10 per cent for each 1°C increase in minimum temperature. There was no significant relationship between crop growth duration and minimum temperature. Therefore, the reduction in biomass production with warm nights was not associated with a decrease in growth duration. There was a tight negative linear relationship between spikelets per m² and minimum temperature in the dry season. Panicles per m² were negatively related to minimum temperature but not to radiation or maximum temperature. Other yield components such as spikelets per panicle, grain-filling percentage, and grain weight were not related to minimum or maximum temperature or radiation. There was a significant negative relationship between minimum temperature and harvest index in the dry season. Therefore, grain yield decreased by at least 10 per cent for each 1°C increase in growing-season minimum temperature. Because

the increase in mean minimum temperature was >3-fold greater than the increase in mean maximum temperature, it was concluded that rice grain yield declined by ≈ 15 per cent for each 1°C increase in growing-season mean temperature (Peng *et al.*, 2004). Besides reducing carbon availability and its rate of translocation to seeds, increases in night temperature induced yield losses through direct impacts on complex reproductive and seed maturity processes, resulting in not only quantitative grain losses but also effects on grain quality and composition. One mechanism that enables yield penalties through high night temperature operates by lowering seed set as a result of poor pollination (Sadok and Jagadish, 2020). This effect has been shown to be mediated by increased accumulation of reactive oxygen species (ROS) leading to increased membrane damage, ultimately leading to lower pollen viability (Prasad *et al.*, 2011; Luria *et al.*, 2019). Exposure to increasing temperatures under high day time temperature, high night time temperature or combined high day and night temperature under chamber (Yamakawa *et al.*, 2007; Cao *et al.*, 2016) or field (Shi *et al.*, 2013; Rehmani *et al.*, 2014; Bahuguna *et al.*, 2017) conditions during grain filling impairs grain growth, leading to poor seed-set and reduced single-grain weight. The high night temperature was strongly and negatively correlated with grain yield in all the four experiments. Grain yields in HNT decreased by 16.7 per cent, 9.1 per cent, 9.6 per cent and 8.0 per cent than LNT in the four seasons. Among the yield components, significant differences were observed in both grain weight and harvest index in the four experiments (Zhang *et al.*, 2013). Seshu and Cady (1984) predicted that rice yield would decrease by 41 g m^{-2} when the average minimum temperature raised from 22 to 23°C .

Conclusion

Production and productivity of rice depends on agrometeorological parameters. By exploring the rice-temperature relation, it can be concluded that, optimum (value of optimum temperature for rice) day and night temperatures are required for the better growth and productivity of crop. From the above reviews, it can be conclude that cool night temperature causes spikelet sterility and hence reduces the yield. Rice crop might get benefit from day and night temperature variations (high or low) during the vegetative growth stages where it is not the same during reproductive stages. Day temperature independently or in combination with high night temperature alters the plant processes and hence decreasing the grain yield and quality.

References

- Abeyasiriwardena, D. S. D. Z., Ohba, K. and Maruyama, A., 2002, Influence of temperature and relative humidity on grain sterility in rice. *Journal of the National Science Foundation of Sri Lanka*. 30: 33 - 41.
- Alam, S. M. M., Islam M.T. and Muhsi A.A.A., 1985, Effect of light and night temperature on some cultivars of rice (*Oryza sativa* L.). *Indian J. Plant Physiol.*, Vol. XXVIII (4): pp. 385-394.
- Alward, R. D., Detling, J. K. and Milchunas, D. G., 1999, "Grassland vegetation changes and nocturnal global warming," *Science*, 283 (5399): pp. 229–231.
- Amanullah, Fahad, S., Anwar, S., Baloch, S. K., Saud, S., Alharby, H., Alghabari, F. and Ihsan, M. Z., 2017, Rice Crop Responses to Global Warming: An Overview. Rice - Technology and Production. *Intech open*. Chapter I, 1-10.
- Anonymous, 2019, Area, production and average yield, Directorate of Economics and Statistics, Department of Agriculture and Cooperation report, New Delhi, available on the website: [http:// www.agricoop.nic.in](http://www.agricoop.nic.in)
- Anonymous, 2020, Area, production and average yield, Food and Agricultural Organization, available on the website: [http:// www.indiastats.com](http://www.indiastats.com).
- Auffhammer, M., Ramanathan, V. and Vincent, J. R., 2006, Integrated model shows that atmospheric brown clouds and greenhouse gases have reduced rice harvests in India. *Proceedings of the National Academy of Sciences*. 103(52): 19668-19672.
- Auffhammer, M., Ramanathan, V. and Vincent, J. R., 2012, Climate change, the monsoon, and rice yield in India. *Climatic Change*. 111(2): 411-424.
- Bahuguna, R. N. and Jagadish, S. V. K., 2015, Temperature regulation of plant phenological development. *Environmental and Experimental Botany*. 111: 83–90.
- Bahuguna, R. N., Solis, C. A., Shi, W., Jagadish, S. V. K., 2017, Post flowering night respiration and altered sink activity account for high night temperature-induced grain yield and quality loss in rice (*Oryza sativa* L.). *Physiologia Plantarum*. 159: 59–73.
- Baker, J. T., Allen, L. H., Boote, K. J., Jones, P. and Jones, J. W., 1990, Developmental responses of rice to photoperiod and carbon dioxide concentration. *Agric. For. Meteorol.*, 50: 201-210.
- Bhaskaran, M. and Sebastian, A., 2017, Understanding effects of high temperature stress on head rice recovery in rice. *International Journal of Current Microbiology and Applied Sciences*. 6(11): 5476-5481.

- Cassman, K. G., 1999, Ecological intensification of cereal production system: Yield potential, soil quality and precision agriculture. *Proc. Natl. Acad. Sci. USA*. 96: 5952–5959.
- Cao, Y., Chen, Y., Chen, M., Wang, Z., Wu, C., Bian, X., Yang, J. and Zhang, J., 2016, Growth characteristics and endosperm structure of superior and inferior spikelets of *indica* rice under high-temperature stress. *Biologia Plantarum*. 60: 532–542.
- Chen, S., Zhang, X., Zhao, X., Wang, D., Xu, C., Ji, C. and Zhang, X., 2013, Response of Rice Nitrogen Physiology to High Night time Temperature during Vegetative Stage. *The Scientific World Journal*. 1-10.
- Cheng, W., Sakai, H. and Yagi K., 2009, Interactions of elevated (CO₂) and night temperature on rice growth and yield. *Agr. Forest. Meteorol.*, 149: 51-58.
- Chowdhury, I. U. A. and Khan, M. A. E., 2015, The impact of climate change on rice yield in Bangladesh: a time series analysis. *Russian Journal of Agricultural and socio-economic sciences*. 4(40).
- Coast, O., Ellis, R. H., Murdoch, A. J., Quiñones, C. and Jagadish, K. S. V., 2015, High night temperature induces contrasting responses for spikelet fertility, spikelet tissue temperature, flowering characteristics and grain quality in rice. *Functional Plant Biology*. 42 (2): pp 149-161.
- De Datta, S. K., 1970, The environment of rice production in tropical Asia. In Rice production manual, International Rice Research Institute, Philippines. pp 53-67.
- Dore, M. H., 2005. Climate change and changes in global precipitation patterns: what do we know? *Environment international*. 31(8): 1167-1181.
- Easterling, D. R., Horton, B., Jones, P. D., Peterson, T. C., Karl, T. R., Parker, D. E., Salinger, M. J., Razuvayev, V., Plummer, N. and Jamason, P., 1997, Maximum and minimum temperature trends for the globe. *Science*, 277: 364–367.
- Evans, L. T. and Fischer, R. A., 1999, Yield potential: Its definition, measurement and significance. *Crop Sci.*, 39: 1544–1551.
- Farrell, T. C., Fox, K. M., Williams, R. L. and Fukai, S., 2006, Genotypic variation for cold tolerance during reproductive development in rice: screening with cold air and cold water. *Field Crops Research*. 98, 178 – 194.
- Guilioni, L., Wery, J. and Tardieu, F., 1997, Heat stress-induced abortion of buds and flowers in pea: Is sensitivity linked to organ age or to relations between reproductive organs? *Ann. Bot.* 80: 159–168.
- Hatfield, J. L. and Prueger, J. H., 2015, Temperature extremes: Effect on plant growth and development. *Weather and Climate Extremes*. 10: 4-10.

- Houghton, J. T., Filho, L. G. M., Bruce, J., Lee, H., Callender, B. A., Haites, E., Harris, N. & Maskell, K., 1995, Climate Change. Intergovernmental Panel on Climate Change: The Science of Climate Change, eds (Cambridge Univ. Press, Cambridge, U.K.).
- Jagadish, S. V. K., Craufurd, P. Q. and Wheeler, T. R., 2007, High temperature stress and spikelet fertility in rice (*Oryza sativa* L.). *J. Exp. Bot.*, 58: 1627-1635.
- Jing, P., Wang, D., Zhu, C. and Chen, J., 2016, Plant Physiological, Morphological and Yield-Related Responses to Night Temperature Changes across Different Species and Plant Functional Types. *Front. Plant Sci.*, 7: 835.
- Johnson, K., 2018, Effects of Day and Night Temperature on Rice Photosynthesis. M. Sc Thesis, Institute of Agricultural Science in the Tropics. University of Hohenheim, Stuttgart, Germany.
- Kanno, K., Tadahiko and Makino, A., 2009, High night temperature stimulates photosynthesis, biomass production and growth during the vegetative stage of rice plants. *Soil Sci Plant Nutr.*, 55:124-431.
- Karl, T. R., Kukla, G. and Razuvayev, V. N., 1991, Global warming: Evidence for asymmetric diurnal temperature change. *Geophys. Res. Lett.* 18: 2253–2256.
- Kumar, K. K., Kumar, K. R., Ashrit, R. G., Deshpande, N. R. and Hansen, J. W., 2004, Climate impacts on Indian agriculture. *International Journal of Climatology*. 24(11): 1375–1393.
- Kumari, P., Kumar, V. P., Kumar, R., Wadood, A. and Tirkey, D. A., 2017, Effect of weather on grain yield of direct seeded upland rice varieties in Jharkhand, India. *Indian J. Agric. Res.*, 51(6): 562-567.
- Kumar, G. S., Ramesh, K., Subrahmaniyan and Ravi, V., 2018, Effect of sprinkler irrigation levels on the performance of rice genotypes under aerobic condition. *Int. J. Curr. Microbiol. App. Sci.*, 7(3):1848-1852.
- Liu, Q., Wu, X., Li, T., Ma, J. and Zhou, X., 2013, Effects of elevated air temperature on physiological characteristics of flag leaves and grain yield in rice. *Chilean Journal of Agricultural Research*. 73(2): 85-90.
- Lobell, D. B., Ortiz-Monasterio, J. I., Asner, G. P., Matson, P. A., Naylor, R. L. and Falcon, W. P., 2005, Analysis of wheat yield and climatic trends in Mexico. *Field Crops Research*, 94(2-3): pp.250– 256.
- Lobell, D. B., Cahill, K. N. and Field, C. B., 2007, Historical effects of temperature and precipitation on California crop yields. *Climatic Change*. 81(2): 187-203.
- Luria, G. *et al.*, 2019, Direct analysis of pollen fitness by flow cytometry: implications for pollen response to stress. *Plant J.*, 98: 942–952.

- Maggs, R. and Ashmore, M. R., 1998, Growth and yield responses of Pakistan rice (*Oryza sativa* L.) cultivars to O₃ and NO₂. *Environ. Pollut.* 103: 159–170.
- Matsui, T., Namuko, O. S., Ziska, L. H. and Horie, T., 1997, Effect of high temperature and CO₂ concentration on spikelet sterility in indica rice. *Field Crops Research*. 51, 213 - 219.
- Mohammed, A. R. and Tarpley, L., 2010, High night temperature and plant growth regulator effects on spikelet sterility, grain characteristics and yield of rice (*Oryza sativa* L.) plants. *Can. J. Plant Sci.*, 91: 283-291.
- Morita, S. Shiratsuchi, H., Takahashi, J. and Furta, K., 2004, Effect of high temperature on ripening in rice plants: analysis of the effect of high night and high day temperatures applied to the panicle and other parts of the plant. *Japanese Journal of Crop Science*. 73, 77-83.
- Nakagawa, H, Horie, T. and Matsui, T., 2003, Effects of climate change on rice production and adaptive technologies. In: Shah, F., Huang, J., Cui, K., Nie, L., Shah, T., and Wang, K. (Eds.) Climate change and agriculture paper. Impact of high temperature stress on rice plant and its traits related to tolerance. (2011). *Journal of Agricultural Science*. Cambridge University Press. pp 1-12.
- Nelson, G. C., Rosegrant, M. W., Koo, j., Robertson, R. D., Sulser, T., Zhu, T., Ringler, C., Msangi, S., Palazzo, A., Batka, M., Magalhaes, M., Valmonte-Santos, R., Ewing, M. and Lee, D. R., 2009, Climate change: Impact on agriculture and costs of adaptation. The International Food Policy Research Institute (IFPRI) Report. Available on the website: [http:// www.ifpri.org](http://www.ifpri.org)
- Peng, S., Huang J., Sheehy J. E., Laza R. C., Visperas R. M., Zhong X., Centeno G. S., Khush G. S. and Cassman K. G., 2004, Rice yields decline with higher night temperature from global warming. *Proc. Natl. Acad. Sci.*, U S A. 101: 9971–9975.
- Prasad, P. V. V. and Djanaguiraman, M., 2011, High night temperature decreases leaf photosynthesis and pollen function in grain sorghum. *Funct. Plant Biol.*, 38: 993.
- Rajeshkumar, A., Ramdass, S. and Thirumeninathan, S., 2019, Influence of drip fertigation in aerobic rice production. *Int. J. Agri. Sci.*, 11(4), pp:8824-8827.
- Ranatunge, E., Malmgren, B. A., Hayashi, Y., Mikami, T., Morishima, W., Yokozawa, M. and Nishimori, M., 2003, Changes in the Southwest Monsoon mean daily rainfall intensity in Sri Lanka: relationship to the El Niño–Southern Oscillation. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 197(1–2): 1-14.
- Rehmani, M. I. A., Wei, G., Hussain, N., Ding, C., Li, G., Liu, Z. and Ding, Y., 2014, Yield and quality responses of two indica rice hybrids to post-anthesis asymmetric day and night

- open-field warming in lower reaches of Yangtze River delta. *Field Crops Research*. 156: 231–241.
- Reidsma, P., Ewert F., Lansink A. O. and Leemans, R., 2010, Adaptation to climate change and climate variability in European agriculture: The importance of farm level responses. *Eur. J. Agron.*, 32: 91–102
- Roberts, R. H., 1943, The role of night temperature in plant performance. *Science.*, 98: 265.
- Rosegrant, M. W., Sombilla, M. A. and Perez, N., 1995, Food, Agriculture and the Environment Discussion Paper No. 5 (International Food Policy Research Institute, Washington, DC).
- Rosenzweig, C. & Parry, M. L., 1994, Potential impact of climate change on world food supply. *Nature*. 367: 133–138.
- Sadok, W. and Jagadish, 2020, The Hidden Costs of Nighttime Warming on Yields. *Trends in Plant Science*. 25(7): 644-651.
- Sakai, H., Yagi, K. and Kobayashi, K., 2001, Rice carbon balance under elevated CO₂ . *New Phytologist*. 150: 241-249.
- Seshu, D. V. and Cady, F. B., 1984, Response of rice to solar radiation and temperature estimated from international yield traits. *Crop Sci.*, 24: 649-654.
- Satake, T. and Yoshida, S., 1978, High temperature induced sterility in indica rice at flowering. *Japanese Journal of Crop Science*. 47, 6 - 17.
- Sheehy, J. E., Elmido, A., Centeno, G. and Pablico, P., 2005, Searching for new plant for climate change. *J. Agrometeorol.*, 60: 463–468.
- Shi, W., Muthurajan, R., Rahman, H., Selvam, J., Peng, S., Zou, Y. and Jagadish, S. V. K., 2013, Source-sink dynamics and proteomic reprogramming under elevated night temperature and their impact on rice yield and grain quality. *New Phytologist*. 197: 825–837.
- Shi, W., Yin, X., Struik, P. C., Solis, C., Xie, F., Schmidt, R. C., Huang, M., Zou, Y., Ye, C. and Jagadish, S. V. K., 2017, High day- and night-time temperatures affect grain growth dynamics in contrasting rice genotypes. *Journal of Experimental Botany*. 68(18): pp. 5233–5245.
- Shimono, H., 2011. Earlier rice phenology as a result of climate change can increase the risk of cold damage during reproductive growth in Northern Japan. *Agric. Ecosyst. Environ.*, 144: 201-207.
- Sillmann, J., Kharin, V. V., Zhang, X., Zwiers, F. W., Bronaugh, D., 2013, Climate extremes indices in the CMIP5 multimodel ensemble part: 1. Model evaluation in the present climate. *Journal of Geophysical Research: Atmospheres*. 118: 1716–1733.

- Singh, S., 2001, Growth, yield and biochemical response of rice genotype to low light and high temperature humidity stress. *Oryza*. 37(1): 35–38.
- Smith and Bruce, D., The Emergence of Agriculture. Scientific American Library, A Division of HP HLP, New York, ISBN 0-7167-6030-4.
- Stuerz, S. and Asch, F., 2019, Responses of Rice Growth to Day and Night Temperature and Relative Air Humidity—Dry Matter, Leaf Area, and Partitioning. *Plants*. 8, 521. 1-12.
- Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R. and Polasky, S., 2002, Agricultural sustainability and intensive production practices. *Nature*. 418: 671–677.
- Tubiello, F. N., Soussana, J. F. and Howden, S. M., 2007, Crop and pasture response to climate change. *Proc. Natl. Acad. Sci. USA*, 104: 19686-19690.
- Upadhyay, J., 2012, Climate Change and its Impact on rice productivity in Assam. Social Science Research Network. Retrieved from [http://ssrn.com/abstract = 2172183](http://ssrn.com/abstract=2172183) or [http://dx.doi.org/ 10.2139/ssrn.2172183](http://dx.doi.org/10.2139/ssrn.2172183).
- Welch, J. R., Vincent, J. R., Auffhammer, M., Moya, P. F., Dobermann, A. and Dawe, D., 2010, Rice yields in tropical/subtropical Asia exhibit large but opposing sensitivities to minimum and maximum temperatures. *Proc. Natl. Acad. Sci. U.S.A.* 107: 14562.
- Yamakawa, H., Hirose, T., Kuroda, M. and Yamaguchi, T., 2007, Comprehensive expression profiling of rice grain filling-related genes under high temperature using DNA microarray. *Plant Physiology*. 144: 258–277.
- Yin, X., Kroff, M. J. and Goudriann, J., 1996, Differential effects of day and night temperature on development to flowering in rice. *Annals of Botany*. 77: 203 - 213.
- Yoshida, S. and Parao, F. T., 1976, Climatic influence on yield and yield components of lowland rice in the tropics. *Climate and Rice*. International Rice Research Institute, Los Banos, Philippines. pp. 471-494.
- Yoshida, S., 1978, Tropical climate and its influence on rice. IRRI Research Paper Series 20. The Philippines: IRRI.
- Yoshida, S., 1981, Fundamentals of rice crop science. The Philippines: IRRI.
- Zhang, Y., Tang, Q., Peng, S., Zou, Y., Chen, S., Shi, W., Qin, J. and Laza, M. R. C., 2013, Effects of high night temperature on yield and agronomic traits of irrigated rice under field chamber system condition. *Australian Journal of Crop Science*. 7(1): 7-13.
- Ziska, L. H., Manalo, P. A. and Ordonez, R. A., 1996, Intraspecific variation in the response of rice (*Oryza sativa* L.) to increased CO₂ and temperature: growth and yield response of 17 cultivars. *J. Exp. Bot.*, 47: 1353-1359.