

# A Narrative Review On Acute Cholecystitis in Pregnancy: Current Update on Management

## Abstract

Acute cholecystitis in pregnancy is one of the common non obstetric causes for acute abdominal pain. The management can be divided into conservative treatment and surgical treatment. Conservative treatment involves the use of intravenous fluids and antibiotics, but patients are prone for developing recurrence of symptoms. Surgical treatment involves performing a cholecystectomy either as an open or laparoscopic method. As there is no consensus on the management of acute cholecystitis in pregnancy, we have conducted this review article to look at the role of conservative treatment, the role of laparoscopic cholecystectomy and when is the best time to perform this surgery. The role of percutaneous cholecystostomy is also mentioned in this review.

**Keywords-**Acute cholecystitis in pregnancy, gallstone disease in pregnancy, conservative treatment of acute cholecystitis in pregnancy, laparoscopic cholecystectomy in pregnancy, cholecystectomy in pregnancy.

## Introduction

Acute cholecystitis accounts for the second most common cause for non-obstetric abdominal pain in pregnant patients. The incidence of acute cholecystitis in pregnancy is about 0.2 to 0.5 per 1,000 pregnancies. The global prevalence of gallstone disease in pregnancy is about 3.6%. During pregnancy the elevated serum estrogen and progesterone predispose to gallstone formation. Estrogen is associated with increased viscosity of bile and cholesterol crystal aggregation. Progesterone is associated with gallbladder smooth muscle relaxation and bile stasis.(1–3)

The clinical presentation is the same as for non-pregnant women, which includes pain over the right hypochondrium and associated symptoms like nausea and vomiting. There may also be symptoms of dyspepsia and abdominal distension after eating. Diagnosis is usually confirmed by using blood investigations like full blood count, liver function test and serum amylase and imaging with an ultrasound of the abdomen. Ultrasound has a high sensitivity and specificity in diagnosing acute cholecystitis in pregnancy. (4)

The anatomical and physiological changes that occur during pregnancy make the diagnosis and treatment of acute cholecystitis in pregnancy difficult. The position of the gravid uterus during the respective trimesters of pregnancy makes clinical examination difficult. The presence of leukocytosis and elevated serum alkaline phosphatase during pregnancy make interpretation of blood investigations difficult in making a diagnosis of acute cholecystitis.(5)

The treatment of acute cholecystitis in pregnancy can be divided into medical and surgical treatment. Medical treatment involves admitting the patients, keeping them fasted, starting intravenous antibiotics and analgesics and once the patient's condition had improved, cholecystectomy was often planned during the post-partum period. If a cholecystectomy needs to be done during the pregnancy it is usually performed during the second trimester, with surgery not performed during the first and third trimester due to the risk of threatened abortion and the size of the gravid uterus.(6–8)

Patients who undergo conservative treatment for acute cholecystitis in pregnancy often have a high readmission rate due to recurrent attacks and this can affect the wellbeing of the mother and the fetus. This has been the reason for performing the cholecystectomy earlier to manage this condition.(9)

If cholecystectomy is to be performed in the third trimester of pregnancy, the severity of acute cholecystitis must be graded and if the clinical condition deteriorates then laparoscopic cholecystectomy may have to be performed. If the symptoms improve then cholecystectomy may be performed in the post-partum period.(10)

The Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) has recommended laparoscopic cholecystectomy in the management of acute cholecystitis in pregnancy as it offers similar benefit to open surgery and it can be safely performed in any trimester of pregnancy if surgery is needed. Technical points like position of the patient, placement of the ports, the use of low-pressure insufflation and intraoperative monitoring of the carbon dioxide are important.(11)

As there is no current consensus on the management of acute cholecystitis in pregnancy, The role of conservative treatment is not well defined, the ideal time for performing a laparoscopic cholecystectomy is also not defined, either laparoscopic cholecystectomy can be performing in any of the trimesters of pregnancy and what is the consequence of delaying laparoscopic cholecystectomy to the post-partum period. We have conducted this review article looking for answers for all these factors in the management of acute cholecystitis in pregnancy. We conducted a literature review using PUBMED, the Cochrane database of systemic reviews, Google scholar and semantic scholar looking for randomized control trials, non-randomized trials, observational and cohort studies, clinical reviews, systemic reviews, case write ups, and meta-analysis from 1985 to 2023. The following keywords were used, "Acute cholecystitis in pregnancy", "Gallstone disease in pregnancy", "Conservative treatment of acute cholecystitis in pregnancy", "Laparoscopic cholecystectomy in acute cholecystitis in pregnancy" and "Cholecystectomy in pregnancy". All articles were in English, and all articles were assessed by manual cross referencing of the literature. Commentaries and editorials were excluded from this review. Only Pregnant patients with symptoms on non-obstetric abdominal pain were included in this study.

## **Discussion**

### **Conservative treatment of acute cholecystitis in pregnancy**

Pregnant patients who present with symptoms of acute cholecystitis are often treated conservatively. This involves the use of intravenous antibiotics and analgesics with the addition of intravenous fluids if the patient has persistent nausea and vomiting. There are limitations regarding the type of antibiotic prescribed due to pregnancy with the second and third generation cephalosporins being commonly used due to their safety.(12–14)

The use of analgesia is usually with opioids and paracetamol with non-steroidal anti-inflammatory drugs (NSAIDS) being contraindicated due to risk of complication to the fetus like oligohydramnios and premature closure of the patent ductus arteriosus. Conservative treatment is associated with a high recurrence rate of 40% to 77% and this can be detrimental to the mother and fetal well-being.(15–17)

A retrospective study by Othman et al on conservative treatment in pregnancy concluded that the number of visits of patients to the emergency department were higher in those who had undergone conservative treatment, and the readmission rates were also higher.(18)

A nationwide analysis of the morbidity associated with the conservative management of acute cholecystitis in pregnancy was conducted by Rios-Diaz et al. A total of 6390 patients were included in this study and they concluded that conservative treatment was associated with increased fetal-maternal complications, premature labor, increased caesarean section rates and poor fetal health.(19)

A review of the management of complicated gallstone disease in pregnancy by Date et al concluded that up to 27% of patients failed to respond to conservative treatment and the maternal and fetal mortality was similar between patients who had undergone conservative or surgical treatment. Another review by Lu et al comparing conservative treatment versus surgical treatment for acute cholecystitis in pregnancy also showed a recurrence rate of 38%, with patients in the second trimester exhibiting the highest recurrence rate..(20,21)

### **Surgical management of acute cholecystitis in pregnancy**

The surgical treatment of acute cholecystitis in pregnancy is cholecystectomy which can be performed as an open or laparoscopic procedure. The timing of surgery is important with the second trimester being the best time to perform it to obtain the best response. The risk of spontaneous abortion is higher during the first trimester and premature labor is high if surgery is done during the third trimester. Laparoscopic cholecystectomy is not a contraindication for patients who are pregnant, and the second trimester is the best time to perform it as trocar placement can be done without interference to size of the uterus.(22–30)

A meta-analysis was conducted by Athwal et al on the safety of surgery for symptomatic gallstone disease in pregnancy. A total of 470 patients were included in this study and the pre-term delivery rate was 6.8% and there was no significant difference with regards to the maternal and fetal mortality rates. This meta-analysis concluded that cholecystectomy can safely be performed in pregnancy.(31)

A systemic review and meta-analysis were conducted by Sedaghat et al comparing laparoscopic versus open cholecystectomy in the surgical management of acute cholecystitis in pregnancy. A total of 10,632 patients were included and laparoscopic cholecystectomy was associated with reduced maternal and fetal complications. The conversion rate was 3.8% and the length of the hospital stay was 3 days. This meta-analysis concluded that laparoscopic cholecystectomy was safe to perform but it could not tell when the best time was to perform surgery as most of the cases were operated on the first and second trimester.(32)

A systemic review on laparoscopic cholecystectomy during pregnancy was conducted by Nasioudis et al. A total of 590 patients were included in the study, The majority of the operations were performed during the second trimester and the post operative complication rates were 4% and the conversion to open cholecystectomy was 2.2%. The preterm delivery rate was 5.7% and this study showed that laparoscopic cholecystectomy can be safely performed in acute cholecystitis in pregnancy.(33)

Laparoscopic cholecystectomy in pregnancy should be performed with the patient in the supine position, port placement should be with the open Hassan technique, pneumoperitoneum should be maintained with a pressure of 8-12mmHg, and tocolytics should be administered and fetal heart rate monitoring be done.(34,35)

Laparoscopic cholecystectomy was found to be safe in pregnancy as it was associated with reduced short term fetal adverse effects, the operative time was shorter, length of hospital stay was reduced and there was a reduced requirement for blood transfusion. There was also a reduced risk of miscarriage. Other factors that can affect the outcome are independent factors like age above 35 years, the presence of jaundice and biliary peritonitis.(36–40)

Study	Year	Study type	N=numbers	Conversion rate (%)	Preterm delivery rate (%)	Complication rate (%)
Sedaghat et al	2016	Systemic review and meta-analysis	10,632	3.8%	8.7%	9.6%

Nasioudis et al	2016	Meta-analysis	590	2.2%	5.7%	4%
-----------------	------	---------------	-----	------	------	----

Table 1: The conversion rates, preterm delivery rates and complication rates of patients who underwent laparoscopic cholecystectomy in pregnancy.

### Percutaneous cholecystostomy in acute cholecystitis in pregnancy

Percutaneous cholecystostomy can be used to stabilize a pregnant patient who presents with acute cholecystitis. It can be used in combination with conservative treatment to treat pregnant patients with acute cholecystitis. Percutaneous cholecystostomy can be used in the first trimester to enable elective cholecystectomy to be performed in the second trimester. For pregnant patients who present with acute cholecystitis in the third trimester it can be used as a bridge to post-partum cholecystectomy. There are limited studies on the use of percutaneous cholecystostomy in acute cholecystitis in pregnancy, primarily case reports and case series. Large scale studies may need to be done to look at its efficacy as there are no consensus on its management.(41–45)

### Conclusion

The management of acute cholecystitis in pregnancy has seen a trend towards early laparoscopic cholecystectomy because of the high recurrence rate of conservative treatment. Laparoscopic cholecystectomy is not a contraindication for performing it in pregnant patients. The position of the gravid uterus makes placement of the port difficult, so surgeons who perform this operation will require some amount of experience in performing a laparoscopic cholecystectomy. Most general surgeons are still reluctant to perform surgery during pregnancy despite the evidence collaborating the safety of surgery in pregnancy.

Conservative treatment is still a popular form of therapy for acute cholecystitis in pregnancy as most hospitals may not have additional monitoring for the fetus and mother, hence this is a

deterrent for performing laparoscopic cholecystectomy in pregnancy. Most patients will ultimately get the cholecystectomy performed in the post-partum period.

Percutaneous cholecystostomy may be used as a bridging procedure to stabilize patients but it is not a popular form of treatment due to the problems of inserting the catheter and the contraindication of performing a tube cholangiography due to the radiation risk to the fetus.

Moving forward laparoscopic cholecystectomy in pregnancy is best to be performed in the second trimester as the risk to both the mother and fetus are the lowest. Pregnant patients with cholelithiasis should be counselled about the risk of complications of gallstone disease and the treatment options. More large-scale studies are needed to establish if surgery is the best form of treatment for acute cholecystitis in pregnancy.

Conflict of interest-There is no conflict of interest.

## References

1. Tseng JY, Yang MJ, Yang CC, Chao KC, Li HY. Acute Cholecystitis During Pregnancy: What is the Best Approach?. *Taiwan J Obstet Gynecol*. 2009;48(3):305-307. doi:10.1016/S1028-4559(09)60311-9
2. Salari N, Hasheminezhad R, Heidarisharaf P, et al. The global prevalence of gallstones in pregnancy: A systematic review and meta-analysis. *Eur J Obstet Gynecol Reprod Biol X*. 2023;19:100237. Published 2023 Sep 6. doi:10.1016/j.eurox.2023.100237
3. Ibitoye BO, Adisa AO, Makinde ON, Ijarotimi AO. Prevalence and complications of gallstone disease among pregnant women in a Nigerian hospital. *Int J Gynaecol Obstet*. 2014;125(1):41-43. doi:10.1016/j.ijgo.2013.09.022
4. Schwulst SJ, Son M. Management of Gallstone Disease During Pregnancy. *JAMA Surg*. 2020;155(12):1162-1163. doi:10.1001/jamasurg.2020.3683
5. Bouyou J, Gaujoux S, Marcellin L, et al. Abdominal emergencies during pregnancy. *J Visc Surg*. 2015;152(6 Suppl):S105-S115. doi:10.1016/j.jviscsurg.2015.09.017
6. Casey BM, Cox SM. Cholecystitis in Pregnancy. Vol. 4, Infectious Diseases in Obstetrics and Gynecology. iss, Inc; 1996:303-309. [https://doi.org/10.1002/\(SICI\)1098-0997\(1996\)4:5%3C303::AID-IDOG12%3E3.0.CO;2-S](https://doi.org/10.1002/(SICI)1098-0997(1996)4:5%3C303::AID-IDOG12%3E3.0.CO;2-S)
7. Grigoriu C, Lutic C, Scurtu I, Calinescu G, Gheorghe CM, Balan AM, et al. Management of biliary lithiasis in pregnancy – an updated overview. *Romanian Medical Journal*. 2021;68(4):433–6. DOI: 10.37897/RMJ.2021.4.2
8. Barut B, Gönültaş F, Gök AFK, Şahin TT. Management of Acute Cholecystitis During Pregnancy: A Single Center Experience. *Management of Acute Cholecystitis During Pregnancy: A Single Center Experience. Ulus Travma Acil Cerrahi Derg*. 2019;25(2):154-158. doi:10.5505/tjtes.2018.82357

9. Hedström J, Nilsson J, Andersson R, Andersson B. Changing management of gallstone-related disease in pregnancy - a retrospective cohort analysis. *Scand J Gastroenterol*. 2017;52(9):1016-1021. doi:10.1080/00365521.2017.1333627
10. Zhang W, Yi H, Cai M, Zhang J. Management strategies for acute cholecystitis in late pregnancy: a multicenter retrospective study. *BMC Surg*. 2023;23(1):340. Published 2023 Nov 10. doi:10.1186/s12893-023-02257-3
11. Pearl J, Price R, Richardson W, Fanelli R; Society of American Gastrointestinal Endoscopic Surgeons. Guidelines for diagnosis, treatment, and use of laparoscopy for surgical problems during pregnancy. *Surg Endosc*. 2011;25(11):3479-3492. doi:10.1007/s00464-011-1927-3
12. Augustin G, Majerovic M. Non-obstetrical acute abdomen during pregnancy. *Eur J ObstetGynecolReprod Biol*. 2007;131(1):4-12. doi:10.1016/j.ejogrb.2006.07.052
13. Ibiebele I, Schnitzler M, Nippita T, Ford JB. Outcomes of Gallstone Disease during Pregnancy: a Population-based Data Linkage Study. *Paediatr Perinat Epidemiol*. 2017;31(6):522-530. doi:10.1111/ppe.12406
14. Messidi A, Alsarraj G, Czuzoj-Shulman N, Mishkin DS, Abenhaim HA. Evaluation of management and surgical outcomes in pregnancies complicated by acute cholecystitis. *J Perinat Med*. 2018;46(9):998-1003. doi:10.1515/jpm-2017-0085
15. Gilo NB, Amini D, Landy HJ. Appendicitis and cholecystitis in pregnancy. *Clin Obstet Gynecol*. 2009;52(4):586-596. doi:10.1097/GRF.0b013e3181c11d10
16. Selzer DJ, Stefanidis D. Surgical Emergencies in the Pregnant Patient. *Adv Surg*. 2019;53:161-177. doi:10.1016/j.yasu.2019.04.008
17. Jorge AM, Keswani RN, Veerappan A, Soper NJ, Gawron AJ. Non-operative management of symptomatic cholelithiasis in pregnancy is associated with frequent hospitalizations. *J Gastrointest Surg*. 2015;19(4):598-603. doi:10.1007/s11605-015-2757-8
18. Othman MO, Stone E, Hashimi M, Parasher G. Conservative management of cholelithiasis and its complications in pregnancy is associated with recurrent symptoms and more emergency department visits. *GastrointestEndosc*. 2012;76(3):564-569. doi:10.1016/j.gie.2012.04.475
19. Rios-Diaz AJ, Oliver EA, Bevilacqua LA, et al. Is It Safe to Manage Acute Cholecystitis Nonoperatively During Pregnancy?: A Nationwide Analysis of Morbidity According to Management Strategy [published correction appears in Ann Surg. 2021 May 1;273(5):e184-e187]. *Ann Surg*. 2020;272(3):449-456. doi:10.1097/SLA.0000000000004210
20. Date RS, Kaushal M, Ramesh A. A review of the management of gallstone disease and its complications in pregnancy. *Am J Surg*. 2008;196(4):599-608. doi:10.1016/j.amjsurg.2008.01.015

21. Lu EJ, Curet MJ, El-Sayed YY, Kirkwood KS. Medical versus surgical management of biliary tract disease in pregnancy. *Am J Surg.* 2004;188(6):755-759. doi:10.1016/j.amjsurg.2004.09.002
22. Chiedozi LC, Al Hadi FN, Salem MM, Al Moaidi FA, Okpere EE. Management of symptomatic cholelithiasis in pregnancy. *Ann Saudi Med.* 2001;21(1-2):38-41. doi:10.5144/0256-4947.2001.38
23. Barber-Millet S, Bueno Lledó J, Granero Castro P, Gómez Gavara I, Ballester Pla N, García Domínguez R. Update on the management of non-obstetric acute abdomen in pregnant patients. Actualización en el manejo del abdomen agudo no obstétrico en la paciente gestante. *Cir Esp.* 2016;94(5):257-265. doi:10.1016/j.ciresp.2015.11.001
24. Hedström J, Nilsson J, Andersson B. Cholecystectomy and ERCP in pregnancy: a nationwide register-based study. *Int J Surg.* 2024;110(1):324-331. Published 2024 Jan 1. doi:10.1097/JS9.0000000000000812
25. Chiappetta Porras LT, Nápoli ED, Canullán CM, et al. Minimally invasive management of acute biliary tract disease during pregnancy. *HPB Surg.* 2009;2009:829020. doi:10.1155/2009/829020
26. Glasgow RE, Visser BC, Harris HW, Patti MG, Kilpatrick SJ, Mulvihill SJ. Changing management of gallstone disease during pregnancy. *Surg Endosc.* 1998;12(3):241-246. doi:10.1007/s004649900643
27. Costantino GN, Vincent GJ, Mukalian GG, Kliefoth WL Jr. Laparoscopic cholecystectomy in pregnancy. *J Laparoendosc Surg.* 1994;4(2):161-164. doi:10.1089/lps.1994.4.161
28. Cheng V, Matsushima K, Sandhu K, et al. Surgical trends in the management of acute cholecystitis during pregnancy. *Surg Endosc.* 2021;35(10):5752-5759. doi:10.1007/s00464-020-08054-w
29. Mazza GR, Youssefzadeh AC, Aberle LS, et al. Pregnant patients undergoing cholecystectomy: nationwide assessment of clinical characteristics and outcomes. *AJOG Glob Rep.* 2024;4(1):100310. Published 2024 Jan 11. doi:10.1016/j.xagr.2024.100310
30. Paramanathan A, Walsh SZ, Zhou J, Chan S. Laparoscopic cholecystectomy in pregnancy: An Australian retrospective cohort study. *Int J Surg.* 2015;18:220-223. doi:10.1016/j.ijssu.2015.05.005
31. Athwal R, Bhogal RH, Hodson J, Ramcharan S. Surgery for gallstone disease during pregnancy does not increase fetal or maternal mortality: a meta-analysis. *Hepatobiliary Surg Nutr.* 2016;5(1):53-57. doi:10.3978/j.issn.2304-3881.2015.11.02
32. Sedaghat N, Cao AM, Eslick GD, Cox MR. Laparoscopic versus open cholecystectomy in pregnancy: a systematic review and meta-analysis. *Surg Endosc.* 2017;31(2):673-679. doi:10.1007/s00464-016-5019-2
33. Nasioudis D, Tsilimigras D, Economopoulos KP. Laparoscopic cholecystectomy during pregnancy: A systematic review of 590 patients. *Int J Surg.* 2016;27:165-175. doi:10.1016/j.ijssu.2016.01.070



34. Halkic N, Tempia-Caliera AA, Ksontini R, Suter M, Delaloye JF, Vuilleumier H. Laparoscopic management of appendicitis and symptomatic cholelithiasis during pregnancy. *Langenbecks Arch Surg*. 2006;391(5):467-471. doi:10.1007/s00423-006-0069-x
35. Rollins MD, Chan KJ, Price RR. Laparoscopy for appendicitis and cholelithiasis during pregnancy: a new standard of care. *Surg Endosc*. 2004;18(2):237-241. doi:10.1007/s00464-003-8811-8
36. Shigemi D, Aso S, Matsui H, Fushimi K, Yasunaga H. Safety of Laparoscopic Surgery for Benign Diseases during Pregnancy: A Nationwide Retrospective Cohort Study. *J Minim Invasive Gynecol*. 2019;26(3):501-506. doi:10.1016/j.jmig.2018.06.008
37. Vujic J, Marsoner K, Lipp-Pump AH, Klaritsch P, Mischinger HJ, Kornprat P. Non-obstetric surgery during pregnancy - an eleven-year retrospective analysis. *BMC Pregnancy Childbirth*. 2019;19(1):382. Published 2019 Oct 25. doi:10.1186/s12884-019-2554-6
38. Palanivelu C, Rangarajan M, Senthilkumaran S, Parthasarathi R. Safety and efficacy of laparoscopic surgery in pregnancy: experience of a single institution. *J Laparoendosc Adv Surg Tech A*. 2007;17(2):186-190. doi:10.1089/lap.2006.0037
39. Itaimi A, Abbassi I, Baraket O, Kotti A, Triki W, Bouchoucha S. Safety of Laparoscopic Cholecystectomy for Cholecystitis during Pregnancy. *Gynecol Minim Invasive Ther*. 2023;12(3):166-169. Published 2023 Aug 10. doi:10.4103/gmit.gmit\_57\_22
40. Nan X, Chan E, Wong KSC, et al. Laparoscopic Cholecystectomy in Pregnancy: A Seven-Year Retrospective Study From an Australian Tertiary Center. *Cureus*. 2023;15(12):e50034. Published 2023 Dec 6. doi:10.7759/cureus.50034
41. Little MW, Briggs JH, Tapping CR, et al. Percutaneous cholecystostomy: the radiologist's role in treating acute cholecystitis. *Clin Radiol*. 2013;68(7):654-660. doi:10.1016/j.crad.2013.01.017
42. Allmendinger N, Hallisey MJ, Ohki SK, Straub JJ. Percutaneous cholecystostomy treatment of acute cholecystitis in pregnancy. *Obstet Gynecol*. 1995;86(4 Pt 2):653-654. doi:10.1016/0029-7844(95)00087-8
43. Hojberg Y, Patel K, Shebrain S. Utilizing Percutaneous Cholecystostomy Tube as a Temporary Minimally Invasive Approach for Acute Cholecystitis during Third Trimester of a High-Risk Pregnancy. *Case Rep Gastroenterol*. 2022;16(1):49-54. Published 2022 Feb 14. doi:10.1159/000522060
44. Hanks LR, Towner MN, Hollenbach SJ, Drennan KJ, Olson-Chen C. Nonsurgical Management of Acute Cholecystitis in Pregnancy: A Cautionary Tale. *ACS Case Reviews in Surgery*. 2020;3(2):53–56.
45. Caliskan K. The use of percutaneous cholecystostomy in the treatment of acute cholecystitis during pregnancy. *Clin Exp Obstet Gynecol*. 2017;44(1):11-13.