

Original Research Article

**EFFECT OF DEHAULMING AND STORAGE CONDITIONS ON POSTHARVEST
PARAMETERS OF POTATO (*Solanum tuberosum*)**

Abstract

The research was conducted in Rampur, Udayapur. Two factorial Complete Randomized Design with two level of treatment combination was applied throughout the experiment. The variety chosen was lalgulab collected from the farmer's field in the month of February. During the experiment, dehaulming and non-dehaulming practice was performed and the potatoes were stored in three different storage conditions (Cold storage, Bamboo basket and Gunny bags). It was noted that parameters like rotting, sprouting, total weight loss and greening were significantly influenced by different storage conditions along with the dehaulming practices. The maximum rotting (8.30%), minimum rotting (0%), maximum sprouting (87.38%), minimum sprouting (0%), maximum final weight loss (14.20%), minimum final weight loss (3.22%), superiority in greening (18.15%) and minimum greening (0%). The interaction effect of dehaulming and non-dehaulming with different storage conditions had significant effect at 20days in rotting and at 40days in sprouting. The interaction effect of final weight loss and greening at 20, 40, 60 and 80days was not found significant with the different storage conditions. Similarly, dehaulmed potatoes is best for minimum rotting percentage, minimum sprouting percentage, minimum final weight loss percentage and minimum greening percentage. Among all the storage conditions, cold storage was found appropriate for storage with minimum storage loss followed by dehaulming practices.

Keywords:??

1. Introduction

Potatoes (*Solanum tuberosum*) occupies the fourth position as the important crops after wheat, maize and rice in the world (Bajracharya & Sapkota, 2017). Potatoes are ranked fifth on area covered, second on total production and first on productivity in Nepal (MoALD, 2021). They are a staple food rich in vitamins, minerals and fibers and have a low glycemic index making them a good option for people with diabetes. According to the recent study, the area, production and yield of potato were found 1876ha, 28900mt and 15.41mt/ha respectively in Udayapur (MoALD, 2021).

Dehaulming is the practice of removing the aerial parts of a plant 10-15 days before harvesting ("Dehaulming in potato", 2018). According to Litaladio et al. (2009), dehaulming can be carried out after the yellowing of aerial parts, as this signifies the potato's maturity. Dehaulming extends the shelf life of potatoes by hardening the tubers. The potato tubers are left in the field for 2-3 weeks after dehaulming so that the skin of potato hardens and reduces the damage during postharvest handling (Potato Vocabularies Explained, 9 Mar 2020). Dry matter weight percentage of potato flesh was measured at 0, 30, 60, 90, and 120 days after storage, and found to be 20.55%, 21.28%, 22.31%, 24.60%, and 27.32% respectively in potatoes harvested 12 days

Commented [N1]: Too much unnecessary para and sentence were included in this section

Commented [N2]: If possible then give update information

Commented [N3]: If possible then give update information

after dehaulming and 12.85%, 13.52%, 15.52%, 16.55%, and 18.81% in potatoes harvested 0 days after dehaulming (Nipa et al, 2013).

Postharvest loss refers to any change in the produce after harvesting that hinders the normal consumptions. Postharvest loss of potato was estimated to be 25% in Nepal, 24% in India and 20% in Bangladesh (Karki, 2002; Prasad et al., 1989; Satter et al., 2002). From the study, the losses of potato during postharvest storage were estimated between 15-20% in Nepal (Bhattarai, 2018). The loss of potato tubers was found to be 12.08% per hectare in average in various supply chain stages (Kuyu et al., 2019).

Harvested potatoes are kept into storage before being distributed to the market. The main goals of storage are to preserve quality, prevent infestation of disease and pest, regulate supply and provide security to the product. Maintaining the unique microclimate characteristics at each stage of potato storage, such as temperature and relative humidity, is necessary. (Potato Storage Technology, 2020). Though traditional methods of storage suffer considerable losses, the number of farmers relying on the traditional methods of potato storage is increasing. The main reason for increasing the use of local methods of potato storage are cheap and easy handling as well as the potatoes taste better than that of stored in cold storage (Dahiya et al., 1997).

1. Botanical description of potato

Potato is a starchy tuber of plant scientifically named as *Solanum tuberosum*. It is a root crop native to America belonging to family Solanaceae (Britannica, 2022 November 11). Potato is mostly cross pollinated by insects but sometimes self-pollination as well occurs (Amador, 2009). Planting tuber fragments allows for vegetative propagation of potatoes, which can also be cultivated from botanical seeds. When cultivated under the right conditions, eyes on the surface of potato tubers produce sprouts. When the tubers are utilized for propagation, the ideal sprouting is a desired characteristic. (Singh and Kaur, 2009). The average composition of potato consists of 77.5% moisture, 19.4% carbohydrate, 2% protein, 0.6% fiber and 1% ash (Gould, 1999).

According to Theisen and Compilan (2010), potatoes are a winter crop in the terai and low hills, a spring and autumn crop in the mid-hills, and a summer crop in the high hills and mountains. In Nepal, potatoes are often sown in late September or early October. Potato harvesting can begin as soon as two months after sowing (Dhital, 2000). Potatoes are the fourth most important food crop in the world. Two thirds of them are consumed by humans, while the other third is used as animal feed, potato starch, and in the manufacturing of pharmaceuticals, textiles, adhesives, paper products, and other products (Ehler, 2015).

2. Potato variety

There are many varieties grown in eastern part of Nepal. The mainly grown commercial varieties are Lal gulab, KurfiSinduri (C140 or C40), Chisapani, llam blue, Phulwa, Letang white and so on. Lal gulab is the terai grown variety which is round red in color with thick and scaly skin. It has uniform size and highly valued for consumption with good keeping quality. It does not deteriorate as rapidly as other varieties and have high market value (Rhoades, 1985).

3. Postharvest loss of potato

Commented [N4]: Give update reference

The losses that occur during the period between maturity of crop and the time of its final consumption are called postharvest losses (Grieig and Reeves, 1985). Potato postharvest losses are mostly caused by water loss, mechanical damage, physiological damage, illness, and pest infestation (www.fao.org/docrep/008/a018e/a01850c.htm). According to Sudheer and Indira (2007), inadequate harvesting, transportation, marketing, and legal frameworks provide the right environment for secondary causes of loss. To avoid losing weight, rotting, shrinking, sweetening, greening, and sprouting, potatoes should be stored in an appropriate environment (Gotts Chalk and Christenbury, 1998).

Postharvest losses can be the result of a qualitative or quantitative losses (Aulakh et al., 2013). A product's acceptability, edibility, and nutrient composition are all impacted by quality losses. Losses of a product's quantity are included in quantity losses (Kaiya, 2014).

4. Impact of dehauling on postharvest and qualitative aspects

Dehauling is the process of removing the potato plant's vegetative portion from its above-ground root tubers (Potato Vocabularies Explained, 9 Mar 2020). Dehauling extends the shelf life of potatoes by hardening the tubers. After harvesting, the potato tubers are left in the field for two to three weeks to allow the skin to harden and minimize harm from handling them after harvest (Potato Vocabularies Explained, 9 Mar 2020).

The dry matter weight percentage of the potato flesh was determined to be 12.85%, 13.52%, 15.52%, 16.55%, and 18.81% in potatoes harvested 0 days after dehauling at 0, 30, 60, 90 and 120 days, and 20.55%, 21.28%, 22.31%, 24.60%, and 27.32% in potatoes harvested 12 days after dehauling respectively at 0- and 120-days following storage (Nipa et al, 2013). When dehauling delayed from 70 to 90 DAP, there was a notable decrease in the reducing sugar level and an improvement in tuber quality (Marwaha et al, 2012). After peeling or cutting, phenolic acid and flavonoids induce enzymatic discoloration (Marwaha et al, 2010).

5. Types of storage systems

Following are the major storage methods practiced in Nepal:

A. Conventional methods for storing potatoes

- Storage in bhakaris
- Pit storage/Clamps dug in the ground
- Heap storage
- Storage with the use of chemicals
- Bamboo baskets lined with soil and straw
- Sack storage (Gautam, 2014)
- Room storage

B. Modern methods of storage

- Cold storage (Theisen and Campilan, 2010)
- Stored in diffused sunlight (Anon, 2004)

The people of Udayapur are facing the problem of postharvest loss and potatoes greening during storage. Hence, this experiment was performed under farmers condition to study the effect of harvesting parameters and postharvest losses of potato under different storage conditions.

6. Room storage

Room storage is the traditional storage practices followed over all the ecological regions of Nepal. In such storage systems potatoes are spread out in a dark room under the beds or corners to dry for about 7-10 days after harvesting (Rhoades, 1985). This storage technique is mainly practiced to devoid sunlight from the potatoes during storage period as when potatoes are exposed to strong light, they got materially injured for table purposes. The injury thus occurred is due to the greening of the tubers and development of alkaloid called solanine in the outer layer and around the eyes causing issues to human health. The problems reported following this storage is, this is space consuming as the potatoes are spread all over the floor which decreases the seed stock vigor and uniformity due to excessive sprouting (Khatana et al., 1997).

7. Storage in bamboo baskets

In this practice, the bamboo baskets locally known as doko (average sized) lined up with sand from outside and mud from inside is left to dry well in the sunlight. Potatoes are then well mixed with dry mustard straw and kept in the bamboo baskets in the dark room. Dry mustard straw is used to create dry and cool conditions within storage area (Mutandwa and Gadzirayi, 2007). It also prevents the further moisture loss from potatoes (Rhoades, 1985). The holes in the basket provides the proper ventilation and prevents overheating.

8. Low temperature storage

Cooling the product after harvesting controls the rate of quality loss by decreasing the rate of respiration. The higher the temperature, faster is the deterioration and lower the shelf life of the product so the temperature should be lowered to increase the storage life of product (Sakare, 2014). In cold stores, seed potatoes should be kept between 2-4°C and 90-95% RH to prevent sprouting and minimize weight loss. Once removed from storage, potatoes kept in a cold environment do not keep well, hence it is recommended to store potatoes at 10–12°C for processing and table use. Potatoes stored at this temperature do not become sweet minimizing the accumulation of reducing sugars (Singh, 2016).

9. Effect of different storage conditions on postharvest loss

Low temperature storage, cold storage, root cellar storage, traditional clamp storage, pit storage, bamboo baskets storage, gunny bags and normal room storage, hanging baskets are the major storage techniques practiced in Nepal (Rhoades, 1985). Postharvest loss decreases the quantity and quality of crop due to improper storage, handling, transportation and market infrastructure. The total postharvest loss was found 4.383%, 8.539% and 13.043% in cold storage, in-basket storage and in-house storage conditions respectively. With cold storage, reducing sugar reached a maximum value of 1.04% as storage time climbed. In potatoes, post-harvest losses ranged from 20 to 30 percent, and occasionally as high as 50 percent (Karki, 2002; Rasheed and Shareef, 2002; Wijeratnam, 2002). According to Khanal and Bhattarai (2020), among the three storage conditions monitored during the experiment, cold storage (2–5°C/85–90%RH) produced the least amount of storage loss, but room storage (25–33°C/75–85%RH) produced the most.

2. Materials and Methods

2.1 Harvesting of potato tubers

Commented [N5]: Reference

Potato tubers were harvested after the yellowing of aerial parts and left for curing before storage. The potatoes were then sorted and kept in storage after 10 days of harvesting. Rotten and deteriorated tubers were discarded and fresh tubers each sample weighting about 5kg were sorted out for the experiment purpose.

2.2 Preparation of storage structures

The storage in normal room temperature was done by placing the potato tubers in gunny bags. Bamboo basket locally known as doko (average sized) was lined with mud and sand and mustard straw were procured for bamboo basket storage and arranged. The primary use of straw is to regulate the humidity and accessibility. For the easy availability of laboratory analysis and convenience during the study, environment of low temperature storage was recreated in cold storage (Model: A7060 – 353A, Capacity: 7ton Power supply: 230V.60Hz). Maintenance of these conditions were confirmed with the use of hygrometer throughout the storage period (Model: HTC-1).

2.3 Equipment used

General equipment like digital weighing balance, hygrometer, cold storage, spade, sickle, etc were used during the entire study period. Digital weighing balance and hygrometer were brought from the laboratory of Mahendra Ratna Multiple Campus, Ilam.

2.4 Experimental procedure

Each storage unit was loaded with 5kg of fresh, healthy potato tubers following variety selection and careful sorting of similar-sized tubers. Tubers were organized in bamboo baskets with chopped mustard straw between layers after the baskets were lined with mud and sand and allowed to dry for a day. For room storage, the tubers were stored in gunny bags; for cold storage, they were packed in sacks. Four replications of such specimens were conducted in order to acquire accurate results. For each sample, the experimental data was recorded every 20 days up to an 80-day period. Analysis was done for final weight loss percentage, greening, sprouting, and rotting during the storage period. At the conclusion of the 80-day storage period, the weight of each sample under storage conditions was measured in order to estimate the overall loss.

2.5 Treatments and Experiment Design

The experiment was started from 14 Magh, 2079 to 18 Jestha, 2080 in farmer's field condition at Belaka municipality of Udayapur district as it is one of the significant potato growing area. Belaka municipality was declared multiple crop zone including potato as a core crop on 2021/2022. The site is located geographically at 26°42'09" N- 86°55'38" E latitude and 26°55'38" N – 87°10'06" E longitude with the elevation of around 150masl.

The variety selected for research was lalgulab. With four replications and six treatments, the design was set up in a two factorial Completely Randomized Design (CRD). Each sample was considered as one treatment. Factor A includes the storage conditions (Cold storage, Bamboo basket and Gunny bags) whereas Factor B includes the dehaulming and non-dehaulming practices with treatment combination A×B.

Commented [N6]: Please use gregorian calendar year

Table 1: Detail of treatments

S.N.	Treatment
T ₁	Dehaulming + bamboo basket lined with mud and straw
T ₂	Dehaulming +gunny bag in normal room temperature
T ₃	Dehaulming + cold storage
T ₄	Without dehaulming + bamboo basket lined with mud and straw
T ₅	Without dehaulming +gunny bag in normal room temperature
T ₆	Without dehaulming + cold storage

Commented [N7]: Please give temperature of cold storage

2.6 Data Collection

Potato variety lalulab was taken as a sample. The sample were collected from the dehaulmed and non-dehaulmed field. After dehaulming, the potatoes were left in the field for 15 days and then harvesting was done. All the samples were harvested in the same day and left for curing for 10days. Then, all the samples were placed in different storage conditions and the changes was noted in the interval of 20day.Total four data were collected.

2.6.1 Parameters Observed

(1) Weight loss Percentage (%)

Total weight loss = Initial weight of tubers – Final weight of tubers

Final weight loss (%) = Total weight loss ÷ Initial weight of tubers × 100

(2) Sprouting Percentage (%)

Total loss (%) = Weight of sprouts ÷ Total initial weight of tubers × 100

(3) Damage due to rotting Percentage (%)

Total rotting (%) = Weight of deteriorated tubers ÷ Total initial weight of tubers × 100

(4) Greening percentage (%)

Greening (%) = Weight of green tubers ÷ Initial weight of tubers × 100

2.7 Data analysis

First, the gathered data was cleaned up, tallied, and entered into MS-Excel LTSC 2021. R-stat Version 4.2.1 statistical software was utilized to analyze the entered data and derive significant conclusions. The Honestly Significant Difference Test (HSD) was used to compare the mean of the examined parameters at the 5% significance level. The significance difference of each parameter was tested using an ANOVA.

3. Results and Discussion

3.1 Rotting Percentage

Rotting percentage generally varies as per the difference in the storage conditions where potatoes are stored. The results obtained during the study showed that in all cases i.e., at 20, 40, 60 and 80 days, the rotting percent was found highly significant in the storage conditions. Non-dehaulmed potatoes showed significant difference in rotting at 20days interval followed by non-significant difference at 40, 60 and 80days interval. The maximum rotting percentage was observed 8.30% in gunny bags and the minimum rotting percentage was recorded 0% in cold storage. The

interaction between two factors was found significantly different in 20days but no variation was observed in 40, 60 and 80days.

Loss due to the changes in moisture and respiration rate was observed in potato stored inside house in gunny bags as the relative humidity and temperature are not controlled which causes a potatoes to rot and directly affects the weight loss of potato greatly reducing the market value of tubers (Keasar et al., 2005). The obtained result was similar to my findings.

Because mustard straw is used instead of dry straw, which inhibits transpired vapour from condensing on the surface of tubers, the percentage of degradation in the bamboo basket was lower than in the gunny bags. Thus, prevents an overly wet environment in the storage area. By doing this, fungal diseases—which typically thrive in warm, humid environments—are prevented from developing (Mutandwa & Gadzirayi, 2007). According to Knutsson (2012), straw absorbs condensed water produced by the tuber's respiration, which may increase the presence of microorganisms and validate the results of my study.

According to Ezekiel and Dahiya (2002), decaying losses in room storage was 4.7%, but losses in heap storage was 3.8%. However, according to Singh et al. (2014), rotting causes roughly 10–12% of the loss in a typical storage method. The observed outcome shows 8.30% of the maximum loss.

Table 2: Effect of dehauling and storage conditions on rotting percentage of potatoes

Rotting				
Treatment	20days	40days	60days	80days
Factor A				
Cold storage	0±0 ^c	2.10±0.18 ^b	3.19±0.19 ^b	4.75±0.26 ^b
Bamboo basket	1.15± 0.07 ^a	2.78±0.20 ^b	3.69±0.21 ^b	4.87±0.20 ^b
Gunny bags	0.66±0.03 ^b	3.60±0.21 ^a	5.94±0.24 ^a	8.30±0.24 ^a
LSD	0.11***	0.76***	0.80***	0.88***
Factor B				
Non-dehauling	0.64±0.16 ^a	2.74±0.27	4.28±0.45	3.66±0.35
Dehauling	0.56±0.12 ^b	2.91±0.20	4.26±0.34	4.60±0.34
LSD	0.07*	0.51NS	0.54NS	0.59NS
Interaction (A×B)				
LSD	0.20**	1.34NS	1.42NS	1.56NS
CV%	15.34	21.21	14.83	11.61
Grand mean	0.60	2.83	4.27	5.97

Table 3: Effect of dehauling and storage conditions on rotting percentage of potatoes (Interaction)

Rotting at 20days

Treatment	Non-dehaulming	Dehaulming
Cold storage	0 ^d	0 ^d
Bamboo basket	1.3 ^a	1 ^b
Gunny bags	0.64 ^c	0.69 ^c
CV (%)	15.34	
LSD (_{0.05})	0.20**	
Grand mean	0.60	

Note: The common letter (s) within the column indicate the non-significant difference based on HSD Test at 5% level of significance. Significance codes: 0 ‘***’ 0.001 ‘**’ 0.01 ‘*’ 0.05 ‘.’ 0.1 ‘.’ 1. The \pm values represent the standard error of the mean.

3.2 Sprouting Percentage

At 20days of storage period, no any signs of sprout growth were seen in both dehaulmed and non-dehaulmed samples followed by storage conditions. The maximum sprouting percentage was observed (87.38%) in bamboo basket and the minimum sprouting percentage was recorded (0%) in cold storage. Dehaulmed and non-dehaulmed potatoes were found significantly different in 40 and 80days. The interaction between two factors was found significantly different in 40days.

After harvest, sprouting takes some time. The dormant phase is a span of time that exists (Dris and Jain, 2004). Because of which, there was no any significant difference over the span of 20 days. Kaul and Mehta (1999) reported that after 15 weeks of storage, the weight loss resulting from sprouting is 0.6% for Kufri Phulwa and 0.67% for Kufri Sinduri. Because variety affects both dormancy and the amount of sprouting, the Lal gulab variety's acquired result differs significantly from the other varieties in this study (Mani & Hannachi, 2015). Given that the sprouting phenomena is primarily temperature-dependent (Khanal and Bhattarai, 2020), the temperature increase throughout the storage period (31°C–44°C) compared to the ambient temperature (29°C) is the reason for the noticeably larger loss in bamboo basket storage.

Table 4: Effect of dehaulming and storage conditions on sprouting percentage of potatoes

Treatment	Sprouting			
	20days	40days	60days	80days
	Factor A			
Cold storage	0 \pm 0	1.23 \pm 0.14	9.85 \pm 0.29 ^c	43.22 \pm 0.59 ^c
Bamboo basket	0 \pm 0	1.01 \pm 0.07	24.95 \pm 0.68 ^a	87.38 \pm 1.02 ^a
Gunny bags	0 \pm 0	1.18 \pm 0.08	15.27 \pm 0.57 ^b	64.66 \pm 1.48 ^b
LSD	0NS	0.24NS	1.80***	3.37***

Factor B				
Non-dehauling	0±0	1±0.06 ^b	17.07±2.11	66.65±5.59 ^a
Dehauling	0±0	1.28±0.08 ^a	16.31±1.72	63.53±5.37 ^b
LSD	0NS	0.16**	1.21NS	2.26**
Interaction (A×B)				
LSD	0NS	0.43**	3.17NS	5.94NS
CV%	0	16.94	8.47	4.06
Grand mean	0	1.14	16.69	65.09

Table 5: Effect of dehauling and storage conditions on rotting percentage of potatoes (Interaction)

Sprouting at 40 days		
Treatment	Non-dehauling	Dehauling
Cold storage	0.88 ^b	1.59 ^a
Bamboo basket	0.89 ^b	1.14 ^b
Gunny bags	1.22 ^{ab}	1.13 ^b
CV (%)	16.94	
LSD (_{0.05})	0.43**	
Grand mean	1.14	

Note: The common letter (s) within the column indicate the non-significant difference based on Honestly Significance Difference test at 5% level of significance. Significance codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1. The ± values represent the standard error of the mean.

3.3 Final Weight loss Percentage

Maximum final weight loss percentage was recorded in bamboo basket (14.20±0.64). The minimum final weight loss percentage was recorded in gunny bags (3.22±0.62). Non-dehauling samples show higher percentage of final weight loss followed by dehauling samples. No any signs of significant difference were observed in the interaction between two factors.

When storage temperature is lowered, tuber weight loss decreases, and cold storage is the best way to maintain tuber weight (Aharoni et al., 2007). It is because as compared to other storage conditions, cold storage has a lower storage temperature (10–12°C). As sprouted tubers lose a greater amount of weight than unsprouted ones (Azad et al., 2017), the higher sprouting rate in the bamboo basket accounts for the larger weight loss that is ultimately experienced in the bamboo basket.

According to Mutandwa and Gadzirayi (2007) a moderately warm and dry atmosphere can be created inside the basket by mixing straw, mud and the inherent qualities of bamboo. Warmth

accelerates the rate of moisture evaporation from the potato resulting in greater weight loss over time. But in cold storage, the temperature was maintained cool and humid due to which the rate of evaporation slow down and enhance the shelf life of potato.

Table 6: Effect of dehauling with different storage conditions on final weight loss percentage of potatoes

Treatment	Final weight loss percentage			
	20days	40days	60days	80days
Factor A				
Cold storage	3.27±0.78 ^b	6.11±0.74 ^{ab}	8.29±0.73 ^b	11.08±0.77 ^b
Bamboo basket	5.32±0.61 ^a	7.53±0.64 ^a	10.86±0.65 ^a	14.20±0.64 ^a
Gunny bags	3.22±0.62 ^b	5.36±0.81 ^b	10.07±0.71 ^a	12.55±0.77 ^{ab}
LSD	1.40**	1.78*	1.62**	1.74***
Factor B				
Non-dehauling	5.40±0.52 ^a	7.80±0.59 ^a	11.16±0.51 ^a	14.14±0.56 ^a
Dehauling	2.48±0.30 ^b	4.87±0.29 ^b	8.32±0.43 ^b	11.08±0.49 ^b
LSD	0.94***	1.20***	1.09***	1.17***
Interaction (A×B)				
LSD	2.47NS	3.15NS	2.86NS	3.06NS
CV%	27.90	22.11	13.07	10.82
Grand mean	3.94	6.33	9.74	12.61

Note: The common letter (s) within the column indicate the non-significant difference based on HSD Test at 5% level of significance. Significance codes: 0 ‘***’ 0.001 ‘**’ 0.01 ‘*’ 0.05 ‘.’ 0.1 ‘.’ 1. The ± values represent the standard error of the mean.

3.4 Greening Percentage

Very highly significant difference was observed in data of 20, 40, 60 and 80days. Maximum greening percentage was recorded in gunny bags (18.15±0.59). ~~The minimum greening percentage was recorded in cold storage (0±0).~~ No greening potato was found in cold storage (0±0). Non-dehaulmed samples shows more greening than dehaulmed samples. No any signs of significant difference were observed in the interaction between two factors.

Buck, R.W., Akeley, and R.V. (1967) came to the conclusion that maturity, storage temperature, storage duration, each influenced the degree of greenness. All of the studied varieties responded to greening in the same way, though to varying degrees.

Tanios et al. (2018) stored potatoes in light or darkness for up to 8 days at 5, 10 and 25°C. The level of chlorophyll accumulation that resulted in the greatest amount of greening was at 20°C. Since, potato stored in gunny bag in a room temperature ranges the temperature more than 25°C during my storage period so it may have caused the maximum greening in gunny bags. Chlorophyll synthesis in potato is slowed down by cold temperatures in the range of 4 to 10°C. The low temperature in cold storage inhibits the formation of chlorophyll, further limiting the greening process with the faster rate of greening at room temperature (Folsom 1947; Kozukue and Mizuno 1990; Edwards and Cobb 1997). Due to the controlled condition inside cold storage, the greening was observed in minimum rate.

Table 7: Effect of dehauling with different storage conditions on greening percentage of potatoes

Treatment	Greening			
	20days	40days	60days	80days
Factor A				
Cold storage	0±0 ^c	3.97±0.21 ^c	3.97±0.21 ^c	4.39±0.22 ^c
Bamboo basket	1.31±0.07 ^b	6.04±0.36 ^b	12.76±0.60 ^b	13.92±0.61 ^b
Gunny bags	3.82±0.14 ^a	8.18±0.39 ^a	16±0.47 ^a	18.15±0.59 ^a
LSD	0.35***	1.23***	1.48***	1.69***
Factor B				
Non-dehauling	1.74±0.50	6.15±0.57	11.41±1.67 ^a	12.67±1.88
Dehauling	1.68±0.46	5.98±0.58	10.41±1.46 ^b	11.63±1.65
LSD	0.23NS	0.83NS	0.99*	1.13NS
Interaction (A×B)				
LSD	0.61NS	2.17NS	2.60NS	2.98NS
CV%	16.04	15.96	10.62	10.93
Grand mean	1.71	6.06	10.91	12.15

Note: The common letter (s) within the column indicate the non-significant difference based on HSD Test at 5% level of significance. Significance codes: 0 ‘***’ 0.001 ‘**’ 0.01 ‘*’ 0.05 ‘.’ 0.1 ‘.’ 1. The ± values represent the standard error of the mean.

4. Conclusion

Recorded parameters such as rotting, sprouting, final weight loss and greening at 20, 40, 60 and 80days were significantly influenced due to the different storage conditions. The superiority of rotting was recorded in gunny bags whereas the minimum rotting was observed in cold storage. Similarly, maximum sprouting was observed in bamboo baskets and minimum sprouting was observed in cold storage. The maximum final weight loss was observed in bamboo basket

followed by minimum final weight loss in cold storage. The superiority in greening was observed in gunny bags whereas minimum greening was observed in cold storage. Non-dehaulmed potatoes showed maximum loss in observed parameters whereas dehaulmed potatoes showed minimum loss.

The interaction effect of dehaulming and non-dehaulming with different storage conditions had significant effect in rotting at 20days and in sprouting at 40days. The interaction effect of final weight loss and greening at 20, 40, 60 and 80days was not found significant with the different storage conditions.

On the basis of result obtained from the study, dehaulmed potato stored in different storage conditions (Cold storage, Bamboo basket and Gunny bags) is best for minimum rotting, sprouting, final weight loss and greening percentage. Among all the storage conditions used, best result was obtained in cold storage with minimum storage loss. The farmers are suggested to use dehaulmed potatoes with the storage under cold store. As the potato was stored in 10-12 degree Celsius, there was no sweetness in the cold stored potatoes so preferred by the consumers for table purpose. Therefore, it is suggested to conduct similar research in different location using other varieties along with dehaulming practices.

References

- Aharoni N, Rodov V, Fallik E, Afek U, Chalupowicz D, Aharon Z, Maurer D, Orenstein J. (2007). Modified atmosphere packaging for vegetable crops using high-water-vapor-permeable films. In CL Wilson (Ed.), *Intelligent and Active Packaging for Fruits and Vegetables* (1st ed., pp. 73–112). Boca Raton: CRC Press Taylor & Francis Group. <https://doi.org/10.1201/9781420008678.ch5>.
- Amador, V. (2009). *International Journal of developmental biology*. [page??](#)
- Anon (2004). Annual Report of the Coconut Research Institute (CRI) of Sri Lanka. Lunuwila: CRI.
- A. Kaur, N. Singh, R. Ezekiel et al. (2009). Properties of starches separated from potatoes stored under different conditions.
- Aulakh, J., Regmi, A., Fulton, J. and Alexander, C. (2013). Estimating post-harvest food losses: developing a consistent global estimation framework.
- Azad, A.K., Kabir, H., Eaton, T.E.-J., Soren, E.B., 2017. Storage Potentialities of Some Exotic Potato Varieties at Farmers' Condition in Bangladesh. *Agricultural Sciences* 8, 183.
- Bhattarai, D.R. (2018). Postharvest horticulture in Nepal. *Horticulture International Journal*, 2, Pp. 458–460.
- Bajracharya, M. & Sapkota, M. (2017). Profitability and productivity of potato (*Solanum tuberosum*) in Baglung district, Nepal. *Agriculture and Food Security*, 6(47), 1-8.
- Britannica, T. Editors of Encyclopaedia (2022, November 11). *potato*. *Encyclopedia Britannica*. <https://www.britannica.com/plant/potato>

Commented [N8]: Follow journal rules of referencing

Commented [N9]: rewrite it

Commented [N10]: rewrite it

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Dahiya, P. S., Khatana, V. S., Ilangantileke, S. G. and Dabas, J. P. S. (1997). "Potato storage patterns and practice in Meerut District, western Uttar Pradesh, India". *International potato center*. India.

Formatted: Font: Italic

"Dehaulming in potato". (2018, December 08). Retrieved from <https://www.agriedu4u.com/2018/11/14/dehaulming-in-potato/>

Dhital, B. K. (2000). Diffused Light Storage of Seed Potatoes in the Hills of Nepal in Relation to Seed Tuber Physiology, Production Environment, and SocioEconomics. (Ph.D. Thesis). University of Newcastle upon Tyne

Dris, R. and Jain, S. M. (2004). "Post Harvest Treatment and Technology". Vol. 4. Kluwer Academic Publishers. New York.

Edwards, E.J., and A.H. Cobb. 1997a. Effect of temperature on glycoalkaloid and chlorophyll accumulation in potatoes (*Solanum tuberosum* L. cv. King Edward) stored at low photon flux density, including preliminary modeling using an artificial neural network. *Journal of Agricultural and Food Chemistry* 45: 1032–1038

Formatted: Font: Italic

Ehler, J. T. (2015). Potato production and consumption. Retrieved from <http://www.foodreference.com/html/f-potato-consumption.html>. (Accessed 8 November 2015)

Ezekiel, R., P.S. Dahiya and G.S. Shekhawat. 2002. Traditional methods of potato storage in the Malwa region of Madhya Pradesh. Technical Bulletin No. 57. Central Potato Research Institute, Shimla, India 39 p (4) (PDF) *Changes in temperature and relative humidity in heap and pit during storage of potatoes*. Available from: https://www.researchgate.net/publication/259151776_Changes_in_temperature_and_relative_humidity_in_heap_and_pit_during_storage_of_potatoes [accessed Jan 08 2024].

Folsom, D. 1947. Permanence of greening of potato tubers. *American Journal of Potato Research* 24: 336–340.

Formatted: Font: Italic

Gautam, I. P. (2014). Methods of potato storage (Video clip, 14:48 min). National potato development project. khumaltar, Lalitpur. Retrieved from <https://www.youtube.com/watch?v=611Hq-W5bwc>. [Accessed 14 November 2015].

Griegel, D. J. and Reeves, M. (1985). "Prevention of Post Harvest Food Losses: a Training Series". Food and Agriculture Organization of United States. Rome, 1985.

Gottschalk, K. and Christenbury, G. D. (1998). A model for predicting heat and mass transfer in filled pallet boxes. Paper (Paper No. 983157) presented at the ASAE Annual International Meeting in Orlando, Florida on 12-16 July, 1998.

Gould, W. A. (1999). "Potato Production, Processing and Technology". CTI Publications, Inc. 2 Oakway Road, Timonium, Maryland 210934247USA.

Kaiya, V. (2014). "Post-Harvest Losses and Strategies to Reduce them". Action Centre la Faim (ACF). pp.3-5. Retrieved from http://www.actioncontrelafaim.org/sites/default/files/publications/fichiers/technical_paper_phl_.pdf. (Accessed 17 November 2015).

- Karki, T.B. (2002). Post-harvest technology in Nepal. Proceedings of SAARC workshop on post-harvest technology, September 24-25, Kathmandu, Nepal, Pp. 64.
- Keasar, T., A. Kalish, O. Becher, and S. Steinberg. 2005. Spatial and temporal dynamics of potato tuberworm (Lepidoptera: Gelechiidae) infestation in field-stored. *Journal of Economic Entomology* 98: 222–228.
- Khanal, S. & Bhattarai, K. (2020). Study on Post-Harvest Losses in Potato in Different Storage Conditions. *Journal of Food Science and Technology Nepal*, 12(12), 14–19.
- Khatana, V., Kumar, P., Ilangantileke, S. and Scott, G. (1997), "Postharvest handling of potatoes by farmers in the hill areas of Uttar Pradesh, India: Problems and Prospects", working paper [1997-9], International Potato Centre, Lima, 3 February.
- Knutsson, J. (2012). Long-term storage of starch potato and its effect on starch yield Master's Thesis. Swedish Univ of Agricultural Sciences. [page??](#)
- Kozukue, N., and S. Mizuno. 1990. Effects of light exposure and storage temperature on greening and glycoalkaloid content in potato tubers. *Journal of the Japanese Society for Horticultural Science* 59: 673– 677.
- Kuyu, C.G., Tola, Y.B. and Abdi, G.G. (2019). Study on post-harvest quantitative and qualitative losses of potato tubers from two different road access districts of Jimma zone, South West Ethiopia. *Heliyon*, 5, e2272, Pp. 1-7
- Lutaladio, NeBambi, and Oscar Ortiz. Sustainable Potato Production Guidelines for Developing Countries. (2009), Retrieved from www.fao.org/3/a-i1127e.pdf.
- Mani, F., Hannachi, C., 2015. Physiology of potato sprouting. *Journal of New Sciences, Agriculture and Biotechnology*, 17 (2), Pp. 591-602.
- Marwaha, R. S. et al., (2010). "Potato Processing Scenario in India: Industrial Constraints, Future Projections, Challenges Ahead and Remedies — A Review." *Journal of Food Science and Technology*, vol. 47, no. 2, pp. 137–156., doi:10.1007/s13197-010-0026-0.
- MoALD. (2020/21). Statistical Information on Nepalese Agriculture. Ministry of Agriculture and Livestock Development, Government of Nepal. 2022.
- Mutandwa, E. and Gadzirayi, C. T. (2007). Comparative Assessment of Indigenous Methods of Sweet Potato Preservation among Smallholder Farmers: Case of Grass, Ash and Soil based Approaches in Zimbabwe. *African Studies Quarterly*. 9 (3).
- Nipa, J & Roy, Tuhin & Amin, A.K.M. & Hasanuzzaman, Md. (2013). Effect of Lifting Time and Tuber Size on Ambient Storage Performance of Potato Derived from True Potato Seed. *Int. J. Sustain. Agric.* 5. 1-9. 10.5829/idosi.ijsa.2013.05.01.314.
- "Potato Vocabularies Explained". (9 Mar. 2020), Retrieved from www.npck.org/potato-vocabularies-explained-2/.
- Rhoades, R. E. (1985). Traditional potatoes production and farmer's selection of varieties in eastern Nepal [Report]. International Potato Center. Retrieved from http://pdf.usaid.gov/pdf_docs/PNAAT487.pdf. [Accessed 10 April, 2015]

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Sedera, F. A., & Shetata, S. (1994). Effect of irrigation frequency and N, K level on growth, yield, chemical composition and storage ability of potato. *Zagazig Journal of Agriculture Research*, 21, 129-149

Formatted: Font: Italic

Singh, B. (2013). Potato storage and physiology of seed. In: "Summer School on Advances in Quality Potato Production and Post-harvest Management". Shimla - 171 001 H.P India. Central Potato Research Institute, Shimla 171 001.

Sudheer, K. P. and Indira, V. (2007). "Post-Harvest Technology of Horticultural Crops". *New India publishing agency*. Pritam pura, New Delhi.

Formatted: Font: Italic

Tanios S, Eyles A, Tegg R and Wilson C. (2018). Potato tuber greening: a review of predisposing factors, management and future challenges. *Am. J. Potato Res.* 2018;95: 248–57.

Theisen, K. and Campilan, D. (2010). International Potato Centre: world potato atlas. Retrieved from <https://research.cip.cgiar.org/confluence/display/wpa/Nepal>. (Accessed 20 October, 2015).

www.fao.org/docrep/008/a018e/a01850c.htm