

## Distribution of dinoflagellates along Purnagad estuary, Ratnagiri, Maharashtra

### Abstract

A study was carried out in the Purnagad estuary, southwest coast of India, to find out the distribution of dinoflagellates. Samples were collected for one year period during 2022–2023. During the study 22 species of dinoflagellates were encountered among which *Ceratium fusus* was dominant. Relatively high density and diversity of dinoflagellates were discovered in the month of May as compared to the other months. Temperature showed a positive correlation with the dinoflagellate community showing its importance in dinoflagellate growth.

**Keywords.** Dinoflagellates. Species diversity. estuary

### 1.0 Introduction


In the near-shore and continental shelf areas of the marine environment, dinoflagellates are an important group of protists that are found in various kinds of aquatic habitats, next to diatoms (Sahu et al., 2014). Dinoflagellates are a type of primarily unicellular creatures that belong to the division or Phylum Pyrrophyta. They are distinguishable from other groups by several unique qualities, including organelles, coloring, flagellar insertion, and features of the nucleus. Dinoflagellates and ciliates are the two most diverse groups of algae. There are currently about 2400 identified species of dinoflagellates (Gomez, 2012b). Environmental molecular surveys uncover more groups of primitive dinoflagellates that are not as well-documented, and tens of thousands of new species are published each year (Guillou et al., 2008). Dinoflagellates are neither plants nor animals, many of them exhibit characteristics associated with plants, including the ability to photosynthesize, have walls made of cellulose, and produce starch, which is used as an energy storage material (Carty et al., 2015).

Estuaries are highly productive, dynamic, semi-enclosed water bodies that are intermittently or permanently connected to the sea. They are nourished by freshwater from river inflows, which produces a unique salinity gradient and unique biota characteristics (Elliott and Mclusk, 2002; Mclusk and Elliott, 2007; Whitfield et al., 2002; Chicharo et al., 2006; Tweedley et al., 2019). The fluctuations in river water mixing with seawater, which produce turbidity, nutrient gradients, and salinity (Svetlichny, 2019), indicate the complexity of a river estuary. Estuaries are dynamic environments that change constantly. They have been recognized as dynamic ecosystems due to their physical characteristics. They get an influx of freshwater during the monsoon, which results in periodic variations in the concentration of nutrients and salinity. A salinity gradient consequently develops along the estuary. A very poor paper has been published on the physico-chemical characteristics in relation to the dinoflagellates in Purnagad estuary (Map. 1) Ratnagiri. Hence the present study was conducted to study the monthly distribution of dinoflagellates and physico-chemical parameters of water in the Purnagad estuary, southwest coast of India.

**Comment [H11]:** Please adjust it to the journal template!

## 2.0 Material and Methods

### 2.1. Study area and sampling



**Map.1** Sampling locations at Purnagad estuary, Ratnagiri

The west coast of India experiences intense rainfalls during the monsoon months of June–September, the summer season (February–May), and winter season (October–January). The study was carried out in the estuarine and coastal locations of Purnagad. Based on the salinity gradient, sampling was conducted at five sampling stations (Map. 1) in Purnagad estuary. The tidal amplitude Surface samples were collected every month from these locations during the high tide times from February 2022 to January 2023.

2.2. Environmental parameters-Field data like atmospheric and surface water temperatures, salinity, dissolved oxygen, and pH were measured during the sampling. Nutrients like inorganic phosphate, nitrate, nitrite, and silicate, were analyzed by adopting the standard methods (Strickland and Parsons, 1972; AOAC, 2006; APHA, 2005; Andrew et al., 2005 and Rice et al., 2012).

2.3. Dinoflagellate assemblage - Monthly samplings for quantitative and qualitative analyses of dinoflagellates were collected from the estuarine water surface and filtered by using the 60 $\mu$  plankton net. The collected dinoflagellate samples were preserved in Lugols iodine and 5% neutralized formalin solution for counting dinoflagellate cells and identification of genera and species Sedgwick-Rafter plankton counting chamber and examined microscopically. Cells were enumerated and expressed as no.l<sup>-1</sup>. (Santhanam et al., 1987, Newell and Newell, 1963, Yamaji, 1979, Wood, 1968, Claudia et al., 2017). Biodiversity indices such as the Shannon index, species richness, evenness and dominance (Bakus, 2007)

### 3.0 Results and Discussion

#### 3.1. Physico-chemical parameters

Atmospheric temperature was maximum in May (37.2 °C) and minimum in the month of January (21.9 °C). Water temperatures with the minimum during the winter in the month of December (25.4 °C) and a maximum of 37 °C was recorded during the summer month of April. The maximum salinity of 37.6psu was found in May and zero salinity was recorded during monsoon season in the months of July and August. The pH was recorded maximum (8.8) in the month of April while minimum in the month of August. The highest dissolved oxygen was observed in the month of February and lowest in the months of April and May respectively (Graph 1).

Nutrients like, maximum nitrate concentration (0.1796 mg l<sup>-1</sup>) were found in the month of January while minimum (0.1747 mg l<sup>-1</sup>) in the month of October. The lowest nitrite concentration (0.0076 mg l<sup>-1</sup>) was recorded in the month of June while the highest (0.0228 mg l<sup>-1</sup>) in the month of January. Phosphate values varied from 0.1038 mg l<sup>-1</sup> to 0.3899 mg l<sup>-1</sup> in the month of February to January respectively. Silicate was maximum (0.6031 mg l<sup>-1</sup>) in the month of February and minimum (0.0050 mg l<sup>-1</sup>) in the month of October (Graph 1).

#### 3.2 Qualitative and quantitative distribution of dinoflagellates along Purnagad estuary

*Ceratium breve* was uncommonly found in the month of May and June, similarly in Cochin backwater (Gopinathan, 1972), *Ceratium furcatus* was infrequently found in the month of May, similar result found in Cochin estuary (Dayala et al., 2014). *Ceratium fusus* (14%) was recorded maximum in numbers in the months of Feb, May, June, July and November, with similar results observed along Tuticorin (Asha et al 2018). *Ceratium fusus* (203 no. l<sup>-1</sup>) was found to be the most abundant species at Purnagad estuary, due to wide ranges of water temperatures, salinities along Kerala coast (Rajashekhar and Rai, 2014). *Ceratium gibberum* was occasionally found in the month of May, similarly in the Cochin water (Sanilkumar, 2009). *Ceratium tripos* was infrequently observed in the months of May and July similar result were found along the Mahanandi estuary (Naik et al., 2009). *Dinophysis caudata* (9%) was rarely found in the months of Feb, May, and January, a similar result was found along the Nethravati – Gurupura estuary (Shruthi and Rajashekhar, 2013). *Dinophysis tripos* (8%) was observed in the months of Feb and November, similarly along Cochin water (Sanilkumar, 2009). *Gymnodinium catenatum* (9%) was dominated in the month of May, similarly along southern Brazil (Tavares et al., 2009). *Ornithocercus magnificus* was found maximum in number in the month of June, similar result reported by (Lavanya Ratheesh et al. 2020) along Kochi waters. *Peridinium* sp. was found in the month of January. *Peridinium oceanicum* was recorded in the month of July and November, similar result found along Bay of Bengal (Mishra et al., 2005). *Peridinium ovatum* (11%) was second abundant species (166 no. l<sup>-1</sup>) in the months of May and November along Purnagad estuary (Yamaji, 1979). *Peridinium rectum* was observed only in the month of November, similarly along Veraval coast, Gujarat (Temkar et al., 2015). *Prorocentrum compressum* was observed in the months of May, July and November, similarly in Cochin water (Sanilkumar, 2009). *Prorocentrum micans* was reported in the months of February and May corresponding

result found in Nethravati – Gurupura estuary (Shruthi and Rajashekhar, 2013). *Protoberidinium crassipes* was found in the months of May and November, similarly in Georgia, Russian Federation and Ukraine, (Slobodnik et al, 2018). *Protoberidinium latissimum* (7%) was found in the months of July, October and November, similarly along the southwestern Gulf of Mexico (Okolodkov et al., 2008). *Protoberidinium quinquecorne* was recorded in the month of November, similarly along Cochin estuary (Dayala et al., 2014). *Protoberidinium venustum* in the month of June, similar result found along Vietnamese coastal waters (Luom et al., 2017). *Pyrocystis lunula* was observed in the month of July, similar result along East Coast of India (Sahu et al., 2013). *Pyrocystis noctiluca* was reported in the month of May, similar result found along Mandovi estuary, Goa (Pednekar et al., 2014). *Pyrophacus horologium* was observed in the months of May, June and July, similar result were recorded along Cochin water (Sanilkumar, 2009), (Table no.1), (Fig.1).

Dinoflagellates were dominant in the month of May (509 no.l<sup>-1</sup>) both quantitatively and qualitatively (Sahu et al., 2014). It was found that high temperature and salinity are favorable to the growth of dinoflagellates (Taylor, 1973), (Graph 2), (Graph 3), (Fig. 1).

**Comment [H12]:** We recommend stating in ind. l<sup>-1</sup>

**Table no.1-** Monthly plankton abundance along Purnagad estuary during February 2022-January 23

Species	Months	F-22	M-22	A-22	M-22	J-22	JI-22	A-22	S-22	O-22	N-22	D-22	J-23
<i>Ceratium breve</i>		-	-	-	+	+	-	-	-	-	-	-	-
<i>Ceratium furca</i>		-	-	-	+	-	-	-	-	-	-	-	-
<i>Ceratium fusus</i>		+	-	-	+	+	+	-	-	-	+	-	-
<i>Ceratium gibberum</i>		-	-	-	+	-	-	-	-	-	-	-	-
<i>Ceratium tripos</i>		-	-	-	+	-	+	-	-	-	-	-	-
<i>Dinophysis caudata</i>		+	-	-	+	-	-	-	-	-	-	-	+
<i>Dinophysis tripos</i>		+	+	-	-	-	-	-	-	-	-	-	-
<i>Gymnodinium catenatum</i>		-	-	-	+	-	-	-	-	-	-	-	-
<i>Ornithocercus serratus</i>		-	-	-	-	+	-	-	-	-	-	-	-
<i>Peridinium</i> sp.		-	-	-	-	-	-	-	-	-	-	-	+
<i>Peridinium oceanicum</i>		-	-	-	-	-	+	-	-	-	+	+	-
<i>Peridinium ovatum</i>		-	-	-	+	-	-	-	-	-	+	-	-
<i>Peridinium rectum</i>		-	-	-	-	-	-	-	-	-	+	-	-
<i>Prorocentrum compressum</i>		-	-	-	+	-	+	-	-	-	+	-	-
<i>Prorocentrum micans</i>		+	-	-	+	-	-	-	-	-	-	-	-
<i>Protoberidinium crassipes</i>		-	-	-	-	-	-	-	-	-	+	-	-
<i>Protoberidinium latissimum</i>		-	-	-	-	-	+	-	-	-	+	-	-
<i>Protoberidinium quinquecorne</i>		-	-	-	-	-	-	-	-	-	+	-	-
<i>Protoberidinium venustum</i>		-	-	-	-	-	+	-	-	+	+	-	-
<i>Pyrocystis lunula</i>		-	-	-	-	-	+	-	-	-	-	-	-
<i>Pyrocystis noctiluca</i>		-	-	-	+	-	-	-	-	-	-	-	-
<i>Pyrophacus horologium</i>		-	-	-	+	+	+	-	-	-	-	-	-

+(Present), -(Absent)

### 3.3 Correlation with environmental parameter

**Table no. 2-** Correlation between physico-chemical parameters and dinoflagellates along Purnagad estuary during February 2022-January 23

A.T- Atmospheric temperature, W.T- water temperature, S- salinity, DO-dissolved oxygen, NO<sub>3</sub>-nitrate, NO<sub>2</sub>-nitrite, PO<sub>4</sub>- phosphate, Si- silicate, D-Dinoflagellates


**Correlations**

	A.T	W.T	S	pH	DO	NO <sub>3</sub>	NO <sub>2</sub>	PO <sub>4</sub>	Si	D
A.T	1									
W.T	.782**	1								
S	-0.1	0.045	1							
pH	.604*	.650*	0.402	1						
DO	-0.063	-0.218	-.639*	-0.38	1					
NO <sub>3</sub>	-0.354	-0.343	0.365	-0.067	0.232	1				
NO <sub>2</sub>	-.576*	-0.407	0.461	-0.158	0.001	0.548	1			
PO <sub>4</sub>	-.648*	-0.522	0.422	-0.331	0.091	0.486	.813**	1		
Si	0.26	-0.122	0.104	0.306	0.067	0.412	-0.28	-0.396	1	
D	.625*	.640*	0.364	.672*	-0.1	0.023	-0.215	-0.272	0.314	1

\*\* . Correlation is significant at the 0.01 level (2-tailed).


\* . Correlation is significant at the 0.05 level (2-tailed).

Environmental factors such atmospheric temperature showed correlation with dinoflagellates( $r = 0.625$ ) (Sahu et al., 2014). pH was correlated with dinoflagellate ( $r = 0.672$ ) (Yoo, 1991). Salinity was negatively correlate with dissolved oxygen ( $r = -0.639$ ) (Supriatna and Mahmudi, 2021 and Rajashekhar and Rai, 2014). The phosphate was positively correlate with nitrate ( $r = 0.813$ ), relationship of nitrate and phosphate was favourable for phytoplankton abundance (Marsela et al., 2021), (Table no.2).




**Graph 1-** Monthly water parameters recorded along Purnagad estuary, during 2022-23


A.T- Atmospheric temperature, W.T- water temperature, S- salinity, DO-dissolved oxygen



**Graph 2-** Monthly quantitative variations of dinoflagellates recorded during 2022-23



**Graph 3-** Monthly quantitative variations of dinoflagellates recorded during 2022-23




**Fig.1-** Percentage distribution of different phytoplankton divisions in the Purnagad estuary during 2022-23.

### 3.4 Species richness and diversity indices

**Comment [H13]:** It is worth discussing further why fluctuations occur

The range of species diversity, dominance, evenness and richness were 0 - 3.4096, 0.0644-1, 0-0.9356, 0 – 18 respectively. The least values of diversity indices were recorded during March, April, August and September but were higher during other periods. Higher values of dominance were reported in the months of March, April, August and September while minimum in the months of February, July and October. The maximum richness values were recorded in the months of May, July and November. The low richness was recorded April, August and September. Graph 2 (Graph 4).



**Graph 4** – Monthly variations of dinoflagellates species diversity, dominance, evenness and richness recorded during 2022-23.

### 4.0 Conclusion

The dinoflagellate community structure at Purnagad estuary was found to constitute 22 species during the one year of study. The species that was found to be most prevalent was *Ceratium fusus*. *Prorocentrum* (six species) was the most diversified genus among all the species, followed by *Ceratium* (five species). Temperature was one of the important parameters that positively influenced the dinoflagellate community.

### 7.0 Reference



- Andrew D. E., Lenore. S. C., Eugene W. R. and Arnold E. G. Standard method for the examination of water and wastewater. Centennial edition, 2005;1-1: (In press).
- AOAC. Official Methods of Analysis, 18th ed. Association of Official Analytical Chemist, Washington, D. C. 1094, 2006.
- APHA. Standard Methods for the Examination of Water and Waste Water 21st ed. American Public Health of Association, American Water Works Association Environmental Federation, Washington DC, USA. 2605, 2005.
- Asha PS, Ranjith L, Diwakar K, Prema D and Krishnakumar PK. Distribution and species diversity of phytoplankton in the inshore waters of Tuticorin in relation to the physicochemical variables. J. Mar. Biol. Ass. India, 2018; 60:1.
- Bakus G J. Quantitative analysis of marine biological communities, ISBN-13; 978-0- 470-04440-7. ISSN-10; 0-470-04440-3.143-148, 2007.
- Carty S and Parrow MW. Dinoflagellates. Freshwater Algae of North America. 2015: (In press).
- Chicharo L, Chicharo MA, Ben-Hamadou R. Use of a hydrotechnical infrastructure (Alqueva Dam) to regulate planktonic assemblages in the Guadiana estuary: Basis for sustainable water and ecosystem services management. Estuar. Coast. Shelf Sci., 2006;70: (In press).
- Claudia C, Neelke F, Stijn V, Dries V and Jessica DM. Intellectual and developmental disabilities, 2017; 55(4): (In press).
- Dayala VT, Salas PM and Sujatha CH. Spatial and seasonal variations of phytoplankton species and their relationship to physicochemical variables in the Cochin estuarine waters, Southwest coast of India. Indian Journal of Geo-Marine Sciences. 2014;43(6): (In press).
- Elliott M, McLusky DS. The need for definitions in understanding estuaries. Estuar. Coast. Shelf Sci., 2002; 55: (In press).
- Gopinathan CP. Seasonal abundance of phytoplankton in the Cochin backwater. J mar. biol. Ass. India. 1972;14 (2): (In press).
- Gomez, FA. checklist and classification of living dinoflagellates(Dinoflagellata, Alveolata). CICIMAR Océanides.2012b; 27: (In press).
- Guillou L, Viprey M, Chambouvet A, Welsh RM, Kirkham AR, Massana R, Scanlan DJ. and Worden AZ. Widespread occurrence and genetic diversity of marine parasitoids belonging to Syndiniales (Alveolata). Environmental Microbiology, 2008;10: (In press).
- Lavanya R, Joseph RV, Parvathy R, Shelton P, Prema D, Kripa V and Kaladharan P. First report of a rare bloom of *Ornithocercus magnificus*, Stein 1883 along the coastal waters of Kochi; A possible indicator of increasing sea surface temperature. J. Mar. Biol. Ass. India, 2020;62 (2): (In press).



- Marsela K, Hamdani H, Anna Z and Herawati H. The Relation of Nitrate and Phosphate to Phytoplankton Abundance in the Upstream Citarum River, West Java, Indonesia. Asian Journal of Fisheries and Aquatic Research, 2021;11(5): (In press).
- Mcluskys DS, Elliott M. Transitional waters: A new approach, semantics or just muddying the waters? Estuar. Coast. Shelf Sci., 2007, 71, 359–363 (In press).
- Mishra S, Sahu G, Mohanty AK, Singh SK and Panigrahy RC.. Impact of the Diatom *Asterionella glacialis* (Castracane) Bloom on the Water Quality and Phytoplankton Community Structure in Coastal Waters of Gopalpur Sea, Bay of Bengal. Asian Journal of Water, Environment and Pollution, 2005; 3(2): (In press).
- Naik S, Acharya BC and Mohapatra A. Seasonal variations of phytoplankton in Mahanadi estuary, east coast of India. Indian Journal of Marine Sciences. 2009; 38:2: (In press).
- Newell GE and Newell RC. Marine Plankton, A practical guide. London, 1963: (In press).
- Pednekar SM, Kerkar V, Matondkar SGP. Spatiotemporal distribution in phytoplankton community with distinct salinity regimes along the Mondovi estuary, Goa, India. Turkish Journal of Botany, 2014; 38: (In press).
- Rajashekhar M and Rai SV. Seasonal assessment of hydrographic variables and phytoplankton community in the Arabian sea waters of Kerala, southwest coast of India. Brazilian journal of oceanography, 2014; 62(4): (In press).
- Rice EW, Baird RD, Eaton AD and Clesceri LS. Standard methods for the examination of water and wastewater. American Public Health Association, NW, Washington. 2012: (In press).
- Sahu G, Mohanty AK, Samantara MK, Satpathy KK. Seasonality in the distribution of dinoflagellates with special reference to harmful algal species in tropical coastal environment, Bay of Bengal. Environmental Monitoring and Assessment 2014; 186(8).
- Santhanam R, Ramanathan N, Venkatramanujam K and Jagatheesan G. Phytoplankton of the Indian seas. Daya Publishing house, Delhi, India, 1987: (In press).
- Sanilkumar MG. Microalgae in the southwest coast of India. Cochin University of Science and Technology. 2009; 1-285: (In press).
- Shruthi MS and Rajashekhar M. Ecological observations on the phytoplankton of Nethravati–Gurupura estuary, south west coast of India. J. Mar. Biol. Ass. India, 2014; 55 (2): (In press).
- Slobodnik J, Alexandrov B, Komorin V, Mikaelyan A, Guchmanidze A, Arabidze M, Korshenko A. National Pilot Monitoring Studies and Joint Open Sea Surveys

- in Georgia, Russian Federation and Ukraine, 2017. Scientific Report – Joint Black Sea Surveys, 2018.
- Strickland JDH., and Parsons TR. A Practical Handbook of Seawater Analysis. Bull. Fish. Res. Bd. Can, 1972; 167-310.
- Supriatna and Mahmudi M. Dynamic model of dissolved oxygen in intensive concrete pond of white leg shrimp (*Litopenaeus vannamei*) in Bomo Village, East Java. Earth and Environmental Science. 2021:919, (In press).
- Svetlichny L, Hubareva E, Khanaychenko A, Uttieri M. Response to salinity and temperature changes in the alien Asian copepod *Pseudodiaptomus marinus* introduced in the Black Sea. J. Exp. Zool., 2019; 331:(In press).
- Temkar GS, Gangan SS, Azeez PA, Sikotaria KM, Brahmane VT, Metar SY, Mathew KL and Desai AY. Correlation of phytoplankton density with certain hydrological parameters along the coastal waters of Veraval, Gujarat. Journal of the Marine Biological Association of India, 2015; 57:2.
- Tweedley JR, Dittmann SR, Whitfield AK, Withers K, Hoeksema SD, Potter IC. Hypersalinity: Global distribution, causes, and present and future effects on the biota of estuaries and lagoons. In Coasts Estuaries; Elsevier: Amsterdam, The Netherlands, 2019; 30: (In press).
- Taylor FJR. General features of dinoflagellate material collected by the Anton Bruun during IIOE. In: Zeitzschel, B. (Ed.). The Biology of the Indian Ocean: Biological Studies, Springer Verlag, Berlin, 1973:(In press).
- Whitfield AK, Elliott M. Fishes as indicators of environmental and ecological changes within estuaries: A review of progress and some suggestions for the future. J. Fish Biol., 2002; 61:(In press).
- Wood EJF. Dinoflagellates of the Caribbean Sea and Adjacent Areas. University of Miami Press. United States of America, 1968: (In press).
- Yamaji I. Illustration of the marine plankton of Japan. Hoikusha Publishing Co. Ltd, 1979: (In press).
- Yoo KI. Population dynamics of dinoflagellate community in Masan Bay with a note on the impact of environmental parameters. Mar. Pollut. Bull, 1991; 23: (In press).
- Zar JH. Biostatistical analysis. Fourth ed. Ten prints (I) Pvt. Ltd. Delhi, India, 2004: (In press).