

IMPACT OF SEASONAL CHANGES ON AVIFAUNA DIVERSITY IN NGURU WETLAND,
YOBE STATE, NIGERIA.

ABSTRACT

The research on impact of seasonal changes on avifauna diversity in Nguru Lake of Hadejia-Nguru Wetland was carried out in two different seasons for the period of six months, from April to June (dry season) and July to September (wet season). The aim was to evaluate the impact of seasonal changes on avifauna diversity and the species richness in the study area. A reconnaissance survey was carried out to select study sites and sampling points. Five (5) different points were selected which comprised of Madauka, Kayayyashi, Farin Ruwa, Badum respectively and Point count census techniques were adopted. In each counting station, bird observation was carried out twice daily; morning between 6:00 am and 10:00 am and evening between 4:00 pm and 6:00 pm. Data compilation was done using Microsoft Excel and were analysed using statistical software Past 326b. Bird diversity was calculated using Shannon-Weiner diversity index. The finding of this research showed that the majority of bird species identified were Resident (12,225), Migratory (8,186) and Palearctic Migrant (10,876) species in the study area. Dominant bird species in Nguru Lake were *Philomachus pugnax*, *Actophilornis africanus* and *Phalacrocorax africanus*. A total of 5,538 birds' species was identified (Dry Season) in Nguru Lake "belonging to 21 families in which Ardeidae have the highest (7) species of bird. Shannon Diversity Index' has 2.55 in dry season, while in Wet Season' has 3.40. This indicated that there was relatively equal and high diversity of bird species in Wet Season in the study area and has the total species richness of 6,711. Overgrazing and other anthropogenic activities which affect the population diversity of avifauna species serve as major threats to the continued existence of Wetlands. Typha grass also remained pressing problem which create microhabitats reducing critical habitat for feeding, nesting and roosting habitat for birds, as well as hamper the smooth flow of water used by many water bird species. In view of the findings of this study, effective monitoring and strategies for conservation to restore the population of bird species, provide public enlightenment of the people around the wetland and management of Hadejia-Nguru Wetland promptly advice the significant importance of the wetland in serving as home for resident and migratory bird

species.

Key Words: Impact, Seasonal, Changes, Avifauna, Diversity, Reconnaissance, Survey, Sampling.

INTRODUCTION

Avifauna is a general name for bird species. Birds are feathered, winged, egg-laying vertebrates. They belong to the Kingdom "Animalia," Phylum "Chordata" and Class "Aves". They have a worldwide distribution, living in and around oceans, rivers, forests and mountains. They are the most noticeable group in the animal kingdom. Their bright colours, distinct songs and calls, and showy displays add fun to human life. Many people derive great pleasure from watching birds and listening to their beautiful songs. Birds are social animals that communicate with visual signs, calls and songs. They display social behaviours such as cooperative breeding and hunting, flocking and mobbing of predators. Birds live and breed in most terrestrial habitats and on all the seven continents. Nigeria is blessed with many species of birds scattered throughout the different ecological regions. As with any natural habitat, wetlands are important in supporting bird species diversity. Wetlands provide food for birds in the form of plants, vertebrates, and invertebrates. Some birds forage for food in the wetland soils, while some find food in the water column and others feed on the vertebrates and invertebrates that live on submerged and emergent plants. (Labe et al., 2018).

Studies on bird diversity by Burgess et al. (2002), Doggart et al. (2005), Frontier-Tanzania (2005), and Yanda and Munishi (2007) in the Uluguru area were confined to the forest emphasis in the general negative effects of forest conversion to human dominated habitats. Nevertheless, human dominated and agricultural habitats vary a lot and therefore the effect on birds can be very different (Tworek, 2002). Responses of birds to habitat changes differ depending on their strategies, some lifestyles benefit from habitat change, while for others it is a principal threat (Tworek, 2002). Birds are



very visible and integral part of the ecosystem that occupy many trophic levels in a food chain ranging from consumer to predators. Their occurrence has been helpful as an environmental health indicator, plant pollinators and seed dispersal as well as a pest controller (Hadley et al., 2012; Ramchandra, 2013). This research work identified avifauna species and determined its diversity in the study area.

MATERIALS AND METHOD

Study Area

This research was carried out at Hadejia-Nguru Wetlands (HNWs) located in the Sahelian Zone of Nigeria at a point where River Hadejia and flow through a fossil dune field before converging and draining into Lake Chad. It lies between latitude $12^{\circ}13'N$ to $12^{\circ}55'N$ and longitude $10^{\circ}15'E$ to $11^{\circ}30'E$ (Ayeni et al., 2019), covering three Nigerian states namely, Bauchi, Jigawa and Yobe (Figure 1). The Hadejia-Nguru Wetlands named after two major towns (Hadejia and Nguru) are extensive floodplain wetlands in the dry land of northern Nigeria which are surrounded by many villages. It supports a wider range of biodiversity and livelihood activities. (Ayeni et al., 2019).

The wetlands extend for approximately 120 km from west to east within Jigawa State and for a further 60-70 km down-stream in adjacent Yobe State (Caro et al., 2011). In width, the wetlands range from 10 km to more than 50 km from north to south, with approximately 8000 km^2 of floodplain. The extent of the floodplain varies considerably from year to year depending on rainfall and complex interactions of river flow, dam releases, flood regimes and topography. (Bourn, 2003).




Figure1:Map of Nigeria Showing the Study Area.

Source: (Ayeni et al., 2019)

Research Design

A reconnaissance survey was carried out with the aid of topographic map and

personnel assistant to select study sites and sampling points in the field after obtaining permission from the authorities of Nguru Wetland. The survey was carried out at study area and five (5) points were been selected namely; Madauka, Kayayyashi, Farin Ruwa, Badum and Guzan lakes respectively. The survey was carried out for the period of six (6) months (three months for dry season and three months for wet season. Birds' observation was carried out once in a month. Birds were counted as birds seen and heard around the area and birds in flight were also be counted within 100 meters radius to aid data collection. A pair of binoculars with magnification 988,000m Model 750 was used for birds viewing alongside with field guide book "Birds of Western Africa" by Borrow and Demey (2014) for identification of birds (Ramsar Convention Bureau, 2000). A Komery Digital Camera 24MP was also used for taking photographs of birds.

Method of Data Collection

Point count census techniques as outlined by Bibby et al. (1992) and Ralph et al. (1993) were adapted to count the birds for the study. Five (5) different points were selected randomly from each study sites, namely; Madauka, Kayayyashi, Farin Ruwa, Badum and Guzan.

In each counting station, bird observation was carried out twice daily; morning between 6:00 am and 10:00 am and evening between 4:00 pm and 6:00 pm. On arrival at each counting station, the observers stayed for five (5) minutes before beginning to count. This was to enable the birds to settle down following disturbance by the arrival of the observers. The observers at each counting point recorded the entire birds encountered for a period of ten (10) minutes with the aid of binoculars alongside with field guide book "Birds of Western Africa" by Borrow and Demey (2014). Data forms were restructured for ease of recording all the information.

Data Analysis

Compilation and analysis of data was done using Microsoft Excel (2013 version). Data were analysed using statistical software Past 3.26b.

i. Table was used to present the types of avifauna species identified in the study area



for rainy season and dry season according to Lammeed (2011).

ii. Avifauna species diversity was calculated using Shannon-Weiner diversity index, H:

$$H = - \sum_{i=1}^S P_i \ln P_i$$

Where:

H = Shannon-Weiner Diversity Index

S = Total number of species of the community (number seen and heard).

P_i = Proportion of each or individual (1th) species in the sample.

ln P_i = Natural Logarithm of the species proportion. (Lammeed, 2011).

RESULTS

Avifauna Species Identified in Nguru Wetland (NGW) in Dry Season

The result of the avifauna species identified in 'NGW' for the 1st, 2nd and 3rd visits are presented in Table 1. The table showed that 2,148 bird species were identified in the first visit, 1,658 bird species were identified in the second visit and 1,732 species were identified in the third visit (Total = 5,538) belonging to 21 different families in dry season in the study area. The families of Bucerotidae, Coraciidae, Cuculidae, Motacillidae, Phalacrocoracidae, Ploceidae, Psittacidae, Pyronotidae, Recurvirostridae, Scolopacidae, Sternidae, Sturnidae and Viduidae contained 1 species each, families of Accipitridae, Alcedinidae, Charadriidae and Jacanidae contained 2 species each, family of Anatidae contained 3 species, while families of Columbidae and Rallidae contained 4 species and Ardeidae contained the highest number of 7 species in the study area.

Avifauna Species Identified in Nguru Wetland (NGW) in Wet Season

The result of the avifauna species identified in the study area for the 1st, 2nd and 3rd visits are presented in Table 2. The table shows that 690, 281 and 202 bird species (Total = 1,173) were identified in the first, second and third visits respectively during rainy season, different families in the study area. The families of Bucerotidae,

Coraciidae, Cuculidae, Motacillidae, Phalacrocoracidae, Ploceidae, Psittacidae, Pyronotidae, Recurvirostridae, Scolopacidae, Sternidae, Sturnidae and Viduidae contained a single bird species each, meanwhile the families of Accipitridae, Alcedinidae, Charadriidae and Jacanidae contained 4 bird species each where only Anatidae family contained 3 species of birds, the families of Columbidae and Rallidae contained 4 bird species while Ardeidae contained 7 different bird species in the study area during wet season.

UNDER PEER REVIEW



Table1: Avifauna Species Identified in the Study Area (Nguru Wetland) for Dry Season

S/ N	Family	Scientific Name	Common Name	Frequency			Total
				1 st Visit	2 nd Visit	3 rd Visit	
1.	Accipitridae	Circus ranivorus	African Marsh Harrier	3	2	1	6
	“	Milvus aegyptius	Yellow-billed Kite	4	2	10	16
2.	Alcedinidae	Halcyon leucocephala	Grey-headed Kingfisher	3	-	2	5
	“	Ceryle rudis	Pied Kingfisher	1	6	19	26
3.	Anatidae	Nettion auritus	African Pygmy Goose	-	84	84	
	“	Dendrocygna bicolor	Fulvous Whistling Duck	9	7	8	24
	“	Dendrocygna viduata	White-face Whistling Duck	558	26	117	701
4.	Ardeidae	Egretta ardesiaca	Black Heron	3	28	2354	
	“	Bubulcus ibis	Cattle Egret	24	22	12	58
	“	Ardea cinerea	Grey Heron	14	19	12	45
	“	Egretta intermedia	Intermediate Egret	1	-	-	1
	“	Egretta garzetta	Little Egret	10	17	33	60
	“	Ardea purpurea	Purple Heron	8	19	17	44
	“	Ardeola alioides	Squacco Heron	128	88	65	281
5.	Bucerotidae	Tockus erythrorhynchus	Red-billed Hornbill	-	2	-	2
6.	Charadriidae	Vanellus tectus	Black-headed Lapwing	9	11	-	20
	“	Vanellus spinosus	Spur-winged Lapwing	-	15	69	84
7.	Columbidae	Streptopelia decipiens	African Mourning Dove	18	24	25	67
	“	Streptopelia senegalensis	Laughing Dove	15	14	7	36
	“	Columba guinea	Speckled Pigeon	34	18	2779	

	“	“	Streptopelia vinacea	Vinaceous Dove	22	34	32	88
8.	Coraciidae		Coracias abyssinica	Abyssinian Roller	3	-	2	5
9.	Cuculidae		Centropus senegalensis	Senegal Coucal	2	5	4	11
10.	Jacanidae		Actophilornis africanus	African Jacana	245	184	475	904
	“	“	Micropodops capensis	Lesser Jacana	14	22	114	150
11.	Motacillidae		Motacilla flava	Yellow Wagtail	141	94	10	245
12.	Phalacrocoracidae		Phalacrocorax africanus	Longtail/Reed Cormorant	227	109	383	719
13.	Ploceidae		Ploceus cucullatus	Village Weaver	10	-	40	60
14.	Psittacidae		Psittacus krameri	Rose-ringed Parakeet	-	7	-	7
15.	Pyronotidae		Pyronotus barbatus	Common Bulbul	4	-	8	12
16.	Rallidae		Amaurornis flavirostris	Black Crake	9	4	16	29
	“	“	Gallinula chloropus	Common Moorhen	32	45	78	155
	“	“	Gallinula angulata	Lesser Moorhen	-	-	10	10
	“	“	Porphyrio madagascariensis	Purple Swamp Hen	6	4	19	29
17.	Recurvirostridae		Himantopus himantopus	Black-winged Stilt	-	8	-	8
18.	Scolopacidae		Philomachus pugnax	Ruff	550	770	10	1330
19.	Sternidae		Sterna nilotica	Gull-billed Tern	36	50	-	86
20.	Sturnidae		Lamprolaima caudatus	Longtail Glossy Starling	-	2	-	2
21.	Viduidae		Vidua chalybeata	Village Indigo	5	-	-	5
Total Number of Species Per Location					2,148	1,658	1,732	5,538

Source: Field Survey, 2024

Table2:AvifaunaSpeciesIdentified intheStudyArea(NguruWetland) forWetSeason

S/N	Family	Scientific Name	CommonName	Frequency			Total
				1 st Visit	2 nd Visit	3 rd Visit	
1.	Accipitridae	Circusranivorus	AfricanMarshHarrier	1	1	-	2
	“	Milvusaegyptius	YellowbilledKite	8	6	2	16
2.	Alcedinidae	Halcyonleucocephala	Grey-headedKingfisher	7	1	-	8
	“	Cerylerudis	PiedKingfisher	24	11	7	42
3.	Anatidae	Nettapusauritus	AfricanPygmyGeese	3			
	“	Dendrocygnabicolor	FulvousWhistlingDuck	10	-	-	10
	“	Dendrocygnaiduata	WhiteFace WhistlingDuck	23	-	-23	
4.	Ardeidae	Egrettaardesiaca	BlackHeron	34	17	10	61
	“	Bubulcusibis	CattleEgret	36	9	6	51
	“	Ardeacinerea	GreyHeron	34	12	9	55
	“	Egrettaintermedia	IntermediateEgret	3	-	-	3
	“	Egrettazarzetta	LittleEgret	24	-	-	24
	“	Ardeapurpurea	PurpleHeron	13	8	7	28
	“	Ardeolaralloides	SquaccoHeron	32	12	7	51
5.	Bucerotidae	Tockuserythrorhynchus	Red-billedHornbill	9	5	5	19
6.	Charadriidae	Vanellustectus	Black-headedLapwing	25	14	10	49
	“	Vanellusspinosus	Spur-wingedLapwing	14	7	4	25
7.	Columbidae	Streptopeliadecipiens	AfricanMourningDove	16	12	9	37
	“	Streptopeliasenegalensis	Laughing Dove	20	11	11	42
	“	Columbaguinea	SpeckledPigeon	41	20	11	72
	“	Streptopeliavinacea	VinaceousDove	28	15	10	53

8.	Coraciidae	Coraciasabyssinica	AbyssinianRoller	15	4	2	21
9.	Cuculidae	Centropussenegalensis	SenegalCoucal	8	7	7	22
10.	Jacaniidae	Actophilornisafricanus	AfricanJacana	67	24	12	103
	“ “	Microparracapensis	LesserJacana	32	20	11	63
11.	Motacillidae	Motacillaflava	YellowWagtail	9	-	-	9
12.	Phalacrocoracidae	Phalacrocoraxafricanus	Longtail/ReedCormorant	28	12	11	51
13.	Ploceidae	Ploceuscucullatus	VillageWeaver	13	4	4	21
14.	Psittacidae	Psittaculakrameri	Rose-ringedParakeet	7	4	3	14
15.	Pyconotidae	Pycnonotusbarbatus	Common Bulbul	21	12	12	45
16.	Rallidae	Amauornisflavirostris	BlackCrake	13	8	8	29
	“ “	Gallinulachloropus	Common Moorhen	5	4	8	17
	“ “	Gallinulaangulata	LesserMoorhen	11	5	4	20
	“ “	Porphyriomadagascariensis	PurpleSwamphen	4	2	2	8
17.	Recurvirostridae	Himantopus	Black-wingedStilt	4	-	-	4
18.	Scolopacidae	Philomachuspugnax	Ruff	18	-	-	18
19.	Sternidae	Sternanilotica	Gull-billedTern	9	6	4	19
20.	Sturnidae	Lamprotorniscaudatus	LongtailGlossyStarling	16	3	1	20
21.	Viduidae	Viduachalybeata	VillageIndigo	5	5	5	15
TotalNumber ofSpecies PerLocation				690	281	2021,173	

Source: FieldSurvey,2024



Avifauna Species Diversity in the Research Area

The result of avifauna species diversity of Nguru Wetland during dry season is shown in Table 3. The result indicated that the Shannon Diversity Index was 2.548 for total individual species in the study area with *Philomachus pugnax* (0.34258) having the highest diversity and *Egretta intermedia* (0.00156) being the lowest diversity.

The result of avifauna species diversity of Nguru Wetland, Wet Season is shown in Table 4. The result indicated that Shannon Diversity Index was 3.40307 for total individual species in the study area with *Actophilornis africanus* (0.2136) having the highest diversity while *Circus ranivorus* (0.01087), *Egretta intermedia* and *Nettion auritus* (0.01527) respectively having the lowest diversity.



Table3: Avifauna Species Diversity in the Study Area (Nguru Wetland) for Dry Season

S/N	Species	Frequency	Pi	lnPi	Pi lnPi
1.	Actophilornis africanus	904	0.163236	1.81256	0.29587
2.	Amaurornis flavirostris	29	0.005237	5.25209	0.0275
3.	Ardeacinerea	45	0.008126	4.81273	0.03911
4.	Ardea purpurea	44	0.007945	4.8352	0.03842
5.	Ardeolaraloides	281	0.05074	2.98103	0.15126
6.	Bubulcus ibis	58	0.010473	4.55895	0.04775
7.	Centropus senegalensis	11	0.001986	6.22149	0.01236
8.	Ceryle rudis	26	0.004695	5.36129	0.02517
9.	Circus ranivorus	6	0.001083	6.82763	0.0074
10.	Columba guinea	79	0.014265	4.24994	0.06063
11.	Coracias abyssinica	5	0.000903	7.00995	0.00633
12.	Dendrocygna bicolor	24	0.004334	5.44133	0.02358
13.	Dendrocygna viduata	701	0.12658	2.06688	0.26163
14.	Egretta ardesiaca	54	0.009751	4.6304	0.04515
15.	Egretta garzetta	60	0.010834	4.52504	0.04903
16.	Egretta intermedia	1	0.000181	8.61939	0.00156
17.	Gallinula angulata	10	0.001806	6.3168	0.01141
18.	Gallinula chloropus	155	0.027988	3.57596	0.10009
19.	Halcyon leucocephala	5	0.000903	7.00995	0.00633
20.	Himantopus himantopus	8	0.001445	6.53995	0.00945
21.	Lamprolornis caudatus	2	0.000361	7.92624	0.00286
22.	Micropodops capensis	150	0.027086	3.60875	0.09775
23.	Milvusaegyptus	16	0.002889	5.8468	0.01689
24.	Motacilla flava	245	0.04424	3.11813	0.13795
25.	Nettion auritus	84	0.015168	4.18857	0.06353
26.	Phalacrocorax africanus	719	0.12983	2.04153	0.26505
27.	Philomachus pugnax	1330	0.240159	1.42645	0.34258
28.	Ploceus cucullatus	50	0.009029	4.70737	0.0425
29.	Porphyrio madagascariensis	29	0.005237	5.25209	0.0275
30.	Psittacula krameri	7	0.001264	6.67348	0.00844
31.	Pycnonotus barbatus	12	0.002167	6.13448	0.01329
32.	Sternanilia	86	0.015529	4.16504	0.06468

33.	Streptopeliadecipiens	67	0.012098	4.4147	0.05341
34.	Streptopeliasenegalensis	36	0.006501	5.03587	0.03274
35.	Streptopeliavinacea	88	0.01589	4.14205	0.06582
36.	Tockuserythrorhynchus	2	0.000361	7.92624	0.00286
37.	Vanellusspinosus	84	0.015168	4.18857	0.06353
38.	Vanellustectus	20	0.003611	5.62366	0.02031
39.	Viduachalybeata	5	0.000903	7.00995	0.00633
Total		5,538	1		2.548

Source: FieldSurvey,2024



Table4: Avifauna Species Diversity in the Study Area (Nguru Wetland) for Wet Season

S/ N	Species	Frequency	Pi	lnPi	Pi lnPi
1.	Actophilornis africanus	103	0.087809	2.43259	0.2136
2.	Amaurornis flavirostris	29	0.024723	3.70002	0.09148
3.	Ardeacinerea	55	0.046888	3.05999	0.14348
4.	Ardea purpurea	28	0.02387	3.73512	0.08916
5.	Ardeolaralloides	51	0.043478	3.13549	0.13633
6.	Bubulcus ibis	51	0.043478	3.13549	0.13633
7.	Centropus senegalensis	22	0.018755	3.97628	0.07458
8.	Ceryle rudis	42	0.035806	3.32965	0.11922
9.	Circus ranivorus	2	0.001705	6.37417	0.01087
10.	Columba guinea	72	0.061381	2.79065	0.17129
11.	Coracias abyssinica	21	0.017903	4.0228	0.07202
12.	Dendrocygna bicolor	10	0.008525	4.76473	0.04062
13.	Dendrocygna viduata	23	0.019608	3.93183	0.07709
14.	Egretta ardesiaca	61	0.052003	2.95645	0.15375
15.	Egretta garzetta	24	0.02046	3.88927	0.07958
16.	Egretta intermedia	3	0.002558	5.96871	0.01527
17.	Gallinula angulata	20	0.01705	4.07159	0.06942
18.	Gallinula chloropus	17	0.014493	4.23411	0.06136
19.	Halcyon leucocephala	8	0.00682	4.98788	0.03402
20.	Himantopus himantopus	4	0.00341	5.68103	0.01937
21.	Lamprolornis caudatus	20	0.01705	4.07159	0.06942
22.	Micropodops capensis	63	0.053708	2.92419	0.15705
23.	Milvus aegyptius	16	0.01364	4.29473	0.05858
24.	Motacilla flava	9	0.007673	4.8701	0.03737
25.	Nettion auritus	3	0.002558	5.96871	0.01527
26.	Phalacrocorax africanus	51	0.043478	3.13549	0.13633
27.	Philomachus pugnax	18	0.015345	4.17695	0.0641
28.	Ploceus cucullatus	21	0.017903	4.0228	0.07202
29.	Porphyrio madagascariensis	8	0.00682	4.98788	0.03402

30.	Psittaculakrameri	14	0.011935	4.42826	0.05285
31.	Pycnonotusbarbatus	45	0.038363	3.26066	0.12509
32.	Sternanilotica	19	0.016198	4.12288	0.06678
33.	Streptopeliadecipiens	37	0.031543	3.4564	0.10903
34.	Streptopeliasenegalensis	42	0.035806	3.32965	0.11922
35.	Streptopeliavinacea	53	0.045183	3.09703	0.13993
36.	Tockuserythrorhynchus	19	0.016198	4.12288	0.06678
37.	Vanellusspinosus	25	0.021313	3.84844	0.08202
38.	Vanellustectus	49	0.041773	3.1755	0.13265
39.	Viduachalybeata	15	0.012788	4.35927	0.05575
	Total		1,173	1	
		3.40307			

Source: FieldSurvey,2024



Table 5: Shannon-WeinerIndicesfor Avifauna SpeciesDiversityin the Study Area (NguruWetland).

Indices	DrySeason	WetSeason
	LAD	LAW
Taxa_S	39	39
Individuals	5538	1173
Dominance_DM	0.1253	0.03948
Simpson	0.8747	0.9605
Shannon_H	2.548	3.403
Evenness_e^H/S	0.3277	0.7707
Equitability_J	0.6955	0.9289

Source: FieldSurvey,2024

Key:LAD–Location'A'DryandLAW–Location'A'WetSeason

DISCUSSION

The finding of this study shows that the majority of bird species identified during the research were Resident (12,225), Migratory (8,186) and Palearctic Migrant (10,876) species in the study area. Some of the resident species recorded include, African Jacana, Longtail/Reed Cormorant, African Mourning Dove, Vinaceous Dove, Pied Kingfisher, Common Moorhen; Migratory birds include, Yellow Wagtail, African Marsh Harrier, Intermediate Egret, Red-billed Hornbill, Garganey; and the Palearctic migrant bird species were White-faced Whistling Duck, Squacco Heron, Fulvous Whistling Duck, Yellow Billed Kite and Green Sandpiper. According to Osunsina et al. (2018) the majority of birds observed during the study were resident species, migratory and Palearctic migrant species. Similar observation was made by Sabo (2016). This shows that bird species found are either residents, intra-migrants, Vagrant and Palearctic Migrant. Also Lameed (2011) in his study, showed that the majority of wetland birds observed were Resident, Migratory and Palearctic species. The finding of this study could be established that the most dominant bird species in Nguru Lake were *Philomachus pugnax* (1,330), *Actophilornis africanus* (904), *Phalacrocorax africanus* (719), *Columba guinea* (79).

The findings of the study on diversity of avifauna species in Nguru Wetland



show that the Shannon Diversity Index for the area was 2.55 in Dry Season, while that of Wet Season the area has 3.40. The results are indicative of relative high diversity of bird species in the Wet Season. This finding strongly agrees with the statement of Bibi and Ali (2013) who clearly stated that the values of Shannon-Weiner Diversity Index usually falls between 1.5 and 3.5, only rarely it surpasses 4.5. This relative equal and high diversity of bird species at Nguru Lake is an indication of quantitative measure that reflects how many different species are in existence in the sites. This result supports Mengesha and Bekele (2008) who reported that the avian diversity is an indication of habitat heterogeneity and the number of species and individuals in an area implies the importance of the area. Each habitat has a specific set of microenvironment that is suitable for the species.

CONCLUSION

This study was conducted to investigate the status and diversity of Avifauna species in the study area, the information on the occurrence of bird every month obtained was used for seasonality analysis. However, species identified in dry season was higher compared to the wet season, due to the presence of migrants and Palearctic migrant avifauna species in the study area (dry season) in which this study coincided with the period of their migration in the area. Species diversity attained concentrated in the area, it could be due to the availability, abundance of food in the area.

RECOMMENDATIONS

In view of the findings of this study, the following recommendations are made:

- i. Effective monitoring and strategies for conservation to restore the declining population of bird species.
- ii. Provide public enlightenment of the people around the wetland on values of Nguru Wetlands.

ACKNOWLEDGEMENTS

This research work was sponsored by Tertiary Education Trust Fund (TETFund) as part of the Institution-Based Research (IBR) Intervention.

REFERENCES

Ayeni, A.O., Ogunsesan, A.A. and Adekola, O.A. (2019). Provisioning Ecosystem Services



Provided by the Hadejia-Nguru Wetlands, Nigeria. Current Status and Future Priorities. *Journal of Science Africa* 5 Pp. 2276-2468. <http://doi.org/10.1016/j.sciaf2019.e00124>.

Bibi, F., & Ali, Z. (2013) Measurement of diversity indices of avian communities at Taunsa Barrage Wildlife Sanctuary, Pakistan. *The Journal of Animal & Plant Sciences*, 23(2), 469–474.

Bibby, C. J., Burgess, N. D. and Hill, D. (1992). *Birds Census Techniques*, 2nd Edition. London, Academic Press 24-41.

Borrow N. and Demey R. (2014) *Field Guide to the Birds of Western Africa*. Princeton University Press, United State.

Bourn, D. (2003). Review of Remote Sensing Options for Environmental Monitoring of the Hadejia-Nguru Wetlands, Nigeria. Environmental Research Group Oxford Limited, Oxford, United Kingdom.

Burgess, N. D., Doggart, N. and Lovett J. (2002) The Uluguru Mountains of Eastern Tanzania; the Effect of Forest Loss on Biodiversity. *Oryx* 36: 140-152.

Caro, R., De-frutos, H., and Kltwan, A. (2011). *Typha Charcoal in Senegal: Changing a National Threat into Durable Wealth*. Massachusetts Institute of Technology, Technical Report.

Doggart, N., Lovett J., Mhoro B., Kiure, J. and Burgess, N. (2005). Biodiversity surveys in the Forest Reserves of the Uluguru Mountains. Technical paper for The Wildlife Conservation Society of Tanzania and Tanzanian Forest Conservation Group. DSM, Tanzania. 8–18 pp.

Frontier-Tanzania (2005). *Uluguru Component Biodiversity Survey 2005 (Volume III): Uluguru North Forest Reserve*. Brace bridge, Fanning, Howell, Rubio, St. John (eds). Society for Environmental Exploration and the University of Dares Salaam; CARE-Tanzania, Conservation and Management of the Eastern Arc Mountain Forests (CMEAMF): Uluguru Component, Forestry and Beekeeping Division of the Ministry of Natural Resources and Tourism, Dar Es-salaam, Tanzania. Pp. 1-78

Hadley, S. J. K., Hadley, A. S. and Betts, M. (2012). Acoustic classification of multiple simultaneous bird species: A multi-Instance multi-label approach. *Journal of Acoustical Society of America* 131(6): 4640-4650.

Labe, L. T., Iwar, I. I., Uloko, I. J. (2018). Species diversity and abundance of avifauna in the University of Agriculture, Benue state, north central Nigeria. *Forest Res Eng. Int. J.* 2018; 2(4): 198-202. DOI: 10.15406/freij.2018.02.00048

- Lameed, G.A. (2011). Species diversity and abundance of wild birds in Dagona Waterfowl Sanctuary, Borno State, Nigeria. *African Journal of Environmental Science and Technology*. Vol. 5(10)855-866.
- Mengesha, G., & Bekele, A. (2008). Diversity and relative abundance of birds of Alatish National Park, North Gondar, Ethiopia. *International Journal of Ecology and Environmental Sciences*, 34(2), 215–222.
- Osunsina, I.O.O., Ogialekhe P., Adebayo O.A. and Yisau, M.A. (2018). Bird Species Diversity and Abundance in Borgu Sector of Kainji Lake National Park, Nigeria. *Nigerian Journal of Wildlife Management*. 2(2):41-50
- Ralph, C.J., Geupel, G.R., Pyle, P., Maetin, T.E. and Desante, D.F. (1993). *Handbook of Field Methods for Monitoring Landbirds*. United States Department of Agriculture for Sero. Gen. Tech. Rep. PSW-GTR144.
- Ramchandra, A.M. (2013). Diversity and richness of bird species in newly formed habitats of Chandoli National Park in Western Ghats, Maharashtra State, India. *Biodiversity Journal* 4(1):235-242.
- Ramsar Convention Bureau, (2000). *Background paper on Wetland values and Function*. Gland, Switzerland: Ramsar Convention Bureau, <http://www.ramsar.org/info/values>.
- Sabo B.B. (2016) Checklist of Bird Species at the Hadejia-Nguru Wetlands, Nigeria. *Global Advanced Research Journal of Agricultural Science* (ISSN:2315-5094) Vol. 5(11)Pg.402-412.
- Tworek, S. (2002). Different bird strategies and their responses to habitat changes in an agricultural landscape. *Ecological Research* 17:339-359.