

Studies on effect of planting dates on seed quality of onion (*Allium cepa* L.) in North eastern dry zone of Karnataka

Abstract

Aim: To study the effect of planting dates on seed quality of onion varieties

Study design: Split plot design

Place and Duration of study: Department of Seed Science and Technology, University of Agricultural Sciences, Raichur during 2023-24.

Methodology: Two varieties with five planting dates

Results: The results revealed that significantly highest seed germination (89.3 %), shoot length (8.97 cm), root length (7.02 cm) and seedling vigour index I and II (1428 and 1294, respectively) in D₁ (1st fortnight of November). While, lowest seed germination (84.7 %), shoot length (7.65 cm), root length (6.04 cm), and seedling vigour index I and II (1250 and 1047, respectively) in D₅ (1st fortnight of January). Between varieties, Bhima super (V₂) registered significantly highest seed germination (87.8 cm), shoot length (8.49 cm), root length (6.18 cm), and seedling vigour index I and II (1291 and 1116, respectively). Whereas, lowest seed germination (86.5 cm), shoot length (8.07 cm), root length (5.89 cm), and seedling vigour index I and II (1209 and 978, respectively) recorded in Bellary red (V₁).

Conclusion: The results of present investigation revealed that, bulbs planted during 1st fortnight of November and Bhima super were found to be superior for above study.

Key words: Planting dates, onion varieties, seed germination and seed moisture.

INTRODUCTION

Onion (*Allium cepa* L.) belonging to family Alliaceae. It is the second important vegetable crop of the world after tomato and one of the major vegetable crops cultivated in India. It is an indispensable item in every kitchen as vegetable and condiment, used to flavour many of the food stuffs. It is also used as salad and pickles. It is known fact that seed is a basic and crucial input in agriculture, but it is the quality of the seed that decides the commercial success of a crop or variety. Obviously, the bumper harvest could be possible only when the planting seed possess high quality standards viz., genetic purity, germination, uniformity in weight and size apart from freedom from insect pest and diseases. These quality traits are known to be influenced largely by interaction of environment, cultural practices, harvest and post-harvest management practices at both field and storage levels.

Quality seed is very much essential for enhancing the productivity of a crop plant. However, the availability of quality seeds of desirable variety is seems to be the major constrain for crop production in developing countries like India. The quality of the seed is highest when it completes structural and functional development on plant itself. Thereafter, the quality deteriorates irreparably and irreversibly at varying rates (Delouche *et al.*, 1973) ^[5]. As most of the vegetable crops are propagated sexually through seeds, thus the productivity of vegetable crops depends primarily not only on the genetic constitution of the variety but also the environmental conditions of a particular region. Both genetical as well as agronomical principles of seed production should be adopted for enhancing the crop productivity successfully.

Onion seeds being very short lived in nature, thus the percent and rate of germination of onion seeds also vary considerably among the seed lots leading to difficulties in establishing optimum plant population in the field condition. Onion seeds loose vigour and viability comparatively faster than other vegetables crop seeds and failed to remain viable for more than one year under ambient condition (Justice and Bass, 1979) ^[7]. Initial seed moisture, storage temperature and relative humidity have been found to affect the seed quality. The time of production and interaction of genotypes also determine the quality of seed as the crop produced at different times being exposed to varying environmental conditions especially during seed maturation stage.

MATERIAL AND METHODS

The field experiment was conducted with five planting dates (D₁-1st fortnight of November, D₂- 2nd fortnight of November, D₃-1st fortnight of December, D₄-2nd fortnight of December and D₅-1st fortnight of January) by using two varieties *viz.*, V₁-Bellary Red and V₂-Bhima Super during Rabi 2023-24 at Agricultural Research Station, Hagari, Ballari. The crop was harvested when it reached physiological maturity and seed from all the treatments under this investigation was collected separately. The seed obtained from field experiment was used to assess the seed quality parameters. The laboratory experiment was conducted at Department of Seed Science and Technology, University of Agricultural Sciences, Raichur using FCRD with four replications to know the effect of planting dates and varieties on seed quality.

Observations

The observations were recorded for seed moisture (%), seed germination (%), shoot and root length (cm), seedling dry weight (mg), seedling vigour index-I and seedling vigour index-II. Germination test was conducted as per ISTA using between paper method, the germinated seedlings were evaluated on 6th and 12th days after incubation. Seedling vigour index-I and seedling vigour index-II was calculated as per the formula given by Abdul-Baki and Anderson (1973) ^[1] and expressed in whole number.

$$\text{SVI-I} = \text{Germination (\%)} \times \text{Seedling length (cm)}$$

$$\text{SVI-II} = \text{Germination (\%)} \times \text{Seedling dry weight (mg)}$$

The data pertaining to several characteristics of seed quality were statistically analysed using the methodology outlined by Panse and Sukhatme (1978) ^[12].

RESULTS AND DISCUSSION

The results showed that varieties and planting dates had shown significance difference for germination (%), shoot and root length, seedling vigour index-I and Seedling vigour index-II. But, for seed moisture and seedling dry weight varieties had shown a non significance difference. Interaction effect of varieties and planting dates found to be non significant.

The variety Bhima Super (V₂) recorded maximum germination (87.8 %) over Bellary Red (V₁) (86.5 %). Among different planting dates, the crop planted on 1st fortnight of November (D₁) registered significantly maximum germination (89.3 %). Whereas, crop planted on 1st fortnight of January (D₅) recorded less germination (84.9 %) (Table 1). The reason for increasing the percentage of seed germination in early planting may be due to the highest seed size, seed weight and their food reserves which enhances the germinated of the seeds. These results are similar to the findings of Tesfaye *et al.* (2018) ^[16] and Jagtap *et al.* (2014) ^[6] in onion. Additionally, Saini *et al.* (1980) ^[14] and Kanwar *et al.* (2000) ^[8] reported that onion seeds produced from late planting adversely affected seed quality.

Significantly higher shoot and root length (8.49 and 6.18 cm, respectively) were found in Bhima Super (V₂) over Bellary Red(V₁) (8.07 and 5.89 cm, respectively). Among different planting dates, resultant seeds of 1st fortnight of November (D₁) recorded higher shoot and root length (8.97 and 7.02 cm, respectively) and lower shoot and root length (7.65

and 5.54 cm, respectively) was found in crop which was planted on 1st fortnight of January (D₅). As the planting delayed, the plants had undergone into severe environmental stresses leading to shortening of the growth cycle of the plants. This has led to partial seed filling where the 1000 seed weight reduced significantly over the planting dates. This might have been led to the reduced shoot and root length of the plants as the planting was delayed. This result is conformity with the findings of Das *et al.* (2021), Lamani and Deshpande (2016)^[10] and kumar *et al.* (2015)^[9] in onion.

Seed moisture and seedling dry weight had shown a non significance difference between varieties. Whereas, planting dates had shown a significance difference. Among different planting dates, the resultant seeds of 1st fortnight of November (D₁) registered highest Seed moisture (8.52 %) and Seedling dry weight (14.48 mg). While, the seeds from the crop which was planted on 1st fortnight of December (D₃) registered lowest seed moisture (7.66 %) due to high temperature that led to loss of seed moisture content at harvest. As the plants were under the heat stress it might have made plants to limit its water uptake during its growth stage hence the sink might have received less moisture content. The similar findings were recorded by Arvind *et al.* (2017)^[12] and Rade *et al.* (2012)^[13] in alfalfa. While, the lowest seedling dry weight (9.53 mg) was registered in the resultant seeds of 1st fortnight of January (D₅). This may be due to accumulation of more photosynthates or food reserve in the seeds of 1st fortnight of November (D₁), ultimately leading to higher seedling dry weight. 1st fortnight of January (D₅) has recorded lowest seedling dry weight may be because the seeds did not accumulate as much food reserves (Singh *et al.*, 2017)^[15].

Table 1. Seed quality parameters of onion as influenced by varieties and planting dates

Treatment	Seed germination (%)	Seed moisture (%)	Shoot length (cm)	Root length (cm)	Seedling dry weight (mg)	Seedling vigour index-I	Seedling vigour index-II
Varieties							
V ₁ - Bellary Red	86.5	8.20	8.07	5.89	11.75	1209	978
V ₂ - Bhima Super	87.8	7.99	8.49	6.18	12.15	1291	1116

S.Em. \pm	0.3	0.09	0.09	0.06	0.12	14	12
CD @ 1%	1.0	NS	0.34	0.24	Seedling dry weight (mg)	54	47
Treatment	Seed germination (%)	Seed moisture (%)	Shoot length (cm)	Root length (cm)	Seedling dry weight (mg)	Seedling vigour index-I	Seedling vigour index-II
D ₁ -1 st fortnight of November	89.3	8.52	8.97	7.02	14.48	1428	1294
Interactions (V x D)							
D ₂ -2 nd fortnight of November	88.1	8.13	8.52	6.23	13.67	1301	1209
V ₁ D ₁	88.4	8.79	8.67	6.65	14.40	1354	1213
D ₃ -1 st fortnight of December	87.5	8.17	8.31	5.78	13.72	1235	1130
V ₁ D ₂	87.5	8.17	8.31	5.78	13.72	1235	1130
D ₄ -2 nd fortnight of December	86.0	8.10	7.94	5.63	10.38	1167	893
D ₅ -1 st fortnight of January	84.9	8.06	7.65	5.54	9.53	1121	810
S.Em. \pm	0.6	0.15	0.14	0.24	0.19	21	18
CD @ 1%	1.6	0.41	0.54	0.38	0.76	86	74

Table 2. Seed quality parameters of onion as influenced by interaction between varieties

V ₁ D ₃	86.4	7.80	8.10	5.76	11.30	1199	940
V ₁ D ₄	85.9	8.16	7.80	5.61	10.25	1151	850
V ₁ D ₅	84.7	8.12	7.48	5.48	9.45	1099	776
V ₂ D ₁	90.2	8.25	9.26	7.38	14.55	1501	1376
V ₂ D ₂	89.3	8.11	8.73	6.48	13.98	1358	1307
V ₂ D ₃	88.6	7.52	8.53	5.80	12.13	1271	1119
V ₂ D ₄	86.1	8.05	8.09	5.65	10.50	1183	936
V ₂ D ₅	85.1	8.00	7.82	5.60	9.60	1142	843
S.Em. \pm	0.8	0.21	0.19	0.13	0.27	30	26
CD @ 1%	NS	NS	NS	NS	NS	NS	NS

Legend:

NS: Non-significant

Planting dates

- D₁:** 1st fortnight of November
D₂: 2nd fortnight of November
D₃: 1st fortnight of December
D₄: 2nd fortnight of December
D₅: 1st fortnight of January

Varieties

- V₁:** Bellary Red
V₂: Bhima Super




Fig. 1. Effect of planting dates on germination (%) in onion varieties




Fig. 2. Effect of planting dates on Seedling vigour indices in onion varieties

Legend:

Planting dates

D₁ : 1st fortnight of November
D₂ : 2nd fortnight of November
D₃ : 1st fortnight of December
D₄ : 2nd fortnight of December
D₅ : 1st fortnight of January

Varieties

V₁ : Bellary Red
V₂ : Bhima Super

Assessment of seedling vigour index-I and seedling vigour index-II revealed wide range of variation for varieties and planting dates (Table 1). In case of varieties, Bhima super (V₂) recorded highest seedling vigour index-I and seedling vigour index-II (1291 and 1116,

respectively) than Bellary Red (V₁) (1291 and 978, respectively). Among planting dates, significantly higher seedling vigour index-I and seedling vigour index-II (1428 and 1294, respectively) was recorded from resultant seeds of 1st fortnight of November (D₁) planting bulbs. While, lowest seedling vigour index-I and seedling vigour index-II (1121 and 810, respectively) recorded from the resultant seeds of 1st fortnight of January (D₅) planted bulbs. This might be because higher germination percentage was obtained in 1st fortnight of November (D₁) and robust seedlings obtained due to proper accumulation of food reserves in the seeds. As the planting gets delayed, seed germination percentage, seedling length and seedling dry weight reduced, this has led to the decrease in seedling vigour index - I and II when date of planting was delayed. This was consistent with Malik *et al.* (1999)^[11], Ashagrie *et al.* (2014)^[3] and Das *et al.* (2021)^[4].

It was found that there was a non-significant difference for the interaction of varieties and planting dates for all the seed quality parameters (Table 2).

CONCLUSION

From the study it can be concluded that Bhima Super was found to superior for all the seed quality parameters than Bellary Red. Among the different date of planting, the seed harvested from 1st fortnight of November sown crop had good seed quality compared to delayed planting *i.e.*, 1st fortnight of January.

REFERENCES

1. Abdul-Baki AA and Anderson JD. Vigor Determination in Soybean Seed by Multiple Criteria. *Crop Sci.*1973; 13: 630-633.
2. Arvind Kumar PR, Channakeshava BC and Siddaraju R. Effect of dates of sowing and cutting time on seed yield and quality of alfalfa (*Medicago sativa* L.) CV. RL-88. *Indian J. Agric. Res.* 2017; 51(5): 493-497.
3. Ashagrie T, Derbew B, Sentayehu A and Yehenew G. Effect of planting time and mother bulb size on onion (*Allium cepa* L.) seed yield and quality at Kobo Woreda, Northern Ethiopia. *Int. J. Agric. Res.* 2014; 9(5): 231-241.

4. Das BC, Tripathy P, Sahu G S, Das AK, Swain SK, Mandal P and Sahoo D. Effect of planting dates and varieties on seed yielding attributes of onion (*Allium cepa* L.) under west central table land zone of Odisha. *The Pharma Inn. J.* 2021; 10(7): 151-154.
5. Delouche JC and Baskin CC. Accelerated aging techniques for predicting the relative storability of seed lots. *Seed Science and Technology.* 1973; 1:427-452.
6. Jagtap KB, Patil SD, Patil MR and Kamble DM. Studies on flowering, yield and quality of onion seed cv. Phule Suvarna as influenced by bulb size and planting dates. *Int. Sci. J.* 1973; 1: 2348-6058.
7. Justice OL and Bass IN. Some practical information for storing and transporting seeds at ambient conditions. *Principles and Practices of Seed Storage.* 1979;174-182. London.
8. Kanwar JS, Gill BS and Bal SS. Response of planting time and density to onion seed yield and quality. *Seed Res.* 2000; 28(2): 212-214.
9. Kumar S, Tomar BS, Jain SK, Singh N, Parsad R and Munshi AD. Effect of planting time and density on plant growth, seed yield and quality attributes in onion (*Allium cepa*) cv. Pusa Riddhi. *Indian J. Agric. Sci.* 2015; 85(12): 1578-1585.
10. Lamani K and Deshpande VK. Effect of dates of planting and application of foliar nutrition on plant growth and seed yield of onion cv. Arka Kalyan. *Int. J. Pure Appl. Biosci.* 2017; 5(5): 121-130.
11. Malik, Y. S., Narendra Singh, N. S. and Nehra, B. K., 1999, Effect of planting time, bulb cut and pinching of bolt treatments on yield and quality of onion seed. *Veg. Sci.*, 26(2): 143-145.
12. Panse VG and Sukhatme PV. *Statistical Methods for Agricultural Workers*, ICAR, New Delhi: 1978;167-174.
13. Rade S, Dragoljub B, Dragan D, Vladeta S, Dragan T, Jasmina M and Dragoslav D. Influence of plant density on yield components, yield and quality of seed and forage yields of alfalfa varieties. *Romanian Agric. Res.* 2012; 29(1): 245-254.
14. Saini SK, Singh JN and Gupta PC. Effect of planting date and variety on seed quality of soybean. *Seed Res.* 1980; 8(2): 141-149.
15. Singh DP, Verma SC, Sharma HK, Mehta DK, Thakur KS and Bharat NK. Effect of different organic inputs and transplanting dates on seed quality parameters of radish (*Raphanus sativus* L.). *J. Appl. Nat. Sci.* 2017; 9(2): 1203-1207.

16. Tesfaye M, Belew D, Dessalegn Y and Shumye G. Effect of planting time on growth, yield components, seed yield and quality of onion (*Allium cepa* L.) at Tehuledere district, northeastern Ethiopia. *Agriculture Food Security*, 2018; 7: 1-8.

UNDER PEER REVIEW