

# THE INFLUENCE OF MICRONUTRIENTS ON GROWTH PERFORMANCE OF GINGER (*Zingiber officinale*) CV. MAHIMA

## Abstract

The present investigation entitled, “The influence of micronutrients on growth performance of ginger (*zingiber officinale*) cv. mahima” was conducted at All India Co-ordinated Research Project on Spices, Asond Block, Central Experiment Station, Wakawali, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. Total nine treatments were observed in Randomized Block Design in three replications. The experimental findings revealed that the treatment, T<sub>1</sub> i.e. RDF + Foliar Spray of ZnSO<sub>4</sub> @ 0.5 % recorded better results in number of tillers per platnt (12.57), plant height (77.65 cm), number of leaves per plant (76.03), leaf length (23.25 cm) and leaf width (3.08 cm). Thus, it was concluded that the treatment T<sub>1</sub> (RDF + Foliar Spray of ZnSO<sub>4</sub> @ 0.5 %) was found to be the best for growth performance of ginger (*Zingiber officinale* rosc.) under Konkan agroclimatic conditions.

**Keywords:** ginger, micronutrient, foliar spray, growth

## 1. Introduction

Ginger (*Zingiber officinale* Rosc.) is one of the oldest known spices valued for its distinctive aroma and pungency. Ginger is a key component in both conventional and modern medicine. It strengthens immunity and is a rich source of minerals and physiologically active compounds. Ginger is utilized in both fresh and dried form. It is mostly used as spice, for candies, pickles and as a medicinal herb for the treatment of gastrointestinal diseases, including dyspepsia, nausea and diarrhea. In India, ginger is cultivated across 1,90,010 hectares, yielding 22,42,780 tons. It is grown in 21 states, with major production concentrated in Madhyapradesh, Karnataka, the

northeastern states, Sikkim, Orissa, West Bengal, Andhra Pradesh and Maharashtra. (Anon., 2024).

Ginger is a long duration and heavy feeder crop and responds well to manuring and is also affected by deficiency of micronutrients. Micronutrients are the essential elements required in smaller amount but importance of these in growth and development of a crop is very much necessary as of major nutrients. Nowadays micronutrient deficiency is a prevalent issue in ginger growing soils and application of micronutrient particularly zinc (Zn) and boron (B) enhance the growth of the crop especially in deficient soils (Parthasarathy *et al.*, 2010). Micronutrients can play an important role in disease control of spice crop like ginger. Thus, there is an urgent need to augment supplies of customized fertilizers supplying the micronutrients sufficiently to support the overall need of nutrient. Application of micronutrients showed positive impact on different crops (Barbosa *et al.*, 2016; Sarker *et al.*, 2018; Alkarawi and Hasan, 2021)

Ginger is a non-traditional crop in Konkan region of Maharashtra. However, the climatic and soil conditions seem to be suitable for its cultivation in this region. The successful introduction of this crop in a large area will not only provide an opportunity to generate income but will also be an option for crop diversification. Very scanty information is available with respect to the effect of zinc, boron, iron and manganese on the growth of ginger. Keeping these points in view, present investigation was undertaken.

## 2. Material and Methods

An experiment was conducted during the year 2023-24, under the climatic condition of Konkan region at All India Co-ordinated Research Project on Spices, Asond Block, Central Experiment Station, Wakavali, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, having hot and humid climate with the moderate temperature ranging from as low as 7.5°C to as high as 38.5°C and average relative humidity ranging from 55 to 100 %. The region receives very high rainfall (above 3000 mm, annually). One month old healthy pro-tray seedlings of Mahima cultivar of ginger were used for planting and are planted at the depth of 5 cm and distance of 30 cm between rows and 25 cm between plants. The experiment was laid out in Randomized Block Design (RBD) with three replications and nine treatments viz. T<sub>1</sub> (RDF + Foliar Spray of ZnSO<sub>4</sub> @ 0.5 %), T<sub>2</sub> (RDF + Foliar Spray of ZnSO<sub>4</sub> @ 0.75 %), T<sub>3</sub> (RDF + Foliar Spray of Borax @ 0.2 %), T<sub>4</sub> (RDF + Foliar Spray of Borax @ 0.3 %), T<sub>5</sub> (RDF + Foliar Spray of MnSO<sub>4</sub> @ 0.25 %), T<sub>6</sub> (RDF + Foliar Spray of MnSO<sub>4</sub> @ 0.5 %), T<sub>7</sub> (RDF + Foliar Spray of FeSO<sub>4</sub> @ 0.3 %), T<sub>8</sub> (RDF + Foliar Spray of FeSO<sub>4</sub> @ 0.4 %), T<sub>9</sub> (RDF (75:50:50 kg NPK ha<sup>-1</sup>). For the foliar application of different doses of zinc (0.5% and 0.75%), boron (0.2% and 0.3%), manganese (0.25% and 0.50%) and iron (0.3% and 0.4%) solutions were prepared by using zinc sulphate, borax, manganese sulphate and ferrous sulphate fertilizer by dissolving the respective weight of chemical fertilizers in 1L of water by continuous stirring. The spraying was carried out early in the morning by using knapsack hand sprayer. The spraying was done at 60, 90 and 120 days after transplanting.

### 3. Result and Discussion

#### 3.1 Number of tillers per plant

According to the results of present investigation, the highest number of tillers per plant at 90, 120, 150 and 180 DAP were observed in treatment T<sub>1</sub> (7.77, 10.70, 12.07 and 12.57, respectively). While the minimum number of tillers per plant were observed in treatment T<sub>9</sub> (5.40, 7.10, 7.53 and 7.63, respectively). The increase in number of tillers per plant by zinc treated plots might be due to better synthesis of tryptophan being the precursor for auxin synthesis and also it might be due to zinc is effective in regulating plant growth because it forms a part of enzymes system (e.g. carbonic anhydrase), which regulates plant growth. Besides, zinc stimulates photosynthetic activity and its presence is important for protein synthesis (Singh and Dwivedi 2007). Roy *et al.* (1992) also reported increased growth of ginger with application of micronutrients such as zinc, boron and iron through foliar spray.

**Table 1. Effect of different micronutrients levels on number of tillers per plant of ginger (cv. Mahima)**

Tr.	Treatment detail	Number of tillers per plant				
		60 DAP	90 DAP	120 DAP	150 DAP	180 DAP
T <sub>1</sub>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.5 %	1.63	7.77	10.70	12.07	12.57
T <sub>2</sub>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.75 %	1.63	5.79	7.60	9.10	9.27
T <sub>3</sub>	RDF + Foliar Spray of Borax @ 0.2 %	1.90	5.74	7.55	9.00	9.13
T <sub>4</sub>	RDF + Foliar Spray of Borax @ 0.3 %	1.87	6.70	8.87	10.43	10.83
T <sub>5</sub>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.25 %	1.80	5.87	7.86	9.20	9.30
T <sub>6</sub>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.5 %	1.73	5.68	7.39	8.73	9.12
T <sub>7</sub>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.3 %	2.20	5.67	7.36	8.53	8.53
T <sub>8</sub>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.4 %	1.93	5.63	7.27	8.37	8.50
T <sub>9</sub>	RDF (75:50:50 kg NPK ha <sup>-1</sup> )	1.53	5.40	7.10	7.53	7.63
	<b>Range</b>	<b>1.53-2.20</b>	<b>5.40-7.77</b>	<b>7.10 - 10.70</b>	<b>7.53 - 12.07</b>	<b>7.63 - 12.57</b>
	<b>Mean</b>	<b>1.80</b>	<b>6.04</b>	<b>7.97</b>	<b>9.22</b>	<b>9.43</b>
	<b>S.Em (±)</b>	<b>0.28</b>	<b>0.35</b>	<b>0.43</b>	<b>0.36</b>	<b>0.43</b>
	<b>C.D. @ 5%</b>	<b>0.84</b>	<b>1.04</b>	<b>1.28</b>	<b>1.09</b>	<b>1.28</b>
	<b>‘F’ test</b>	<b>NS</b>	<b>SIG</b>	<b>SIG</b>	<b>SIG</b>	<b>SIG</b>

#### 3.2 Height of plant (cm)

Among the different treatments, the maximum plant height at 90, 120, 150 and 180 DAP were noted in treatment T<sub>1</sub> (39.46 cm, 52.89 cm, 71.00 cm and 77.65 cm, respectively). While minimum plant height was observed in treatment T<sub>9</sub> (33.59 cm, 48.18 cm, 66.81 cm and 70.51 cm,

respectively). Zinc is effective in regulation of plant growth. Foliar nutrition usually penetrate the cuticle of the leaf or stomata, enter the cells rapidly and fulfill the nutrient demand of the growing plant and thus ameliorate nutrient deficiency (Devi and Shanti 2013). Zinc is the structural constituent of few enzymes including RNA polymerase (Marschner, 2003) and carbonic anhydrase (Hafeez *et al.*, 2013) besides activating a large number of enzymes influencing metabolisms of carbohydrate, protein (Hafeez *et al.*, 2013) and auxins (Marschner, 2003). Similar increase in plant height of ginger with the application of different doses of zinc, boron and iron was also reported by Shadap *et al.* (2018).

**Table 2. Effect of different micronutrients levels on height of plant of ginger (cv. Mahima)**

Tr.	Treatment detail	Plant height (cm)				
		60 DAP	90 DAP	120 DAP	150 DAP	180 DAP
T <sub>1</sub>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.5 %	22.87	39.46	52.89	71.00	77.65
T <sub>2</sub>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.75 %	22.24	34.73	49.69	68.18	73.04
T <sub>3</sub>	RDF + Foliar Spray of Borax @ 0.2 %	22.94	34.89	48.96	67.71	72.09
T <sub>4</sub>	RDF + Foliar Spray of Borax @ 0.3 %	22.60	38.80	51.37	69.86	75.84
T <sub>5</sub>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.25 %	23.14	35.67	49.82	68.31	74.37
T <sub>6</sub>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.5 %	22.38	34.41	48.47	67.55	71.46
T <sub>7</sub>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.3 %	22.62	34.24	48.42	67.41	71.38
T <sub>8</sub>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.4 %	22.63	34.42	48.87	67.67	71.81
T <sub>9</sub>	RDF (75:50:50 kg NPK ha <sup>-1</sup> )	23.15	33.59	48.18	66.81	70.51
	<b>Range</b>	<b>22.24 - 23.15</b>	<b>33.59 - 39.46</b>	<b>48.18- 52.89</b>	<b>66.81 - 71.00</b>	<b>70.51 - 77.65</b>
	<b>Mean</b>	<b>22.73</b>	<b>35.58</b>	<b>49.63</b>	<b>68.26</b>	<b>73.13</b>
	<b>S.Em (±)</b>	<b>0.55</b>	<b>0.70</b>	<b>0.62</b>	<b>0.38</b>	<b>1.06</b>
	<b>C.D. @ 5%</b>	<b>1.66</b>	<b>2.11</b>	<b>1.85</b>	<b>1.15</b>	<b>3.18</b>
	<b>‘F’ test</b>	<b>NS</b>	<b>SIG</b>	<b>SIG</b>	<b>SIG</b>	<b>SIG</b>

### 3.3 Number of leaves per plant

The maximum number of leaves per plant (76.03) was noticed in treatment T<sub>1</sub> [RDF + Foliar Spray of ZnSO<sub>4</sub> @ 0.5 %] whereas, the minimum number of leaves per plant (61.57) was observed in treatment T<sub>9</sub> [RDF (75:50:50 kg NPK ha<sup>-1</sup>)]. Application of different micronutrient especially zinc (Zn) improves growth, yield and quality of the ginger crop (Parthasarathy *et al.*, 2010). Zinc is critical for chlorophyll synthesis, which enhances photosynthesis. Improved photosynthesis contributes to better growth and yield (Roy *et al.*, 1992).

**Table 3. Effect of different micronutrients levels on number of leaves per plant of ginger (cv. Mahima)**

Tr.	Treatment detail	Number of leaves per plant				
		60 DAP	90 DAP	120 DAP	150 DAP	180 DAP
T <sub>1</sub>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.5 %	8.87	33.67	64.63	72.57	76.03
T <sub>2</sub>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.75 %	7.93	27.50	57.57	61.67	63.80
T <sub>3</sub>	RDF + Foliar Spray of Borax @ 0.2 %	7.27	26.40	56.87	61.53	63.00
T <sub>4</sub>	RDF + Foliar Spray of Borax @ 0.3 %	8.87	33.07	61.10	70.00	72.73
T <sub>5</sub>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.25 %	8.93	27.57	58.13	62.80	64.37
T <sub>6</sub>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.5 %	7.53	26.67	56.97	61.63	63.17
T <sub>7</sub>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.3 %	7.90	26.50	56.37	61.37	62.77
T <sub>8</sub>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.4 %	8.20	27.27	55.43	61.17	62.37
T <sub>9</sub>	RDF (75:50:50 kg NPK ha <sup>-1</sup> )	7.57	26.03	53.50	60.07	61.57
	Range	7.27 - 8.93	26.03 - 33.67	53.50 - 64.63	60.07 - 72.57	61.57 - 76.03
	Mean	8.12	28.30	57.84	63.64	65.53
	S.Em (±)	0.47	0.74	1.29	0.79	0.84
	C.D. @ 5%	1.42	2.21	3.87	2.36	2.52
	'F' test	NS	SIG	SIG	SIG	SIG

### 3.4 Leaf length (cm)

The data showed that, highest leaf length at 90, 120, 150 & 180 DAP were observed in treatment T<sub>1</sub> (14.62 cm, 20.02 cm, 21.70 cm and 23.25 cm, respectively). The minimum leaf length was noticed in treatment T<sub>9</sub> (26.03, 53.50, 60.07 and 61.57, respectively). Improved photosynthesis efficiency, facilitated by optimal micronutrient levels, can contribute to increased leaf growth. Higher levels of chlorophyll, which are supported by micronutrients, enhance the plant's ability to capture light and convert it into energy, promoting longer leaf development. Zinc stimulates photosynthetic activity. Boron plays an important role in cell division as well as nitrogen metabolism, carbohydrate metabolism and water relation in plants. The results in this experiment are also in agreement with the findings of (Halder *et al.*, 2007a) who reported increased growth of ginger plants with the application of micronutrients.

**Table 4. Effect of different micronutrients levels on leaf length of ginger (cv. Mahima)**

Tr.	Treatment detail	Leaf length (cm)
-----	------------------	------------------

		<b>60 DAP</b>	<b>90 DAP</b>	<b>120 DAP</b>	<b>150 DAP</b>	<b>180 DAP</b>
<b>T<sub>1</sub></b>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.5 %	10.11	14.62	20.02	21.70	23.25
<b>T<sub>2</sub></b>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.75 %	10.39	12.11	18.74	20.20	21.21
<b>T<sub>3</sub></b>	RDF + Foliar Spray of Borax @ 0.2 %	9.95	11.97	18.71	20.09	21.12
<b>T<sub>4</sub></b>	RDF + Foliar Spray of Borax @ 0.3 %	10.33	14.15	19.95	21.47	23.03
<b>T<sub>5</sub></b>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.25 %	9.99	12.12	18.82	20.45	21.66
<b>T<sub>6</sub></b>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.5 %	10.14	11.82	18.67	19.75	21.01
<b>T<sub>7</sub></b>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.3 %	10.18	11.61	17.91	19.35	20.20
<b>T<sub>8</sub></b>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.4 %	10.22	11.78	18.15	19.72	20.90
<b>T<sub>9</sub></b>	RDF (75:50:50 kg NPK ha <sup>-1</sup> )	10.18	11.59	17.88	19.15	19.26
	<b>Range</b>	<b>9.95 - 10.39</b>	<b>11.59 - 14.62</b>	<b>17.88 - 20.02</b>	<b>19.1 - 21.70</b>	<b>19.26 - 23.25</b>
	<b>Mean</b>	<b>10.17</b>	<b>12.42</b>	<b>18.76</b>	<b>20.21</b>	<b>21.29</b>
	<b>S.Em (±)</b>	<b>0.20</b>	<b>0.31</b>	<b>0.38</b>	<b>0.35</b>	<b>0.42</b>
	<b>C.D. @ 5%</b>	<b>0.60</b>	<b>0.92</b>	<b>1.12</b>	<b>1.06</b>	<b>1.27</b>
	<b>'F' test</b>	<b>NS</b>	<b>SIG</b>	<b>SIG</b>	<b>SIG</b>	<b>SIG</b>

### 3.5 Leaf width (cm)

The highest leaf width at 120, 150 & 180 DAS were observed in treatment T<sub>1</sub> (2.85 cm, 2.97 cm and 3.08 cm, respectively). The minimum leaf width was noticed in treatment T<sub>9</sub> (2.41 cm, 2.46 cm and 2.56 cm, respectively). Zinc and iron are very effective in regulating plant growth because it forms a part of enzymes system (e.g. carbonic anhydrase), which regulates plant growth. Besides, zinc stimulates photosynthetic activity and its presence is important for protein synthesis (Singh and Dwivedi 2007). Similar increase in leaf width of ginger with the application of different doses of zinc, boron and iron was also reported by Shadap *et al.* (2018).

**Table 5. Effect of different micronutrients levels on leaf width of ginger (cv. Mahima)**

<b>Tr.</b>	<b>Treatment detail</b>	<b>Leaf width (cm)</b>				
		<b>60 DAP</b>	<b>90 DAP</b>	<b>120 DAP</b>	<b>150 DAP</b>	<b>180 DAP</b>
<b>T<sub>1</sub></b>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.5 %	1.53	2.46	2.85	2.97	3.08
<b>T<sub>2</sub></b>	RDF + Foliar Spray of ZnSO <sub>4</sub> @ 0.75 %	1.46	2.35	2.65	2.67	2.85
<b>T<sub>3</sub></b>	RDF + Foliar Spray of Borax @ 0.2 %	1.36	2.41	2.60	2.65	2.83
<b>T<sub>4</sub></b>	RDF + Foliar Spray of Borax @ 0.3 %	1.58	2.53	2.75	2.87	3.03
<b>T<sub>5</sub></b>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.25 %	1.49	2.39	2.67	2.68	2.85
<b>T<sub>6</sub></b>	RDF + Foliar Spray of MnSO <sub>4</sub> @ 0.5 %	1.51	2.45	2.56	2.60	2.80

<b>T<sub>7</sub></b>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.3 %	1.39	2.33	2.44	2.50	2.67
<b>T<sub>8</sub></b>	RDF + Foliar Spray of FeSO <sub>4</sub> @ 0.4 %	1.39	2.46	2.57	2.64	2.80
<b>T<sub>9</sub></b>	RDF (75:50:50 kg NPK ha <sup>-1</sup> )	1.49	2.31	2.41	2.46	2.56
	<b>Range</b>	<b>1.36 - 1.58</b>	<b>2.31 - 2.53</b>	<b>2.41 - 2.85</b>	<b>2.46 - 2.97</b>	<b>2.56 - 3.08</b>
	<b>Mean</b>	<b>1.47</b>	<b>2.41</b>	<b>2.61</b>	<b>2.67</b>	<b>2.83</b>
	<b>S.Em (±)</b>	<b>0.06</b>	<b>0.05</b>	<b>0.05</b>	<b>0.07</b>	<b>0.05</b>
	<b>C.D. @ 5%</b>	<b>0.18</b>	<b>0.16</b>	<b>0.16</b>	<b>0.20</b>	<b>0.16</b>
	<b>‘F’ test</b>	<b>NS</b>	<b>NS</b>	<b>SIG</b>	<b>SIG</b>	<b>SIG</b>

#### 4. Conclusion

On the basis of results found from the present experiment, it can be concluded that foliar application of different micronutrient was found beneficial for growth performance of ginger crop. Treatment T<sub>1</sub> - RDF + Foliar Spray of ZnSO<sub>4</sub> @ 0.5 % was significantly superior in terms of number of tillers per plant (12.57), plant height (77.65 cm), number of leaves per plant (76.03), leaf length (23.25 cm) and leaf width (3.08 cm).

#### Disclaimer (Artificial Intelligence)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOYT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### References

- Alkarawi, H. H. and Hasan, S. Q. (2021). Influence of foliar application of nano-zinc and iron fertilizers on growth and yield of bell pepper (*Capsicum annuum* L.). *Indian J. Ecology*, **48**(4): 1115-1119.
- Anonymous, (2024). Spices Board, India & Ministry of Agriculture and Farmers Welfare, Govt. of India. <https://www.indianspices.com>.
- Barbosa J. M., Rezende, C. F. A., Leandro, W.M., Ratke R. F., Flores, R. A. and Silva, A. R. (2016). Effects of micronutrients application on soybean yield. *Australian J. Crop Science*, **10**(8): 1092-1097.
- Hafeez, B., Khanil, Y. M and Saleem, M. (2013). Role of zinc in plant nutrition-A review, *American J. Expt. Agri.*, **3**: 374-91.

- Halder, N. K., Shill, N. C., Siddiky, M. A., Gomes, R. and Sarkar, J. (2007a). Response of ginger to zinc and boron fertilization. *Asian J. Pl Sci.*, **6**: 394-398.
- Halder, N. K., Shill, N. C., Siddiky, M. A., Sarkar, J. and Gomes, R. (2007b), Response of turmeric to zinc and boron fertilization, *J. Biol. Sci.*, **7**(1): 182-187.
- Marschner, H. (2003). Mineral Nutrition in Higher Plants. Academic Press, London, pp. 889.
- Parthasarathy, V. A., Dinesh, R., Srinivasan, V. and Hamza, S. (2010). Integrated Nutrient Management in Major Spices. *Indian J. Fert.*, **6**: 110–128.
- Roy, A., Chatterjee, R., Hassan, A. and Mitra, S.K. (1992). Effect of Zn, Fe and B on growth, yield and nutrient content in leaf of ginger. *Indian Cocoa, Arecanut Spices J.*, **15**: 99-101.
- Sarker MMH, Moslehuddin AZM, Jahiruddin M. and Islam, M. R. (2018). Effects of micronutrient application on different attributes of potato in floodplain soils of Bangladesh. *SAARC J. Agriculture*, **16**(2): 97-108.
- Shadap, A., Pariari, A. and Lyngdoh, Y. A. (2018). Effect of foliar sprays of micronutrients on the performance of ginger (*Zingiber officinale* Rosc.). *Ann. Biol.* **34**(2): 212-214.
- Singh, S. P. and Dwivedi, D. K. (2007). Impact of zinc, boron and iron elements on yield and economics of ginger (*Zingiber-officinale* Ros.). *Int. J. agric. Sci.*, **3**(1): 136-138.