

Original Research Article

Effect of Biofertilizer and Phosphorus on Growth and Yield of Chickpea (*Cicerarietinum*)

ABSTRACT

A field experiment was conducted at Crop Research Farm (CRF), Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj (UP) during Rabi 2021 to study the “Effect of ~~Biofertilizer~~ ~~biofertilizer~~ and ~~Phosphorus~~ ~~phosphorus~~ on ~~Growth~~ ~~growth~~ and ~~Yield~~ ~~yield~~ of ~~Chickpea~~ ~~chickpea~~ (*Cicerarietinum*)”. It was consisting of three combinations of ~~Biofertilizer~~ ~~biofertilizer~~ and ~~Phosphorus~~ ~~phosphorus~~. The experiment was laid out in Randomized Block Design with ten treatments each replicated thrice. The experiment results were revealed that the growth and yield parameters such as plant height (38.52 cm), number of nodules per plant (36.28), plant dry weight (9.03 g/plant), number of pods per plant (34.29), number of seeds per pod (1.89), test weight (245.76 g) at harvest significantly recorded in T₉ with the application of Rhizobium+PSB + ~~Phosphorus~~ ~~phosphorus~~ 60kg/ha. Moreover, grain yield (2.07 t/ha), stover yield (3.58 ~~t/ha~~ ~~ha~~⁻¹), gross return (108523.7 INR/ha), net return (75364.02 INR/ha) and B:C ratio (2.27) were also recorded higher in T₉ which is Rhizobium+PSB + ~~Phosphorus~~ ~~phosphorus~~ 60kg/ha among all other treatments.

Keywords: Rhizobium, PSB, Phosphorus, Chickpea, Yield

1. INTRODUCTION

Pulses play an important and diverse role in the farming system and in the diets of people around the world. Besides being rich in protein, they sustain the productivity of cropping systems. Pulse crop plays an important role in Indian Agriculture. (Yadav et al 2017) India is the largest producer and consumer of pulses in the world. Pulses contain high percentage of quality protein nearly three times as much as cereals. (Umadevi and Ganeshan 2007). Pulses are known for increasing productivity of soil through fixation of nitrogen from atmosphere, addition of biomass to soil and secretion of growth promoting substances.

Chickpea (*Cicerarietinum*.L) is the most important Rabi pulse crop of India. Among the pulse crops, chickpea occupies an important position due to its nutritious value (17-23%). Area under chickpea cultivation in India is around 8.3mha and in Andhra Pradesh it is 4.7lakh ha. All India production is around 7.1 million tonnes and it is 5 lakh tonnes in Andhra Pradesh. (Annual Report 2016-17, MoAF, GOI). The current average global yield of chickpea is 0.9 t ha⁻¹, which is much

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Superscript

Formatted: Font: Not Italic

Formatted: Font: Italic

lower than its estimated potential of 6 t ha⁻¹ under the optimum cultivated conditions,(FAO, 2012). The low production of chickpea is due to improper use of fertilizers and least importance given to biofertilizers such as Rhizobium and PSB.

Plant nutrients are the main sources for improving quantity and quality of chickpea production. Phosphorus is known to improve crop quality. It is also an important constituent of vital substances like phospholipids and phosphoproteins. Sufficient supply of phosphorus to plant, hastens the maturity and increases the rate of nodulation and pod development. The increase in the nodulation might be due to the enhanced and established good rooting system with the application of Phosphorus. Phosphorus is one of the critical nutrient deficiency in Indian soils and may cause upto 29-45% yield losses in chickpea. (Ahlawat et al 2007). Most of the phosphorus present in the soil is unavailable to plants which are made available through the activities of efficient ~~microorganisms~~ microorganisms like bacteria, fungi and cyanobacteria with production of organic acid and increasing phosphatase enzyme activity.(Rajneesh et al 2018)

Biofertilizers may colonizes the rhizosphere and promotes growth by increasing the availability and supply of nutrients to crop. Nitrogen fixer and phosphate solubilizing microorganisms play an important role in supplementing nitrogen and phosphorus to plants, allowing a sustainable use of nitrogen and phosphate fertilizer (Tambekar et al 2009). Chickpea establishes a symbiotic association with Rhizobium. Rhizobium is one of the nitrogen fixing bacteria which fix atmospheric nitrogen by the symbiotic association with leguminous plants. Rhizobium originally called *Bacillus redicicola* was first isolated by Beijerinck (1901). However, PSB also increase the yield of chickpea by 10-30%. Both Rhizobium and PSB inoculation significantly increased the nodules mainly due to the fact that the nitrogenase enzyme present in the bacteria get introduced through infection causes nodule formation. Application of PSB facilitates the root development vis-à-vis nodule formation and proper development of nodules by increasing the availability of phosphorus through mobilising the unavailable phosphorus present in the soil. The use of bio-fertilizers with reduced quantity of chemical fertilizers is therefore recommended for better yield of chickpea. (Dinesh et al 2015)

2. MATERIALS AND METHODS

The experiment was carried out during *Rabi* 2021-2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh, which is located at 25.28°N latitude, 81.54°E longitude and 98m altitude above the mean sea level. The experiment laid out in Randomized Block Design which consisting of ten treatments with T₁: Rhizobium 20g/kg seed + Phosphorus 20kg/ha, T₂: Rhizobium 20g/kg seed+ Phosphorus 40kg/ha, T₃: Rhizobium 20g/kg seed + Phosphorus 60kg/ha, T₄: PSB 20 g/kg seed + Phosphorus 20kg/ha, T₅: PSB 20g/kg seed + Phosphorus 40kg/ha, T₆: PSB 20g/kg seed + Phosphorus 60kg/ha, T₇: Rhizobium+PSB 20g/kg seed + Phosphorus 20kg/ha, T₈: Rhizobium+PSB 20g/kg seed + Phosphorus 40kg/ha, T₉: Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha, T₁₀: Control in three replications. The observations on different growth parameters are Plant height (cm), ~~Number-number~~ of nodules

per plant, Plant dry weight (g) and yield parameters are ~~Number-number~~ of pods per plant, Number of seeds per pod, ~~Test-test~~ weight (g), Seed yield (kg/ha), Stover yield (kg/ha) and Harvest index (%) were recorded and statistically analysed using analysis of variance (ANOVA) as applicable to Randomized Block Design.

Table 1: Details of Treatment Combinations.

Treatment No.	Treatment Combinations
T ₁	Rhizobium 20g/kg seeds + Phosphorus 20kg/ha
T ₂	Rhizobium 20g/kg seeds + Phosphorus 40kg/ha
T ₃	Rhizobium 20g/kg seeds + Phosphorus 60kg/ha
T ₄	PSB 20g/kg seeds + Phosphorus 20kg/ha
T ₅	PSB 20g/kg seeds + Phosphorus 40kg/ha
T ₆	PSB 20g/kg seeds + Phosphorus 60kg/ha
T ₇	Rhizobium + PSB 20g/kg seeds + Phosphorus 20kg/ha
T ₈	Rhizobium + PSB 20g/kg seeds + Phosphorus 40kg/ha
T ₉	Rhizobium + PSB 20g/kg seeds + Phosphorus 60kg/ha
T ₁₀	Control

3. RESULTS AND DISCUSSION

3.1 Growth ~~Attributes~~attributes

3.1.1 Plant ~~Height-height~~ (cm)

As shown in Table 2, significantly highest plant height (54.94cm) was recorded in T₉ Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha, than all other treatments. However, T₈ Rhizobium+PSB 20g/kg seed + Phosphorus 40kg/ha was statistically at par with T₉ Rhizobium+PSB 20g/kg seed + Phosphorus 60 kg/ha. The minimum plant height was recorded in control plot which is 49.41cm. The plant height of chickpea was significantly influenced by biofertilizers. The increase in growth might be due to enhanced photosynthetic efficiency of Rhizobium+PSB inoculated plant. Inoculations of PSB which are known to produce growth

hormones are likely to favour increased plant height. Phosphorus applications increased plant height and number of branches, Dahiya et al 1993.

3.1.2 Number of ~~Nodules~~nodules per plant

The number of nodules per plant in chickpea were significantly influenced by Rhizobium and PSB inoculation at flowering stage of the crop growth. Maximum number of nodules per plant (36.28) was recorded with the application of Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. However, T₈ Rhizobium+PSB 20g/kg seed + Phosphorus 40kg/ha was statistically at par with T₉ Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. The minimum number of nodules per plant were recorded in control plot which is 28.46. Inoculation of seed with Rhizobium+PSB produce significantly higher number of nodules in comparison to other inoculants. (Akansha 2018). The data (Table 2) indicates that the increase of phosphorus levels has increased the number of nodules per plant with combined application of Rhizobium+PSB.

3.1.3 Plant ~~Dry-dry~~Weight-weight (g)

Treatment with Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha was recorded with significantly maximum plant dry weight (23.71g) than all other treatments. However, T₈ Rhizobium+PSB 20g/kg seed + Phosphorus 40kg/ha was statistically at par with T₉ Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. The minimum plant dry weight (16.15g) was observed in control plot. Application of Phosphorus significantly increased the dry matter accumulation in the plant and differences were seen in 60 kg P/ha. The response of phosphorus in terms of dry matter accumulation up to 60kg P/ha have also been reported Singh et al 2005 in chickpea. Seed inoculation with Rhizobium+PSB also produced significantly higher dry matter of chickpea. Thenua et al (2010) reported that application of phosphorus as single super phosphate (SSP) recorded significantly higher plant height, branches per plant and dry matter accumulation in chickpea.

Table 2: Effect of Biofertilizer and Phosphorus on Growth attributes of Chickpea

S.No	Treatments	Plant Height (cm)	Number of nodules/plant	Plant Dry weight (g)
1	Rhizobium+Phosphorus 20kg/ha	53.25	33.24	20.58
2	Rhizobium+Phosphorus 40kg/ha	52.16	32.87	19.75
3	Rhizobium+Phosphorus 60kg/ha	53.92	35.80	21.82
4	PSB+Phosphorus 20kg/ha	52.59	32.19	19.79
5	PSB+Phosphorus 40kg/ha	51.81	34.47	19.83
6	PSB+Phosphorus 60kg/ha	52.68	31.68	18.45
7	Rhizobium+PSB+Phosphorus 20kg/ha	52.75	33.13	20.67
8	Rhizobium+PSB+Phosphorus 40kg/ha	54.46	35.93	22.33
9	Rhizobium+PSB+Phosphorus 60kg/ha	54.94	36.28	23.71
10	Control	49.41	28.46	16.15
	F test	S	S	S
	SEm(±)	0.48	0.39	0.49
	CD (P=0.05)	1.42	1.15	1.45

3.2. Yield Attributes

As shown in Table 3, yield parameters are summarized statistically. At the time of harvest, significantly effective number of pods per plant (34.29) was recorded with combined application of Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. However, Rhizobium+PSB 20g/kg seed + Phosphorus 40kg/ha was statistically at par with Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. The minimum effective pods per plant (26.65) was recorded in control plot. Significantly maximum number of seeds per pod (1.89) was recorded with application of Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. However, Rhizobium+PSB 20g/kg seed + Phosphorus 40kg/ha was statistically at par with Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. The minimum number of seeds per pod (1.11) was recorded in control plot. Significantly, maximum test weight (245.76g) was recorded in treatment combination Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. However, Rhizobium+PSB 20g/kg seed + Phosphorus 40 kg/ha was statistically at par with Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. The minimum test weight was recorded in control plot which is 194.39g.

Significantly, maximum grain yield (2.07t/ha) was recorded with Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. However, Rhizobium+PSB 20g/kg seed + Phosphorus 40kg/ha was statistically at par with Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. The minimum grain yield (1.65t/ha) was recorded in control plot. Significantly, maximum stover yield (3.58t/ha) was recorded in the treatment combination of Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. However, Rhizobium+PSB 20g/kg seed + Phosphorus 40kg/ha was

statistically at par with Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha. The minimum stover yield (2.50t/ha) was recorded in control plot. Interaction of inoculation of biofertilizer and different levels of phosphorus was also found to be highly significant.

Application of Phosphorus + Rhizobium + PSB recorded higher value of growth as well as yield contributing characters similar result was given by Jarande et al 2006 than all other treatments. Application of phosphorus 60kg/ha significantly increased yield attributes like pods per plant, seeds per pod, test weight, seed yield and stover yield (Arya et al 2002, Meena et al 2002, Pramanik and Singh 2003). Combined effect of Rhizobium+PSB seed inoculation recorded significantly higher yield attributes than Rhizobium or PSB inoculation alone. The beneficial effect of Rhizobium and PSB inoculation was also reported by Singh et al 2011 observed significantly higher seed yield of chickpea with rhizobium inoculation than untreated treatment.

Table 3: Effect of Biofertilizer and Phosphorus on Yield attributes of Chickpea

S.No	Treatments	Pods/plant (No)	Seeds/pod (No.)	Test weight (g)	Seed Yield (t/ha)	Stover Yield (t/ha)
1	Rhizobium+Phosphorus 20kg/ha	29.74	1.33	197.83	1.67	2.57
2	Rhizobium+Phosphorus 40kg/ha	28.84	1.22	203.81	1.83	2.98
3	Rhizobium+Phosphorus 60kg/ha	31.74	1.56	229.89	1.92	3.26
4	PSB+Phosphorus 20kg/ha	28.19	1.11	218.58	1.87	2.78
5	PSB+Phosphorus 40kg/ha	30.94	1.11	198.86	1.69	2.61
6	PSB+Phosphorus 60kg/ha	27.17	1.22	212.48	1.88	2.84
7	Rhizobium+PSB+Phosphorus 20kg/ha	29.26	1.11	206.18	1.72	2.75
8	Rhizobium+PSB+Phosphorus 40kg/ha	32.86	1.78	237.04	1.98	3.47
9	Rhizobium+PSB+Phosphorus 60kg/ha	34.29	1.89	245.76	2.07	3.58
10	Control	26.65	1.11	194.39	1.65	2.50
	F test	S	S	S	S	S
	SEm(±)	0.49	0.12	5.29	31.39	66.01
	CD (P=0.05)	1.46	0.36	15.72	93.27	196.1

4. SUMMARY

Highest plant height (54.94cm), Number of Nodules per plant (36.28), Plant dry weight (23.71g), maximum number of pods per plant (34.29), number of seeds per pod (1.89), Test weight (245.76g), Seed yield (2.07t/ha) and Stover yield (3.58t/ha) were recorded with the treatment Rhizobium+PSB 20g/kg seed + Phosphorus 60kg/ha.

At the same time higher Gross return (108523.7 INR/ha), Net return (75364.02 INR /ha) and Benefit Cost Ratio (2.27) was obtained in treatment Rhizobium+PSB + Phosphorus 60kg/ha.

5. CONCLUSION

From the above study it is concluded that the Treatment 9 with Rhizobium+PSB + Phosphorus 60kg/ha gave best results and performed better in growth and yield parameters.

REFERENCES

1. Ahlawat I P S, Gangaiah B and Zahid M A. Nutrient Management in Chickpea. Chickpea Breeding and Management, CABI, Walling Ford, UK. 2007; PP: 213-232.
2. Akansha Singh, A.K. Sachan, Vipin Kumar, R.K. Pathak and ShubhamSrivastav. Effects of Phosphorus with Biofertilizers on Yield and Nutrient Content of Chickpea (*Cicerarietinum L.*) under Central Uttar Pradesh Condition. *International Journal of Current Microbiology and Applied Sciences*. 2018;10(02):2228-2234.
3. Annual Report 2016-17. Ministry of Agriculture, Cooperation and Farmers Welfare (MoA&F), Government of India
4. Arya RL, Kushwaha BL, Singh BN. Effect of phosphorus management on growth, yield attributes and yield of maize –chickpea cropping system. *Indian J. Pulses Res.* 2002; 15(2):161-165.
5. Dahiya S, Mehar S, Singh M. Relative growth performances of chickpea genotypes to irrigation and fertilizers application. *Haryana J. Agron.* 1993; 9(2): 172-175.
6. Dinesh Kumar, L. K. Arvadiya, K. L. Desai, V. P. Usadadia and A. M. Patel. Growth and yield of chickpea (*Cicerarietinum L.*) as influenced by graded levels of fertilizers and biofertilizers. *An International quarterly journal of life sciences*. 2015; **10**(1): 335-338.
7. FAO 2012 Agriculture Data. <http://faostat.fao.org/site/567>

8. Jarande NN, Mankar PS, Khawale VS, Kanse AA, Mendhe JT. Response of chickpea (*Cicerarietinum*.L) to different levels of phosphorus through inorganic and organic sources J.Soils Crops. 2006; 16:240-243.
9. Jitendra Kumar Yadav, Mahendra Sharma, RN Yadav, SK Yadav and SarojYadav. Effect of different organic manures on growth and yield of chickpea (*Cicerarietinum* L.) Journal of Pharmacognosy and Phytochemistry. 2017; 6(5): 1857-1860.
10. Meena LR, Singh RK, Gautam RC. Effect of moisture conservation practices, phosphorus levels and bacterial inoculation on chickpea under rainfed condition. Indian J. Agron. 2002; 47: 398-404.
11. Pramanik K, Singh RK. Effect of levels and mode of phosphorus and biofertilizers on chickpea under dryland conditions. Indian J. Agron. 2003; 48: 294-296.
12. Rajneesh Singh, TejPratap, Durgesh Singh, Ghanshyam Singh and Abhinav Kumar Singh. Effect of phosphorus, Sulphur and biofertilizers on growth attributes and yieldof chickpea (*Cicerarietinum* L.) *Journal of pharmacognosy and phytochemistry*2018; 7(2): 3871-3875.
13. Singh, A.K.; Singh, B. and Singh, H.C. Response of chickpea (*Cicerarietinum*L.) to fertilizer phosphorus and zinc application under rainfed condition of eastern U.P. *Indian J. Dryland Agric. Res. & Dev.*,2005;20(2): 114-117.
14. Singh, G.; Sekhon, H.S. and Sharma, P. Effect of irrigation and biofertilizer on water use, nodulation, growth and yield of chickpea (*Cicerarietinum* L.). *Agron. Soil Sci.*, 2011; 57(7): 715-726.
15. Tambekar DH, Gulhane SR, Somkuwar DO, Ingle KB, Kanchalwar SP. Potential Rhizobium and phosphate solubilizers as a biofertilizers from saline belt of Akola and Buldhana district, India. *Research Journal of Agriculture and Biological Sciences*. 2009; 5(4):578-582.
16. Thenua OVS, Singh SP, Shivakumar BG. Productivity and economics of chickpea (*Cicerarietinum*. L) + fodder sorghum (*Sorghum bicolor*) cropping systems as influenced by phosphorus sources, biofertilizers and irrigation to chickpea. *Indian J. Agron.* 2010; 55: 22-27.
17. Umadevi M, Ganesan NM. Analysis for yield and quality characters in blackgram (*Vignamungo*(L.) Hepper). *Legume Res.* 2007; 30(3):197-200.