

Original Research Article

Hepatotoxic Nature of Potash (Kaun) in Wistar Rats

Augustine I. Airaodion^{1*}, Aanu P. Agunbiade¹, Ayodeji A. Adediji², Anthony U. Megwas³, Emmanuel B. Ayita⁴, Simeon O. Oluba⁵ and Ojo J. Osunmuyiwa⁶

¹Department of Biochemistry, Federal University of Technology, Owerri, Imo State, Nigeria.

²Department of Biochemistry, Federal University of Technology, Akure, Ondo State, Nigeria.

³Department of Optometry, Federal University of Technology, Owerri, Imo State, Nigeria.

⁴Department of Biochemistry, Federal University Oye-Ekiti, Ekiti State, Nigeria.

⁵Clinical Biochemistry Unit, Department of Medical Laboratory Service, Inland Medical Centre, Ikare-Akoko, Ondo State, Nigeria.

⁶Department of Biomedical Science and Public Health Technology, Margaret Mosunmola College of Health Science and Technology, Owo, Ondo State, Nigeria.

Authors' Contributions

This work was carried out in collaboration with all authors. Author AIA conceptualized and designed the study, and also wrote the manuscript. Author OJO and SOO managed the analyses of the study. Author APA and AUM managed the literature searches. Author EBA wrote the protocol while Author AAA performed the statistical analysis. All authors read and approved the final manuscript.

ABSTRACT

Background: The use of potash as food additive without a recourse to its adverse effect is on the increase in Nigeria.

Aim: This study is designed to assess its effect on hepatic indices of Wistar rats.

Methodology: Potash was locally sourced in a market in Owerri, Imo State, Nigeria. Thirty Wistar rats were acclimatized for seven days, grouped into five and it comprises of six animals respectively. Group A were given distilled water, whereas the treatment groups received 250 mg/kg, 500 mg/kg, 750 mg/kg and 1000 mg/kg body weight of potash for twenty-eight days via oral route of administration. The Wistar rats were anaesthetized using diethyl ether, sacrificed then whole blood needed for the study were obtained through cardiac puncture. Biochemical parameters needed for liver function test were analyzed using standard protocol from the manufacturer.

Results: It was revealed that potash administration at higher dose is toxic and perturbs hepatic biomarkers.

Conclusion: From the results of this study, potash is hepatotoxic; therefore, discontinuation of potash consumption needs to be recommended.

Keywords: Food Addictive, Hepatotoxic, Potash

1. INTRODUCTION

Liver serves as one of the major organs that perform some essential functions for proper

Comment [KM1]: Delete of

Comment [KM2]: Change to was

Comment [KM3]: The oral

Comment [KM4]: The spelling of anaesthetized should be changed

Comment [KM5]: a higher dose

Comment [KM6]: The liver

body functionality such as ability to metabolized and detoxified various compounds, balance homeostasis, ensured essential growth coupled with provision of adequate energy and nutrient needed by the body [1]. Injury of the liver could be as a result of toxicity produced by various toxicants and infectious agents that do affect the liver [2]. Many diseases ensued from the hepatic dysfunction such as jaundice, cirrhosis and fatty liver, these have greatly threatened the health of so many people across the world [3]. The occurrence of lingering liver diseases across the world is about 18.5%, with cirrhosis taken about 4.5-9.5% which eventually led to the death of about 2 million people yearly. What we consumed serves as one of the contributing factors to the occurrence of liver problem.

Potash refers to one of the salts that been mined; which is made up of potassium that can easily dissolve in water, of which the name potash was obtained from the plant burn to ashes which is then dissolved in water housed by a pot, and this serves as the major production method on before the use of technology [4]. Potash production was so great globally with is annual production greater than 30 million tonnes because of is widely application as fertilizer. The major component of various kinds of fertilizer-potash is made up of potassium. And the first production of potassium was achieved through electrolysis of caustic potash (potassium hydroxide) in the year 1807 [5]. Ash burners are the set of people that do produces pot ash called potassium carbonate with the use of an old method by burning wood into ashes, leach and evaporate it in a big pot with deposit of a white residue tagged as pot ash [6]. Most of the wood that are burn into ashes, 10% of it are recovered back as pot ash. Recently development led to the name potash as an acceptable name worldwide rather than potassium salts and byproducts [7]. Potash refers to "Kaun" or "Akanwu" by some ethnics group in Nigeria which is majorly used for cooking food. Beans are one of the foods that need the used of potash to make it done quickly [8]. And also, to increase the texture and retain the green colour of some vegetables soup such as ewedu and Okro [9]. But yet there is still no recommended dosage of potash to be consume in a daily meal

of every Nigerian. Hence, this study was designed to examine the likely effect of potash on the liver.

2. METHODOLOGY

2.1 Experimental Design

Potash was locally sourced in a market in Owerri, Imo State, Nigeria and was carefully preserved to avoid contamination. A thirty Wistar rat with a weight range between 145 and 160 grams were purchase and allowed to acclimatized for seven (7) days with free access to food and drinkable water. They were housed in a clean and well-ventilated cage environment under a standard atmospheric condition for laboratory animals. This treatment was in accordance with the guide prepared for experimental animals by the National Academy of Science [10]. The animals were selected into five (5) major groups according to their body weight; with group A received only distilled water while the treatment groups were given 250 mg/kg, 500 mg/kg, 750 mg/kg and 1000 mg/kg body weight of potash for good twenty-eight days through oral administration. The Wistar rats were anaesthetized using diethyl ether, sacrificed, then whole blood needed for the study were obtained through cardiac puncture.

2.2 Determination of Hepatic Indices

Commercially available enzyme Randox kits were purchased to determine the activities of Aspartate Aminotransferase (AST) and Alanine Aminotransferase (ALT) according to the procedure described by Reitman and Frankel [11]. Alkaline Phosphatase (ALP) activity was determined by Phenolphthalein Monophosphate method described by Babson et al. [12]. Amylase inhibition assay was determined by the method of Bernfield [13]. Biorex diagnostic kit was used to quantified activities of lipase based on the method stated by Lorentz [14]. Total bilirubin concentration was determined by diazo method described by Royden and Alfred [15]. Conjugated bilirubin concentration was determined by the method of Compennolle [16]. Subtraction of conjugated bilirubin from total bilirubin produced the quantity of unconjugated bilirubin to be quantified.

2.3 Statistical Analysis

Comment [KM7]: The ability

Comment [KM8]: an incorrect form of the verb- metabolize

Comment [KM9]: an incorrect form of the verb - detoxify

Comment [KM10]: an incorrect form of the verb- ensure

Comment [KM11]: the provision

Comment [KM12]: delete as

Comment [KM13]: delete the

Comment [KM14]: change to which

Comment [KM35]: correct to Thirty Wistar rats

Comment [KM36]: correct to were

Comment [KM37]: change to be

Comment [KM15]: problems

Comment [KM16]:

Comment [KM17]: correct into potash

Comment [KM18]: change to its

Comment [KM19]: change to its

Comment [KM20]: correct into wide

Comment [KM38]: correct to was

Comment [KM21]: change to produce

Comment [KM22]: correct to potash

Comment [KM23]: change to leaching

Comment [KM24]: correct to evaporating

Comment [KM25]: a deposit

Comment [KM39]: add the

Comment [KM26]: potash

Comment [KM27]: change to are burned

Comment [KM28]: correct to is

Comment [KM29]: correct to potash

Comment [KM40]: correct to quantify

Comment [KM30]: change to etnic groups

Comment [KM31]: correct to the use

Comment [KM32]: correct to color

Comment [KM33]: change to vegetable

Comment [KM34]: change to be consumed

One way analysis of variance was used to compare mean while the results were expressed as mean \pm standard deviation and the graph were drawn using Graph Pad Prism software version 5.00. The results were considered to be significant when $p < 0.05$.

3. RESULTS

The results of this study are presented in figures 1-11. No significant difference was observed when the activities of ALT and AST in animals treated with lower doses (250 and 500 mg/kg) of potash were compared with those in the control group at $P < 0.05$. A significant increase was however observed in the activities of ALT and AST in animals treated with higher doses (750 and 1000 mg/kg) of potash when compared with those in the control group (Figures 1 and 2). ALP activity was observed to increase in experimental animals when compared with those of the control animals. This elevation was however not significant when animals treated with 250 mg/kg body weight of potash were compared with the control group at $P < 0.05$ (Figure 3). No significant difference was observed in the concentrations of total protein and albumin in animals treated with lower doses

(250 and 500 mg/kg) of potash when compared with that of the control group at $P < 0.05$. A significant increase was however observed in the concentrations of total protein and albumin in animals treated with higher doses (750 and 1000 mg/kg) of potash when compared with those in the control group (Figures 4 and 5). The concentration of globulin was only significant when animals treated with 500 and 1000 mg/kg body weight of potash were compared with those of the control animals (Figure 6). Administration of potash increased total bilirubin concentration when compared with those in control animals. The increase was significant when animals treated with 500 and 1000 mg/kg of potash were compared with those in the control group at $P < 0.05$ respectively (Figure 7). No significant difference was observed in the levels of conjugated bilirubin in experimental animals when compared with those in control group at $P < 0.05$ (Figure 8). A significant increase was observed in the level of unconjugated bilirubin (except the group treated with 750 mg/kg) when compared with those in control group (Figure 9). The potash was observed to inhibit the activities of amylase and lipase (Figures 10 and 11) respectively in a dose-dependent manner.

Comment [KM41]: change to One-way

Comment [KM42]: change to meanvalues,

Comment [KM43]: correct to was

b




Figure 1: Effect of Potash on the Activity of Alanine Amino Transferase (ALT) of Animals after 28 days of Treatment

Results are presented as mean \pm SD with $n = 6$. Bars with different letters are significantly different at $P < 0.05$




Figure 2: Effect of Potash on the Activity of Aspartate Amino Transferase (AST) of Animals after 28 days of Treatment

Results are presented as mean \pm SD with $n = 6$. Bars with different letters are significantly different at $P < 0.05$




Figure 3: Effect of Potash on the Activity of Alkaline Phosphatase (ALP) of Animals after 28 days of Treatment

Results are presented as mean \pm SD with n = 6. Bars with different letters are significantly different at $P < 0.05$




Figure 4: Effect of Potash on the Concentration of Total Protein of Animals after 28 days of Treatment

Results are presented as mean \pm SD with $n = 6$. Bars with different letters are significantly different at $P < 0.05$




Figure 5: Effect of Potash on the Concentration of Albumin of Animals after 28 days of Treatment

Results are presented as mean \pm SD with n = 6. Bars with different letters are significantly different at $P < 0.05$




Figure 6: Effect of Potash on the Concentration of Globulin of Animals after 28 days of Treatment

Results are presented as mean \pm SD with $n = 6$. Bars with different letters are significantly different at $P < 0.05$




Figure 7: Effect of Potash on the Concentration of Total Bilirubin of Animals after 28 days of Treatment

Results are presented as mean \pm SD with n = 6. Bars with different letters are significantly different at $P < 0.05$




Figure 8: Effect of Potash on the Concentration of Conjugated Bilirubin of Animals after 28 days of Treatment

Results are presented as mean \pm SD with n = 6. Bars with different letters are significantly different at $P < 0.05$




Figure 9: Effect of Potash on the Concentration of Unconjugated Bilirubin of Animals after 28 days of Treatment

Results are presented as mean \pm SD with n = 6. Bars with different letters are significantly different at $P < 0.05$




Figure 10: Effect of Potash on the Activity of Amylase of Animals after 28 days of Treatment

Results are presented as mean \pm SD with n = 6. Bars with different letters are significantly different at $P < 0.05$




Figure 11: Effect of Potash on the Activity of Lipase of Animals after 28 days of Treatment

Results are presented as mean \pm SD with n = 6. Bars with different letters are significantly different at $P < 0.05$

4. DISCUSSION

Administration of potash to Wistar rat for 28 days revealed significant elevation of alanine aminotransferase (ALT), aspartate aminotransferase (AST) and alkaline

phosphatase (ALP) activities (Figures 1, 2 and 3) at higher dose of 750 mg/kg and 1000 mg/kg body weight than at lower dose of 250 mg/kg and 500 mg/kg body weight. It has been reported that an increase in the enzymatic

Comment [KM45]: change to the higher dose

Comment [KM46]: correct to doses

Comment [KM44]: correct to rats

Comment [KM47]: correct to in comparison with

Comment [KM48]: change to the lower dose

Comment [KM49]: correct to doses

Formatted: Strikethrough

activity of ALT, AST and ALP in the serum directly reflects hepatocellular damage [17]. These findings, therefore suggested that potash may be hepatotoxic at high doses, which is consistence with the finding of Iweka *et al.* [8] who revealed in their findings that as the dosage of potash increases, concomitantly increases activities of AST, ALT and ALP after administration for 21 days. This could be that exposure of animals to potash stimulated the transcription of the genes involved in glucose uptake, glycolysis and lipogenesis [18]. It has been documented that synthesis of cyclic adenosine monophosphate (cAMP) can be inhibited through the repression of inducible operon by glucose such as Lac operon [19]. Allosteric protein needs the help of nucleotide called cAMP for proper activation to effectively binds to the promoter catabolite activator protein (CAP) site, then enhances the joining of ribonucleic acid polymerase with the promoter to start transcription processes and then, the presence of cAMP is needed before joining CAP, this needs to join deoxyribonucleic acid to enable transcription processes [20]. Adenylase cyclase (AC) activity is blocked when glucose is available, hence the production of cAMP from Adenosine Triphosphate (ATP) needed the activity of AC [21]. But low synthesis of cAMP will eventually lead to CAP inactiveness therefore, altered the processes of transcription. Thus, these inducible enzymes are reduced through direct effect on glucose, therefore caused reduction of cyclic AMP level. Administration of potash at high doses might have elevated cAMP in treated rats, thus the significant increase in these inducible enzymes. ALT is considered as one of the major biomarkers for liver damage because it is solely confined to the liver, unlike AST which is also abundantly present in other body organs such as the kidneys, brain, and hearts [22,23]. The significant increase in the activities of ALT, AST and ALP in animals treated with high doses of potash revealed that potash is considered toxic at high dose to the liver.

Similarly, concentrations of total protein and albumin were significantly elevated in animals treated with high dose of potash. This elevation suggested that metabolizing ability of the liver ensued from the administration of potash at higher doses has been compromised. This might have increased the functional activity of the liver by interfering with the state of synthesis, degradation, removal or clearance of total

protein and albumin from the animals' body systemic circulation [24]. This increase in total protein could lead to dehydration which is harmful to the cellular homeostasis [25]. And have a negative effect on the metabolic activities of the liver, which will eventually result in disturbances of the health of the animal. The major function of albumin includes binding and transportation of metal ions, bilirubin, and drugs. Its quantification is required to evaluate the synthetic function of the liver [26]. Significant increase in the level of these parameters indicates that potash stimulated their synthesis in the liver at a higher dosage of 750 mg/kg and 1000 mg/kg body weight. Serum protein levels are controlled through synthesis in the liver and its levels thus reflect the synthetic ability of the liver [26].

Bilirubin refers to as one of the products produced during degradation of hemoglobin. However, high level of bilirubin in the blood lead is toxic to the body system, which we eventually cause different medical conditions such as jaundice, hyperbilirubinemia-induced auditory dysfunction and neurotoxicity resulting in brain damage [27]. Whereas slight elevation of unconjugated bilirubin in the blood serves as an antioxidant, that do safeguard the body from heart related diseases coupled with development of tumor [28]. New investigation about reduction in the concentration of direct bilirubin may cause heart and brain defect. Serum bilirubin levels are often enhanced under a variety of clinical conditions. In the circulation of blood, bilirubin is bound to serum albumin, which prevents its potential toxicity thought to be caused by free bilirubin [29]. Despite its high-affinity of binding to albumin, bilirubin is rapidly and selectively taken up by the liver, biotransformed upon conjugation with glucuronate, and secreted into bile [28]. Thus, bilirubin is converted into bilirubin glucuronic acid in the liver and excreted along with bile. In this study, there was no significant change in the serum conjugated (direct) bilirubin concentrations. However, there was a significant increase in the serum levels of total and unconjugated (indirect) bilirubin in the serum of the Wistar that received potash. The elevation in the concentration of indirect serum bilirubin (unconjugated bilirubin) recorded might due to tissue injuries or damage caused by the toxic substance the animals were exposed to [30].

Comment [KM50]: correct to consistent

Comment [KM51]: change to consistent

Comment [KM52]: correct style

Comment [KM65]: correct to a significant

Comment [KM53]: add the synthesis

Comment [KM54]: add the synthesis

Comment [KM55]: add the synthesis

Comment [KM56]: change to the inducible

Comment [KM57]: correct to the inducible

Comment [KM58]: correct to bind

Comment [KM59]: correct to bind

Comment [KM60]: correct to Adenylate

Comment [KM61]: add the low

Comment [KM66]: add the developmant

Comment [KM62]: correct to altering

Comment [KM63]: correct to causing the reduction

Comment [KM67]: correct to there were not significant changes...

Comment [KM64]: the high

Comment [KM68]: add be due

Administration of potash significantly reduced activities of amylase and lipase as the dosage of potash increases, and the lower is effect on each enzyme activities respectively. Indicating that at higher dose potash reduced amylase and lipase activities in the serum of the experimental animals. Amylase serves as one of the major enzymes needed during degradation of starch [20]. β -cells in the Islets of Langerhans are been destroyed by autoimmune reactions that occurred in the pancreas, which is facilitated by formation of reactive oxygen species in the leucocytes [31, 32]. During digestion and uptake of carbohydrates; when some of the key major enzymes were inhibited, this can easily reduce blood sugar level after taken carbohydrate diet. Hence, this may serve as one of the means of managing hyperglycemia associated with type 2 diabetes [33,34]. Whereas, Lipase is the enzyme responsible for digestion and absorption of triglycerides [20,26]. Its inhibition is one of the widest studied methods used to determine the potential activity of natural products to inhibit dietary fat absorption. Decrease in energy intake from dietary fat through inhibition of this enzyme may be an excellent strategy to prevent and treat obesity [35].

4. CONCLUSION

It was revealed in our findings that potash administration at higher dose is toxic and perturbs hepatic biomarkers, therefore discontinuation of potash consumption needs to be recommended.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing

company rather it was funded by personal efforts of the authors.

REFERENCES

1. Mahmood DN, Mamat SS, Kamisan HF , Yahya F, Kamarolzaman FFM, Nasir N , Mohtarrudin N, Tohid, and Zakaria AZ . Amelioration of Paracetamol-Induced Hepatotoxicity in Rat by the Administration of Methanol Extract of *Muntingia calabura* L. Leaves. BioMed Research International. 2014 , 1 -10.
2. Airaodion AI, Ogbuagu U, Ekenjoku JA, Ogbuagu EO, Airaodion EO, Okoroukwu VN. Hepato-protective efficiency of ethanol leaf extract of *Moringa oleifera* against hydrocarbon exposure. *International Journal of advances in Herbal and Alternative Medicine*. 2019;03(01):32-41.
3. Airaodion AI, Ogbuagu EO. Effect of *Cyperus esculentus* L. (tiger nut) milk on hepatic and renal indices of Wistar rat. *Asian Journal of Research in Nephrology*. 2020;3(2):10-16.
4. Davy H. On some new phenomena of chemical changes produced by electricity in particular the decomposition of the fixed alkalies, and the exhibition of the new substances that constitute their bases; and on the general nature of alkaline bodies". *Philosophical Transactions of the Royal Society of London*; 1808;98: 32.
5. Knight D. Humphry Davy; Science and Power. Oxford: Blackwell. 1992; p 66.
6. Dennis K. "Potash". 2005 Minerals Handbook. United States Geological Survey. 2006; p. 58.1.
7. The World Potash Industry. Past, Present and Future. New Orleans, LA: 50th Anniversary Meeting; the Fertilizer Industry Round Table 2000.
8. Iweka FK, Dic-Ijiewere OE, Oaikhena F, Bankole JK, Festus OO, Dada FL. The Effect Of Potash On Liver Function Of Wister Rats. *International Journal of Herbs and Pharmacological Research*. 2016;5(1): 13 – 20.
9. Okpala B. Benefits of Kaun Potash (Akanwu). Blog by Blessing Okpala. Global Food book Recipes for life. 2015; <https://globalfoodbook.com/benefits-of->

Comment [KM80]: add the personal

Comment [KM69]: unclear

Comment [KM70]: add the higher

Comment [KM71]: add the degradation

Comment [KM72]: correct to levels

Comment [KM73]: correct to taking

Comment [KM74]: add the digestion

Comment [KM75]: add A decrease

Comment [KM76]: correct to standards

Comment [KM77]: standards

Comment [KM78]: correct to used

Comment [KM79]: the preposition for is redundant

- kaun-potash-akanwu/
10. NAS. National Academy of Science Guide for the Care and Use of Laboratory Animals. Eighth Edition. 2011.
11. Reitman S, Frankel S. A colorimetric method for determination of serum glutamate oxaloacetate and glutamic pyruvate transaminase. *American Journal of Clinical Pathology*. 1957;28:56-58.
12. Babson AL, Grecley SJ, Coleman CM, Phillips GE. The use of phenolphthalein in monophosphate as a substrate for serum alkaline phosphatase. *Clinical Chemistry*. 1966;12:482.
13. Bernfield P. Enzymes of starch degradation and synthesis. *Adv Enzymol*, 1951;12:379–428.
14. Lorentz K. Lipase. In: Thomas L, editor. Clinical laboratory diagnostics. 1st ed. Frankfurt: TH-Books Verlagsgesellschaft. 1998; p. 95-97.
15. Royden NR, Alfred P. A New Diazo Method for the Determination of Bilirubin. *Clinical Chemistry*. 1962;8(6):570-578.
16. Compennolle F. Bilirubin conjugates: isolation, structure analysis and synthesis. Bilirubin, Vol 1: Chemistry. Boca Raton, FL: CRC Press, 1982;2:59-74.
17. Ogbuagu EO, Airaodion AI, Ogbuagu U, Airaodion EO. Prophylactic propensity of methanolic extract of *Vernonia amygdalina* leaves against acute ethanol-induced oxidative stress in Wistar rats. *International Journal of Bio-Science and Bio-Technology*. 2019;11(7):37-46.
18. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Ogbuagu U, Airaodion EO. Therapeutic effect of methanolic extract of *Telfairia occidentalis* leaves against acute ethanol-induced oxidative stress in Wistar rats. *International Journal of Bio-Science and Bio-Technology*. 2019;11(7):179-189.
19. Airaodion AI, Akunne PN, Njoku OC, Oladosu NO, Megwas AU. Effect of Bambara nut on hepatic biomarkers of Wistar rats. *International Research Journal of Gastroenterology and Hepatology*. 2021;4(1): 26-38.
20. Njoku OC, Airaodion AI, Osuagwu OL, Oladosu NO Megwas AU. Hepatoprotective Potential of Alkaloid Extracts from *Vitex doniana* and *Ficus thonningii* Leaves in Alloxan-Induced Diabetic Rats. *International Research Journal of Gastroenterology and Hepatology*. 2021;4(1): 48-63.
21. Airaodion AI, Akinmolayan JD, Ogbuagu EO, Esonu CE, Ogbuagu U. Preventive and therapeutic activities of methanolic extract of *Talinum triangulare* leaves against ethanol-induced oxidative stress in Wistar rats. *International Journal of Bio-Science and Bio-Technology*. 2019;11(7):85-96.
22. Ogbuagu EO, Airaodion AI, Okoroukwu VN, Ogbuagu U, Ekenjoku JA. Effect of Monosodium Glutamate on Body Weight and Alanine Aminotransferase Activity in Wistar Rats. *International Research Journal of Gastroenterology and Hepatology*. 2019;2(2):1-8.
23. Airaodion AI, Ngwogu AC, Ekenjoku JA, Ngwogu KO. Hepatoprotective potency of ethanolic extract of *Garcinia kola* (heckel) seed against acute ethanol induced oxidative stress in Wistar rats. *International Research Journal of Gastroenterology and Hepatology*. 2020;3(2):1-10.
24. Airaodion AI, Ogbuagu EO, Ewa O, Ogbuagu U, Awosanya OO, Adekale OA. Ameliorative efficacy of methanolic extract of *Corchorus olitorius* leaves against acute ethanol-induced oxidative stress in Wistar rats. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2019;7(6):1-9.
25. Airaodion AI, Ogbuagu EO, Ogbuagu U, Adeniji AR, Agunbiade AP, Airaodion EO. Hepatoprotective effect of *Parkia biglobosa* on acute ethanol-induced oxidative stress in Wistar rats. *International Research Journal of Gastroenterology and Hepatology*. 2019;2(1):1-11.
26. Ogbuagu EO, Uneke PC, Airaodion AI, Nweke IN, Ogbuagu U. Hepatotoxic effect of *Xylopiia aethiopica* fruit in Wistar rats. *International Research Journal of Gastroenterology and Hepatology*. 2021;4(1):1-16.
27. Shapiro SM. Bilirubin toxicity in the developing nervous system. *Pediatr neurol.*, 2003;29: 410-421.
28. Airaodion AI, Ene AC, Ogbuagu EO, Okoroukwu VN, Ekenjoku JA, Ogbuagu U. Biochemical changes associated with consumption (by rats) of “garri” processed by traditional and instant mechanical methods. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2019;2(4): 1-11.
29. Perlstein TS, Pande RL, Creager MA, Weuve J, Beckman JA. Serum total

- bilirubin level, prevalent stroke, and stroke outcomes: NHANES 1999-2004. *American Journal of Medicine*. 2008;121: 781–788.
30. Ogbuagu EO, Airaodion AI, Ogbuagu U, Nweke IN, Uneke PC. Nephrotoxicity of ethanol extract of *Xylopi aethiopica* fruit in Wistar rats. *International Journal of Advances in Nephrology Research*. 2021;4(1):1-16.
31. Oberley LW. Free radicals and diabetes. *Free Radic Biol Med*. 1998;5:113–124.
32. Airaodion AI, Emaekwu SA, Osunmuyiwa OJ, Megwas AU, Ayita EB, Oluba SO. Nephrotoxic Nature of Potash (Kaun) in Wistar Rats. *International Journal of Health, Safety and Environment*. 2021; 4(7):27-33
33. Pinto MS, Ranilla LG, Apostolidis E, Lajolo FM, Genovese MI, Shetty K. Evaluation of antihyperglycemia and antihypertension potential of native Peruvian fruits using *in vitro* models. *J Med Food*. 2009;12:278–291.
34. Shim YJ, Doo HK, Ahn SY, Kim YS, Seong JK, Park IS, Min BH. Inhibitory effect of aqueous extract from the gall of *Rhus chinensis* on α -glucosidase activity and postprandial blood glucose. *Journal of Ethnopharmacology*. 2003;85, 283–287.
35. Sosnowska D, Podsędek A, Redzynia M, Zyzelewicz D. Effects of Fruit Extracts on Pancreatic Lipase Activity in Lipid Emulsions. *Plant Foods Hum Nutr*. 2015;70(3):344-550.