

# **Fortification of Nano Whey Protein-Meniran Phyllanthine (*Phyllanthus niruri* L.) Encapsulated Casein Hydrolysates on Physicochemical Properties of Skimmed Milk Powder**

## **ABSTRACT**

The research was held from August 2021 until January 2022 in Laboratory Livestock Product Technology Universitas Brawijaya, Malang, Indonesia. The research was conducted to obtain skimmed milk powder fortified nano whey protein-meniran phyllanthine encapsulated casein hydrolysates with better physicochemical properties. The research material used was skimmed milk powder with different concentrations of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates, designed by 5 treatments and 3 replications consisting of T0: 0% (control), T1: 2% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates, T2: 4% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates, T3: 6% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates, T4: 8% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates. The variables tested were solubility, turbidity, sedimentation, water activity (aw) and hygroscopicity. The research method was a laboratory experiment analyzed by Analysis of Variance (ANOVA) using Duncan's Multiple Range Test (DMRT) if there were differences. The results showed a significant difference ( $P < 0.05$ ) on water activity (aw). However, there was no significant difference ( $P > 0.05$ ) on skimmed milk powder's solubility, turbidity, sedimentation, and hygroscopicity. Average data of physicochemical properties in this study were solubility (57.983-75.295%), turbidity (0.459-0.518), sedimentation (0.036-0.043), water activity (0.434-0.467), and hygroscopicity (4.601-5.528%). In conclusion, fortified nano whey protein-meniran phyllanthine encapsulated casein hydrolysates in skimmed milk powder can be used as a product with better physicochemical properties.

**Keywords:** *casein hydrolysates; encapsulated; fortified; nano whey protein-meniran phyllanthine; skimmed milk powder*

## **1. INTRODUCTION**

Skimmed milk powder is one of the dairy products through a heating process by hot air to produce low moisture content. Lower moisture content will prevent microorganism growth and extend the shelf life of

skimmed milk powder. Skimmed milk powder is usually processed using spray drying, which produces from a liquid material into dried powder using hot air [1]. The principle of the spray drying method was atomization, which sprayed liquid into fine droplets with hot air and became dried particles.

However, the heating process during manufacture skimmed milk powder reduced water content and other nutritional contents such as protein and vitamins, which would require fortification. The fortification was an effort to enrich nutrients to improve nutritional quality and human health benefits [2].

Meniran is an herbal plant from Indonesia that has health benefits with its bioactive compounds. Bioactive compounds of meniran consist of lignans (phyllanthine, hypophyllanthine, nirantin, linetetratin), flavonoids (quercetin and rutin), phenolics, alkaloid, saponin and tannin. The quality of meniran is based on a single marker compound from the lignan group [3,4]. Phyllanthine bioactive can be obtained by the extraction method. Microwave-assisted extraction is one of the extraction methods often used to extract phytochemicals in the plant using water solvent and microwave energy [5]. Microwave energy will be converted into heat energy, plant tissue and cell walls will cause lysis and extracted the phyllanthine bioactive. The application of meniran phyllanthine to one of the dairy by-products, such as whey protein, can be used as a functional food and is easy to absorb.

Whey protein is obtained from cheese making and casein production by-products, which still have high protein content. Whey protein mainly by  $\beta$ -lactoglobulin ( $\beta$ -Lg, 50%),  $\alpha$ -lactalbumin ( $\alpha$ -La, 20%), bovine serum albumin (BSA, 10%), immunoglobulins (10%) and peptones protease (<10%) [6]. Whey protein was rich in essential amino acids and complex structures consisting of primary, secondary, tertiary and quaternary structures [7], so if interacted with meniran bioactive can change the structure and functional properties of the protein. Amino acids from whey protein are composed of one hydrogen atom when heated at 65-70°C, which can interact with hydroxyl group from meniran through covalent and non-covalent binding (hydrophobic interactions, van der Waals and hydrogen bonds) [8]. One of the

disadvantages of whey protein is easily denatured when heated up to 70°C, so an encapsulation process is needed.

Encapsulation is alternate protection by adding active substances into particles to coat and protect core material from high pressure and temperature during the heating process and reduce product stickiness [9]. This research used casein hydrolysates as encapsulant composed of  $\alpha$ s1,  $\alpha$ s2,  $\beta$ - dan  $\kappa$ -casein fragments [10]. The properties of casein hydrolysates that do not react to other food constituents were a concern if applied as a functional food or supplement [11]. Based on the description above, the analysis of solubility, turbidity, sedimentation, water activity ( $a_w$ ) and hygroscopicity of fortification nano whey protein-meniran phyllanthine encapsulated casein hydrolysates was carried out to evaluate the quality of skimmed milk powder based on their physicochemical properties.

## 2. MATERIALS AND METHODS

### 2.1 Materials

The materials used in the research were skimmed milk powder, nano whey protein-meniran phyllanthine encapsulated casein hydrolysates, SDS 0.1%,  $\text{NaN}_3$ ,  $\text{Na}_2\text{SO}_4$  and aquades. The equipment used was  $a_w$  meter (Rotronic® Instruments), microwave (Sharp), spray dryer (Buchi), glass funnel, beaker glass, measure glass, centrifugator, centrifuge tube, analytical scale, oven 105°C (WT-Binder), desiccator, hotplate and magnetic stirrer (SBS-06), mini hand mixer, UV-Vis spectrophotometer (721-Faithful), micropipette (HumaPette) and filter paper (Whatman No.1 and 4).

### 2.2 Methods

The research method was a laboratory experiment with 5 treatments and 4 replications. Fortification skimmed milk powder with different concentrations of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates consisting of T0: 0% (control), T1: 2% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates, T2:

4% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates, T3: 6% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates, T4: 8% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates.

### 2.2.1 Meniran Phyllanthine

#### Extraction

Extraction of meniran phyllanthine based on [12]. Meniran powder as much as 3 g diluted in 100 mL aquades and macerated for 24 hours. Meniran solution was extracted with microwave at 70°C for 10 minutes (one-minute radiation, two minutes off and so on until ten minutes). After that, The extract of meniran phyllanthine was filtered using filter paper (Whatman No.4).

### 2.2.2 Interaction Whey Protein and Meniran Phyllanthine

The interaction according to [8], 5 g of whey protein were heated using hotplate stirrer at 65-70°C for 40 minutes and phyllanthine meniran was gently added into whey protein solution until it reached 100 mL volume.

### 2.2.3 Encapsulation Nano Whey Protein-Meniran Phyllanthine with Casein Hydrolysates

Four grams of casein hydrolysates were added into 100 mL nano whey protein-meniran phyllanthine solution and homogenized. The solution was allowed to stand at 4°C for 24 hours. After that, spraying was carried out using a spray dryer (Buchi) until all of it became powder.

### 2.2.4 Fortification Nano Whey Protein-Meniran Phyllanthine Encapsulated Casein Hydrolysates in Skimmed Milk Powder

Fortification of skimmed milk powder can be designed by following different concentrations of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates consisting of T0

(0%), T1 (2%), T2 (4%), T3 (6%) and T4 (8%).

### 2.2.5 Solubility

Solubility was analyzed according to [13] with slight modification. Filter paper (Whatman No.1) and centrifuge tube were oven at 105°C for 1 hour, and then removed to desiccator for 15 minutes. The filter paper (A) and centrifugation tube (B) were weighed. A sample of as much as 6 mL was taken and stirred for 30 minutes, centrifuged later for 10 minutes at 400 rpm to separate supernatant and precipitate. The supernatant was filtered using filter paper (Whatman No.1). After that, filter paper with solid sample and centrifuge tube containing precipitate were placed into the oven for 24 hours and desiccated for 15 minutes. The filter paper with solid sample (A') and centrifugation tube with precipitate (B') was weighed and analyzed using the formula :

$$\text{Solubility (\%)} = \frac{A' - A}{(A' - A) + (B' - B)} \times 100\%$$

Description :

A = Filter paper

B = Centrifugation tube

A' = Filter paper + Solid sample

B' = Centrifuge tube + Precipitate

### 2.2.6 Turbidity

Turbidity was determined using [14,15] method. Sample was taken as much as 3 mL and allowed stand for 2 hours. The analysis was carried out using UV-Vis spectrophotometer with a wavelength of 600 nm. The result was observed and recorded.

### 2.2.7 Sedimentation

Sedimentation was described by [16]. Filter paper (Whatman No.1) and centrifuge tube were oven at 105°C for 1 hour, and then removed to desiccator for 15 minutes. The filter paper (A) was weighed. Sample as much as 5 mL was taken and filtered using filter paper (Whatman No.1). The

filter paper with solid sample was weighed and analyzed using formula :

$$\text{Sedimentation} = \frac{B - A \text{ (gram)}}{\text{mL sample}}$$

Description :

A = Filter paper

B = Centrifugation tube

### 2.2.8 Water Activity (aw)

The measurement water activity based on [17] research using aw meter (Rotronic® Instruments) which has been calibrated with  $\text{BaCl}_2$  /  $\text{Mg}(\text{NO}_3)_2$  /  $\text{NaCl}$  /  $\text{KCl}$ . Set the sample in sample container and waited until aw meter rang. The water activity value shown in digital display, then recorded and analyzed.

### 2.2.9 Hygroscopicity

Hygroscopicity analysis was carried out [18] method. The sample was weighed about 1 gram and placed in petri dish, then analyzed the water content's sample. Put the analyzed sample into desiccator which containing saturated solution of  $\text{Na}_2\text{SO}_4$  and allowed stand for 7 days. Hygroscopicity percentage calculated using formula:

$$\text{Hygroscopicity (\%)} = \frac{\frac{x}{ax} - Wi}{1 - \frac{x}{ax}} \times 100\%$$

Description :

X = The increase in weight of sample

Ax = The amount sample

Wi = Water content of sample before exposure humid environment (%)

### 2.2.10 Data Analysis

The data of solubility, turbidity, sedimentation, water activity (aw), and hygroscopicity were analyzed by analysis of variance (ANOVA) was carried out using the Microsoft Excel Software. The level of significance was determined at  $P < 0.05$ .

## 3. RESULTS AND DISCUSSION

### 3.1 Solubility

The research showed that fortification nano whey protein-meniran phyllanthine encapsulated casein hydrolysates had no significant differences ( $P > 0.05$ ) on the solubility of skimmed milk powder. The lowest solubility was obtained on T3 (6% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates) at 57.983% and the highest on T2 (4% nano whey protein-meniran phyllanthine encapsulated casein hydrolysates) at 75.295%. The difference of water-binding ability on the hydrophilic side from whey protein and casein causes an increase or decrease in solubility. Solubility is based on protein's hydrophilic and hydrophobic sides [19].

Solubility percentage reduction on T1 was 68.480% to 61.854% on T2. It happened because of the non-covalent interaction of phenolic compounds from meniran and amino acids from whey protein and casein. Non-polar interaction on polyphenol with protein increases the surface hydrophobicity of protein molecules by reducing solubility properties [20]. Casein hydrolysates utilization as wall material that processed using spray drying method is one of the encapsulation materials with heat tolerance, however heating process up to  $120^\circ\text{C}$  casein gradually becomes insoluble and form precipitate [21]. Insoluble particles cause turbidity and, if continuous, will produce precipitate at the bottom solutions or form sediment.

### 3.2 Turbidity

Turbidity with different concentrations of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates in Table 1 did not significantly affect ( $P > 0.05$ ). Therefore, the product's material composition was an indication of determining the turbidity value of skimmed milk powder. T1 without nano whey protein-meniran phyllanthine encapsulated casein hydrolysates has a turbidity value of 0.482. Bioactive meniran, which interacts with milk protein (whey protein and casein),

provides partial conformational change to protein structure which can change the physicochemical properties of the protein. The processing is one of the causes of turbidity, the higher temperature during the heating process impact lower water content. The lower water content indicated skimmed milk powder products were more hygroscopic. Thus, the particles combine to form larger and insoluble particles causing turbidity. However, in research by [22], increasing particle size resulted in reduced turbidity value. Therefore, the particle size only can not be used indicator of turbidity.

Skimmed milk powder fortified nano whey protein-meniran phyllanthine encapsulated casein hydrolysates on T1-T2 decrease turbidity with an average 0.518-0.459. Whey proteins are proteins that have been isolated and filtered to be easily absorbed by the body. Casein hydrolysates have been involved in hydrolysis, which partly changes protein to amino acids. This amino acids group is more soluble in water and did not give solution turbidity. The addition of phenolic compounds from meniran is not involved in turbidity, but it will be oxidized to protein. The excessive polyphenol concentration causes the protein not to provide cross-linking sites, so the addition phenolic compounds must match [23].

### 2.3 Sedimentation

Sedimentation is physical property to identify the quality of skimmed milk powder. The decreased solubility of skimmed milk powder presents poor solubility, which results in high sedimentation. The different concentrations of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates had no significant differences ( $P>0.05$ ) on sedimentation of skimmed milk powder, shown in Table 1. Skimmed milk powder without nano whey protein-meniran phyllanthine encapsulated casein hydrolysates had the lowest sedimentation value at 0.036 because milk powder contains fat, water, lactose, protein and mineral that have to affect milk sedimentation properties [24–26].

The differences concentration of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates in skimmed milk powder resulted in sedimentation value on T1 (2%), T2 (4%), T3 (6%) and T4 (8%) consecutive at 0.041, 0.042, 0.043, and 0.043. The encapsulation process with casein hydrolysates protected core materials (whey protein and meniran phyllanthine) during spray drying. Casein hydrolysates are assumed can stabilize protein by reducing the tendency of protein particles to combine or form sediment. The fortification nano whey protein-meniran phyllanthine encapsulated casein hydrolysates did not significantly different of skimmed milk powder sedimentation value. However, there have been increased sedimentation trends.

### 2.4 Water Activity (aw)

The result of fortification nano whey protein-meniran phyllanthine encapsulated casein hydrolysates with different concentrations showed a significant difference ( $P<0.05$ ) on skimmed milk powder water activity (Table. 1). The highest to lowest average value of water activity was T3 ( $0.467^b \pm 0.001$ ), T2 ( $0.463^b \pm 0.0006$ ), T4 ( $0.462^b \pm 0.0071$ ), T1 ( $0.460^b \pm 0.0032$ ), T0 ( $0.434^a \pm 0.001$ ). It can be due to covalent bonds from carboxyl groups of amino acids reacting with other amino acid groups by releasing water molecules through hydrogen bridges. Chemically, an amino acid has one carboxyl group and one hydrogen atom, so it will play a role in determining water activity value.

The water activity values in this study were 0.434-0.467, which can not be used for microorganism growth. Microorganism growth in food products identified according to free water content inside. The more free water in food products means the higher water activity. Water activity value to the growth of microorganisms such as bacteria: 0.90; yeast: 0.80-0.90; mold: 0.60-0.70 [27].

Water activity showed a tendency of increasing value on T0 to T3. Processing with drying method using

spray dryer aims to produced low water activity product. According to [28,29] research, the spray drying method is a drying method by hot air to reduce significant water content. The higher temperature and longer time processing impact the lower water activity. The

reduction of water activity (aw) on T4 is thought to be due to the addition of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates was too much. It is known that the functional properties of protein are as a water binder.

**Table I. Physicochemical Analysis of Fortification Skimmed Milk Powder with Nano Whey Protein-Meniran Phyllanthine Encapsulated Casein Hydrolysates**

Treatment	Analysis				
	Solubility (%)	Turbidity	Sedimentation	Water Activity (aw)	Hygroscopicity (%)
T0	68.480±13.835	0.482±0.0029	0.036±0.0049	0.434 <sup>a</sup> ±0.001	4.933±0.275
T1	61.854±9.864	0.518±0.0684	0.041±0.0079	0.460 <sup>b</sup> ±0.0032	4.896±0.407
T2	75.295±17.554	0.489±0.0171	0.042±0.0037	0.463 <sup>b</sup> ±0.0006	4.601±0.669
T3	57.983±8.247	0.478±0.0064	0.043±0.0039	0.467 <sup>b</sup> ±0.001	5.376±0.412
T4	59.179±6.411	0.459±0.0078	0.043±0.0053	0.462 <sup>b</sup> ±0.0071	5.528±0.145

## 2.5 Hygroscopicity

The fortification of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates had no effect ( $P>0.05$ ) in hygroscopicity of skimmed milk powder with values of T0 (4.933%), T1 (4.896%), T2 (4.601%), T3 (5.376%) and T4 ( 5.528%). The protein content of skimmed milk powder is related to their water content. The higher protein content means drier products [30]. The production of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates was carried out through a spray drying process by the rapid evaporation into dried powder. The lower water content will indicate that the product is more hygroscopic. Increasing the hygroscopicity of skimmed milk powder is related to free water content (water activity/aw), which triggers the growth of microorganisms and then impacts the physicochemical properties.

The increased value of hygroscopicity is evidenced by more addition of nano whey protein-meniran phyllanthine encapsulated casein hydrolysates on T2 (4%), T3 (6%) and T4 (8%), resulting in 4,601%, 5,376% and 5,528%. However, some characteristics of protein encapsulants (especially casein hydrolysates) in the food industry must be limited because of their hygroscopic, hydrophobic, reactive and common

allergy triggers [31]. Encapsulation using protein (whey and casein) fortified in skimmed milk powder impact to hygroscopicity reduction, showed on T2 (2%) and T2 (4%) at 4,896% and 4,601%. This may be caused by interaction between polyphenols from meniran with protein and then the change of protein structure and functional properties during encapsulation using the spray drying method. The core material in this study was whey protein and meniran that cause an interaction through hydrophobic bonds. The hydrophobic side of polyphenols binds whey protein, so if casein hydrolysates were added, these hydrophobic bonds were weakened eventually, which reduces hygroscopic of skimmed milk powder. Casein has low hygroscopic properties, therefore unable to absorb moisture from the air again. [32] reported that encapsulation of *curcumin* in casein micelle powder produced by spray-drying impacted its hygroscopicity, which their less hydrophobic bonds because curcumin bind casein.

## 4. CONCLUSION

This research showed that fortification nano whey protein-meniran phyllanthine

encapsulated casein hydrolysates 2% can improve physicochemical properties of skimmed milk powder. Based on this research, which resulted solubility at 61.854%, turbidity 0.518, sedimentation 0.041, water activity (aw) 0.460, and hygroscopicity 4.896%.

#### **COMPETING INTERESTS DISCLAIMER:**

**AUTHORS HAVE DECLARED THAT NO COMPETING INTERESTS EXIST. THE PRODUCTS USED FOR THIS RESEARCH ARE COMMONLY AND PREDOMINANTLY USE PRODUCTS IN OUR AREA OF RESEARCH AND COUNTRY. THERE IS ABSOLUTELY NO CONFLICT OF INTEREST BETWEEN THE AUTHORS AND PRODUCERS OF THE PRODUCTS BECAUSE WE DO NOT INTEND TO USE THESE PRODUCTS AS AN AVENUE FOR ANY LITIGATION BUT FOR THE ADVANCEMENT OF KNOWLEDGE. ALSO, THE RESEARCH WAS NOT FUNDED BY THE PRODUCING COMPANY RATHER IT WAS FUNDED BY PERSONAL EFFORTS OF THE AUTHORS.**

#### **REFERENCES**

1. Call K, Sollohub KC. Spray Drying Technique. I: Hardware and Process Parameters. J. Pharm Sci. 2010;99(2):575-586.
2. Organization WH. The World Health Report 2006: Working Together for Health. 2006; [https://www.google.com/books?hl=en&lr=&id=taYsDwAAQBAJ&oi=fnd&pg=PR13&dq=WHO,+2006&ots=9fk8Zoziz4&sig=98xGy6mi7YqsxKz2t\\_PfHrn0bK0](https://www.google.com/books?hl=en&lr=&id=taYsDwAAQBAJ&oi=fnd&pg=PR13&dq=WHO,+2006&ots=9fk8Zoziz4&sig=98xGy6mi7YqsxKz2t_PfHrn0bK0)
3. Oktavidiati E, Chozin MA, Wijayanto N, Ghulamahdi M, Darusman LK. Plant Growth and Total Phyllanthin and Hypophyllanthin Contents of *Phyllanthus* sp. L Accession on Various Shading Levels. J Littri. 2011;17(1):25-31.
4. Tambunan RM, Swandiny GF, Zaidan S. Antioxidant Activity Test of 70% Ethanol Extract Herbs Meniran (*Phyllanthus niruri* L.) Standardized. J Ilmu Kefarmasian. 2019;12(2):60-64.
5. Shakinaz AES, Refaat AA, Sheltawy STE. Production of Biodiesel using Microwaves Technique. J Adv Res. 2010;1:309-314.
6. Harna C, Kusharto CM, Roosita K. Intervention of High-Protein Milk on Consumption Level of Macronutrient and Nutritional Status in Adult Age Group. J MKMI. 2017;13(4):354-361.
7. Manab A, Sawitri ME, Awwaly KUA. Edible Whey Protein Film: Addition of Egg Lysozyme and Application in Cheese. UB Press; 2017.
8. Saragih WF, Manab A, Sawitri ME, Rahayu PP, Andriani RD. The Effect of Meniran Extract (*Phyllanthus niruri* L.) Addition on Viscosity, Foam Overrun, Foam Stability and Foam's Microscopic of Nano Whey Protein Isolate. In: E3S Web of Conferences. ; 2022:7.
9. Yaakub NM, Zaini HM, Xiang LJ. Effect of Spray Drying Temperature Differences on the Gross Nutritional Composition and Solubility and Sinkability of Goat Milk Powder. J Bangladesh Agric Univ. 2019;17(2):206-210.
10. Diak OA, Bani-Jaber A, Amro B, Jones D, Andrews GP. The Manufacture and Characterization of Casein Films as Novel Tablet Coatings. J Food Bioprod Process. 2007;85(3):284-290.
11. Kusumaningtyas E. The Role OF Milk Peptide As Antimicrobial Agent In Supporting Health Status. WARTAZOA. 2013;23(2):63-75.
12. Rahayu PP, Purwadi, Radiati LE, Manab A. Physico Chemical Properties of Whey Protein and Gelatine Biopolymer Using Tea Leaf Extract as Crosslink Materials. Current Research in Nutrition and Food Science Journal. Curr Res Nutr Food Sci J. 2015;3(3):224-

- 236.
13. Haque E, Chand R. Antihypertensive and Antimicrobial Bioactive Peptides of Milk Proteins. *Eur Food Res Technol.* 2008;227:7–15.
14. Liu Z, Qin L. Research on Turbidity and Surface Hydrophobicity of Milk Protein at Different Conditions. *J Phys Conf Ser.* 2021;1893:1-8.
15. Sikand V, Tong PS, Vink S, Roy S. Physicochemical Properties of Skim Milk Powders Prepared with The Addition of Mineral Chelators. *J Dairy Sci.* 2016;99:4146-4153.
16. Airoldi L, Beck MW. Loss, Status and Trends for Coastal Marine Habitats of Europe. *Oceanogr Mar Biol An Annu Rev.* 2007;45:345-405.
17. Ramadhani PD, Setiani BE, Rizqianti H. The Quality of Avocado Jam (*Persea americana* Mill) with Various Natural Sweetener Flavouring. *J Teknol Pangan.* 2017;1(1):8-15.
18. Tontul I, Topuz A, Ozkan C, Karacan M. Effect of Vegetable Proteins on Physical Characteristics of Spray-Dried Tomato Powders. *Food Sci Technol Int.* Published online 2016:1-9.
19. Al-Awwaly KU. Animal Food Proteins and Its Applications. UB Press; 2017.
20. El-Messery TM, Mwafy EA, Mostafa AM, El-Din HMF, Mwafy A, Amarowicz R, et al. Spectroscopic Studies of the Interaction Between Isolated Polyphenols from Coffee and the Milk Proteins. *Surf Interfaces.* 2020;20:100558.
21. Walstra P, Geurts TJ, Noomen A, Jellema A, van Boekel MAJS. Dairy Technology. In: *Principles of Milk Properties and Processes.* Marcel Dekker; 1999.
22. Siebert KJ, Lynn PY. Effects of Alcohol and pH on Protein–Polyphenol Haze Intensity and Particle Size. *J Am Soc Brew Chem.* 2018;61(2):88-98.
23. Wang Y, Ye L. Haze in Beer: Its Formation and Alleviating Strategies, from a Protein–Polyphenol Complex Angle. *Foods.* 2021;10(3114):1-11.
24. Rusdi R, Sidi TBP, Pratama R. The Effect of Concentration and Deposition Time of Moringa Seeds pH, Turbidity and Color of Krengeng Reservoir Water. *J Integr Proses.* 2014;5(1):46-50.
25. Le Tran, Saveyn TP, Hoa HD, van der Meeren P. Determination of Heat Induced Effects on the Particle Size Distribution of Casein Micelles by Dynamic Light Scattering and Nanoparticle Tracking Analysis. *Int Dairy J.* 2008;18(2):1090–1096.
26. Gaur, Schalk VJ, Anema SG. Sedimentation in UHT milk. *Int Dairy J.* 2018;78:92-102.
27. Belitz HD, Grosch W, Schieberle P. *Springer Food chemistry 4th Revised and Extended Edition.* Annu Rev Biochem. 2009;79:655-681.
28. Dwika RT, Ceningasih T, Sasongko SB. The Effect of Dryer Temperature and Air Flow Rate on Carrageenan Drying Using Spray Dryer Technology. *J Teknol Kim dan Ind.* 2012;1(1):298-304.
29. Leviana W, Paramita V. Effect of Temperature on Water Content And Water Activity In Material On Turmeric (*Curcuma Longa*) With Tools Electrical Oven Dryer. *Metana.* 2017;13(2):37-44.
30. Puspita DA, Agustini TW, Purnamayanti L. The Effect of Different Concentration of Salt Towards The Glutamic Acid in Catfish Bakasang Powder. *J Teknol Pangan.* 2019;3(1):110-115.
31. Ortiz SEM, Mauri A, Monterrey-Quintero ES, Trindade MA, Santana AS, Favaro-Trindade CS. Production and Properties of Casein Hydrolysate Microencapsulated by Spray Drying with Soybean Protein Isolate. *LWT-Food Sci Technol.* 2009;42:919-923.
32. Khanji AN, Michaux F, Petit J, Salameh D, Rizk T, Jasniowski J, et al. Structure, Gelation, and Antioxidant Properties of Curcumin-Doped Casein Micelle Powder Produced by Spray-Drying. *Food Funct.* 2017;9(2):971-981.



UNDER PEER REVIEW