

# Original Research Article

## Modelling Water Absorption Kinetics of *Abrohema*, *Omankwa*, and *Akposoe* Maize Varieties

### ABSTRACT

The study investigated the effect of temperature and variety on water absorption kinetics of three Ghanaian maize varieties (*Abrohema*, *Omankwa*, and *Akposoe*) during soaking and also to generated the required moisture diffusivity data for the varieties. The study employed a gravimetric method at four different temperatures of 30°C, 40°C, 50°C, and 60°C. Varying soaking temperature affected the water absorption behaviours and rates of the varieties. The Fick's diffusion law satisfactorily predicted the absorption kinetics of the varieties at all temperatures and that variety and temperature were the most important factors controlling water absorption rate of the maize kernel. The moisture diffusivities of the kernels varied in the order of *Abrohema* > *Omankwa* > *Akposoe* and increased as the soaking temperature increases with values ranging from 2.91- 3.93 x 10<sup>-10</sup> m<sup>2</sup>/s for *Akposoe*, 3.19 - 4.33 x 10<sup>-10</sup> m<sup>2</sup>/s for *Omankwa* and 3.66–4.74 x 10<sup>-10</sup> for *Abrohema*. The Arrhenius equation was able to described the effects of temperature on the diffusion coefficient with activation energy values of 7.04 for *Abrohema* and 8.31 kJ/mol for both *Akposoe* and *Omankwa*

**KEYWORDS:** Modelling, Water absorption, Kinetics, Maize, Soaking

### 1 Introduction

Maize (*Zea mays L.*) is one of the many cereal crops grown and marketed in many parts of the world and plays major economic role in Sub-Saharan Africa (SSA)[1]. It is regarded as dominant cereal and staple crop produced and consumed across Africa regions and in many parts of America. Govaerts [2][3] put the daily consumption per capita as 290g in Guatemala and Mexico while consumption per capita in Venezuela and Colombia was estimated as 100g/day. The Maize kernel serves as inexpensive source of protein and cheap energy and therefore becomes a good replacements or add-on to the major food staples. FAOSTAT [4] maintained that maize constitute 40% of the total cereal production and 80% of the total production is consumed as food in the SSA regions. It is also believed that maize could become the food for the next generation and [3] argued that the consumption of maize in the Sub-Saharan regions is likely to increase threefold by the year 2050. Nuss et al [5] observed that Sub Saharan Africans derived 30% of the energy intake from maize. Cereals contribute immensely to agricultural production and help to provide the needs of Africa's growing population. This accession indicates, that the consumption of maize will remain extremely important for years to come. In Ghana, more than 1 million (20%) of estimated 5 million small scale farming households gain main income from the production of maize [6]. In 2018/2019 growing season, Ghana achieved a production level of 1.09 billion metric tons in maize.

Maize processing often requires rehydration of the kernels to make it easy for further operations such as milling, cooking and also to achieve desired palatability. Thus, the knowledge of absorption of water by the kernels is extremely important to the food industries [7].

The rate and amount of moisture imbibed by kernels during rehydration is determined by numerous factors including the initial moisture content, thickness of outer bran layer, variety of kernel, kernel size, soaking duration as well as temperature and soaking water acidity [8]. Studies have shown that the rate at which water is absorbed by kernels is greatly influenced by temperature and soaking period [9]. A rise in soaking temperature produces higher water absorption rate [10][11].

From a processing and engineering prospective, food processors are not only interested in determining how fast water absorption can be achieved, but also, how processing parameters including temperature affects the process [7][12] as well as how soaking time could be predicted at a specific processing condition. The models obtained from Fick's laws of diffusion and the Peleg model are widely employed to characterize the soaking of cereals and legumes kernels [13]. Modelling the absorption kinetics during maize soaking is important since it offers industry players the opportunity to manipulate the operation to optimize industrial processing of maize-based dishes.

Hence, availability of quantitative data on processing parameters are necessary for optimization of the soaking conditions, design of food processing equipment as well as predicting water absorption behaviour of maize kernels. Nevertheless, the available literature provides little or no data on the moisture absorption kinetics of *Abrohemaa*, *Omankwa*, and *Akposoe* maize varieties grown in Ghana. Consequently, the current study was designed to investigate the temperature and variety effects on these varieties during soaking and also to generate the required moisture diffusivity data for these three Ghanaian maize varieties.

## 2 Materials and Methods

### 2.1 Collection and preparation of maize kernels

Three maize varieties (*Abrohemaa*, *Omankwa*, and *Akposoe*) were obtained from the experimental farm of Crops Research Institute (CRI), Fumasua, Kumasi. The kernels of the varieties were prepared manually and only whole and good kernels were selected for the experiment. Each sample was packed separately into zip-lock polythene bags and placed in a refrigerator to prevent moisture loss or gain.

### 2.2 Estimation of initial moisture content

The initial moisture content of the kernels were estimated by the standard oven drying procedure described by [14]. The three varieties of sample size 5g was weighed using Electronic Compact Scale, SF-400C and placed in labelled dishes. The dishes, together with the samples were placed in standard oven and uniformly heated to a temperature of  $103^{\circ}\text{C} \pm 2^{\circ}\text{C}$  for 4h. The weight loss of the sample was monitored with the electronic Compact Scale, until subsequent weight loss was less than 0.01g. The weight loss by the sample was taken as the weight of water in the sample before drying. The procedure was triplicated and the average value was estimated.

The initial moisture content ( $M_d$ ) on dry basis was then computed as follows:

$$M_d = \frac{W_w}{W_d} \times 100\% \quad (1)$$

$$W_d = W_t - W_w$$

Where:

$W_t$  = weight of sample before drying,

$W_d$  = weight of sample after drying (dry matter)

$W_w$  = weight of water loss

### ***2.3 Measurement of length, width and thickness of kernel***

The length, width and thickness of the kernel was determined by the method described by [15]. Hundred (100) kernels were picked at random and their principal diameters were measured along three directions with a digital micrometer screw gauge with accuracy of 0.01 mm. The major diameter was equated to the length of the kernel, the intermediate diameter equated to the width, and the minor diameter equated the thickness of the kernel. The micrometer screw gauge was held at right angle to the direction of the diameter being measured. Hundred kernels were selected and used to determine the average value of the length while 50 kernels were used for both width and thickness as proposed by [16]. For the sake of irregularity and non-uniformity of the shape of the kernels, the highest values of the dimensions were recorded and used.

### ***2.4 Estimation of equivalent radius (r)***

The equivalent radius was evaluated from the average volume of the kernels. The volume of a kernel was calculated based on the assumption that the volume of the kernel can be approximated to the volume of a spherical object.

The volume of the kernel was determined by filling a 100-ml measuring cylinder with 50 ml of water. Then, 50 kernels were immersed in the water. The amount of water displaced was recorded. The procedure was replicated and the true volume of a kernel was calculated.

The average volume of a kernel was equated to the volume of a sphere ( $V = \frac{4}{3} \pi r^3$ ) and equivalent radius ( $r$ ) was obtained.

Thus,

$$r = \sqrt[3]{\frac{3v}{4\pi}} \quad (2)$$

### ***2.5 Determination of saturation moisture content (Ms)***

Three set of sample size  $5g \pm 0.0$  for each variety was soaked at four different water temperatures in a portable water bath fixed with temperature control system. The samples were situated separately in a labelled nylon mosquito net and placed into the water bath at a pre-determined temperature of  $30^\circ\text{C} \pm 2^\circ\text{C}$ ,  $40^\circ\text{C} \pm 2^\circ\text{C}$ ,  $50^\circ\text{C} \pm 2^\circ\text{C}$ , and  $60^\circ\text{C} \pm 2^\circ\text{C}$ . The sample was removed at time interval of 1h, and quickly wiped out the residual water on the surface of the kernels [17] and then reweigh [18]. The weight gain of the sample was monitored at each time interval using the electronic Compact Scale. The moisture content of the sample at specific time interval was

calculated and recorded. The saturation moisture content ( $M_s$ ) was computed when the subsequent weight gain of the sample was  $\leq 0.01$ g.

## 2.6 Modelling of absorption kinetics

At constant temperature, it is assumed that diffusion process follows Fick's second law of diffusion. Therefore, Fick's three-dimensional equation can be applied for an axisymmetric diffusion:

$$\frac{\partial M}{\partial t} = D \left( \frac{\partial^2 M}{\partial x^2} + \frac{\partial^2 M}{\partial y^2} + \frac{\partial^2 M}{\partial z^2} \right) \quad (3)$$

Where,  $D$  is the diffusion coefficient and  $M$  is the water content at time  $t$ . The equation (3) was re-arranged to obtain a solution for an object with a sphere shape (Seyhan-Curtas *et al.*, 2001),

$$MR = \frac{M - M_i}{M_e - M_i} = 1 - \left( \frac{6}{\pi^2} \right) \sum_{i=1}^{\infty} \left( \frac{1}{i^2} \right) \exp(-Di^2 \pi^2 \frac{t}{r^2}) \quad (4)$$

where,

MR = moisture ratio,

$M_i$  = initial moisture content,

$M_e$  = equilibrium moisture contents.

$Di$  = effective diffusion coefficient,

$i$  = number of terms in the summation

$r$  = equivalent radius of the kernels and

$t$  = soaking time.

In this case, only a finite number of Eq. (4) was used for estimating the MR values. All moisture terms were computed on dry basis.

The effective diffusion coefficient,  $Di$  was evaluated from the slope of a curve,  $1-Mr$  against the soaking time  $t$ .

The temperature dependency of diffusion coefficient ( $D_i$ ) was evaluated using an Arrhenius equation:

$$Di = D_o e^{-E_a/RT} \quad (5)$$

where:

$D_o$  = diffusion constant ( $m^2/s$ ),

$E_a$  = activation energy (kJ/mol),

$R$  = general gas constant (8.314 J/mol K) and

$T$  = absolute temperature (K).

The  $E_a$  values for the selected varieties were calculated from the linear regression of  $D_i$  against reciprocal of soaking time ( $1/T$ ). The constant  $D_0$  and the slope  $E_a/R$  were computed. The  $E_a$  values were obtained by multiplying the gas constant (8.314 J/mol K) by the value of the slope.

### 3 RESULTS AND DISCUSSION

#### 3.1 Moisture content and dimensions

The initial moisture contents ( $M_i$ ), length (L), width (W), thickness (T), equivalent radius (r) and surface area (SA) of the kernels are presented in **Table 1**.


**Table 1:** Initial Moisture Content and Physical Properties of Maize Varieties

Maize	$M_i$ (%db)	L(mm)	W(mm)	T(mm)	r(mm)	SA(mm <sup>2</sup> )
<i>Aborohema</i>	11.76	11.03	8.98	4.74	4.04	596.65
<i>Omankwa</i>	11.76	10.47	9.02	4.53	4.02	560.79
<i>Akposoe</i>	13.66	10.20	8.80	4.10	3.65	507.23


**Table 1** shows values for length (l), width (W), thickness (T) and surface area (SA) of the kernels used in this study. The SA ranged from 507.23 mm<sup>2</sup> to 596.65 mm<sup>2</sup>. These values are in agreement with values documented by [19]. In general, for different varieties one would anticipate an inverse relation between the absorption rate and kernel size, since a larger kernel yields a smaller specific surface area for moisture movement [20].

#### 3.2 Saturation moisture content and diffusion coefficient


The water absorption curves of maize varieties soaked at temperatures are illustrated in **Figure 1a, 1b and 1c**.



**Figure 1a: Water absorption curves of *Abrohema***



**Figure 1b: Water absorption curves of *Omankwa***



**Figure 1c: Water absorption curves of *Akposoe***

The maize varieties showed moisture absorption behaviour similar to what has been published. The initial water absorption was rapid, immediately followed by a slower rate and at later stages asymptotically got to the saturation point ( $M_s$ ). Moisture saturation was reached within 12h of soaking at temperature of 30°C but soaking time was shortened to 8h as soaking temperature rises to 60°C. This observation was in line with reports by other published studies [21].

The rate at which water was absorbed by these kernels were found to increase with increasing soaking temperature as a result of increased diffusion rates. Also, high water temperature could cause damage to cell tissues leading to an increased opening of the pores which will consequently increase the rate of water absorption. Identical findings have been documented by

[22] [23][24][25][26]. Reports by other researchers such as [17][27] derived similar hydration curves for red kidney beans, selected Turkish legumes, wheat and barley.

The study also found that water absorption capacity increased as the soaking temperature increases and that the rates of moisture increment differ for the varieties considered. The observed phenomenon could be attributed to the difference in varietal characteristic such as nutritional composition of the varieties. This confirms earlier reports on the water absorption behaviour of rice by [28][29].

The study further observed that Akposoe had moderately higher water gain than the other varieties. At temperatures of 30°C and 40°C, *Abrohema* recorded the least water gain as anticipated while *Omankwa* recorded minimum water gain at 60°C. Generally, water absorption rates were seen to be influenced by temperature. The observed effect of temperature on absorption rates agreed with other published studies [30][17][27]. Shafaei et al. [21] made identical observation after studying water absorption characteristics of chickpea and soybean.

An increased specific surface promotes higher absorption capacity due to smaller nature of kernel size [13]. This phenomenon was noticed in the current study. *Akposoe* had a small kernel size in relation to *Omankwa* and *Abrohema* therefore recorded higher water absorption capacity as expected. In contrast, *Abrohema* kernel was relatively bigger but recorded the upmost absorption capacity at temperatures 50°C and 60°C. This behaviour could be attributed to variations in severity of disorderness brought about by different temperature levels to bear on the nutritional composition. The observed occurrence could suggest a high probability of *Abrohema* containing more starch content than *Akposoe* and *Omankwa*. Addo et al. [7] made similar observation that high starch content of *Obatanpa* was accountable for its higher water absorption capacity although *Mamaba* kernel was relatively smaller than *Obatanpa*. Many reports including Ituen et al. [31] have shown that kernel characteristics and temperature are the most important factors responsible for variations in equilibrium moisture content values. Brennan et al. [32] also made similar assertion that the extent to which protein and starch (water holding constituents) have been destroyed by temperature during drying principally influenced reconstitutability of dried food products.

### 3.3 Water absorption rates of maize Hybrids

The data on moisture ratio obtained from the soaking experimental was plotted against the square root of soaking period. Based on eqn. (4), the effective diffusivity values for the varieties during water absorption process was determined. The moisture contents ( $M_s$ ) employed in eqn. (4) and the coefficient of determination ( $R^2$ ) are indicated in **Table 2**.


**Table 2:** Effective moisture Diffusivity ( $D_i$ ) values and Saturation Moisture Content ( $M_s$ )

Variety	30°C			40°C			50°C			60°C		
	$D_i^*$	$R^2$	$M_s$	$D_i^*$	$R^2$	$M_s$	$D_i^*$	$R^2$	$M_s$	$D_i^*$	$R^2$	$M_s$
<i>Abrohema</i>				3.9	0.9	53.4	4.2	0.9			0.9	73.2
<i>a</i>	3.66	0.99	48.95	6	8	7	1	9	64.5	4.74	8	7
				3.4	0.9	54.0	3.7	0.9	60.2		0.9	66.4
<i>Omankwa</i>	3.19	0.99	50.02	8	9	0	5	7	7	4.33	4	1

				3.1	0.9	61.1	3.3	0.9	70.1		0.9	72.7
<i>Akposoe</i>	2.91	0.97	55.37	3	9	0	8	9	2	3.95	6	4
	* $\times 10^{-10} \text{ m}^2/\text{s}$											


Estimated  $D_i$  values of the selected varieties at various water temperatures are indicated in **Table 2** above. Generally, the experimental data fitted the diffusion equation with high accuracy. The  $R^2$  values obtained from the linear regression analysis, varied from 0.94 to 0.99. The  $D_i$  of *Abrohemaa* varied from  $3.66 - 4.74 \times 10^{-10} \text{ m}^2/\text{s}$ ; *Omankwa*,  $3.19-4.33 \times 10^{-10} \text{ m}^2/\text{s}$  and *Akposoe* varied  $2.91-3.95 \times 10^{-10} \text{ m}^2/\text{s}$ . The observed differences in  $D_i$  could be attributed to differences in kernel characteristics due to varietal variations of the maize. Agarry et al [33] attributed the kernel's propensity to absorb water to the two main constituent (protein and carbohydrate) with protein possessing a higher capacity to absorb water than carbohydrate.

Therefore, variations in nutritional characteristics of the kernels could provide another explanation for *Abrohemaa* variety having higher diffusion coefficient than *Omankwa*, and *Akposoe* varieties. It is also probable that *Abrohemaa* kernel contain high protein concentrations than *Omankwa* and *Akposoe*, causing it to soak up more water than the two varieties within the same temperature and time conditions. In comparison with other studies, the  $D_f$  values obtained in this work is marginally higher than those documented by [7] for *Mamaba* and *Obatanpa* but were lower than values reported by [8]. The observation could possibly be attributed to variations in the physiochemical and nutritional characteristics of the varieties studied. Addo et al. [7] and Agidi et al. [34] agreed to this assertion when they reported that seed's size and nutrient are primarily responsible for water uptake in seeds soaking. Additionally, Webb [35] also disclosed that among the starch, amylose content has been found to be the most well know property for forecasting rate of moisture gain. The rates of water absorption of the selected varieties are presented in **Fig. 2a, 2b and 2c**.




**Fig. 2a: Water absorption rate for *Abrohemaa***






**Fig. 2b: Water absorption rate for *Omankwa***



**Fig. 2c: Water absorption rate for *Akposoe***

### 3.4 Activation Energies ( $E_a$ ) of selected maize varieties

The values of diffusion coefficients of the selected maize was fitted into an Arrhenius-type relationship (eqn. (5)) in other to relate the temperature effect to moisture diffusivity of the maize kernels. A plot of  $D_f$  versus reciprocal of absolute temperature are presented in **Figure 3**. The  $R^2$  obtained varied from 0.96 to 0.98 which represents a good fit of the experimental data.



**Figure 3:** The relationship between water Diffusivity and reciprocal of temperature

**Figure 3** portrays the correlation between temperature and the diffusion coefficients of the varieties (*Akposoe*, *Abrohemaa* and *Omankwa*). The graphs indicates that the Arrhenius equation was able to describe the strong effects of temperature on the water absorption behaviour of the maize varieties. The  $E_a$  obtained for the selected Maize varieties are presented in **Table 3**

**Table 3:** Estimated values of Activation Energy ( $E_a$ ) of selected maize varieties

Variety	$E_a$ (kJ/mol)	$R^2$
<i>Akposoe</i>	8.31	0.96
<i>Abrohemaa</i>	7.04	0.98
<i>Omankwa</i>	8.31	0.98

The  $E_a$  estimated from the slopes of the curves in **figure 3** shows  $E_a$  values of 8.31 kJ/mol for *Akposoe*; 7.04 kJ/mol for *Abrohemaa* and 8.31 kJ/mol for *Omankwa*. The marginally low energy values obtained in this study suggest that the varieties were less sensitive to heat and that there was a heat build-up during the absorption process. In comparison,  $E_a$  values obtained in this study were identical to values reported by Addo et al. (2006) for *Obatanpa* (6.54kJ/mol) and *Mamaba* (6.82kJ/mol). Idun-Acquah et al. [36] obtained similar  $E_a$  values for *Asontem* (7.27 kJ/mol), *Hewale* (7.26 kJ/mol) and *Asomdwee* (6.26 kJ/mol). In contrast, Verma and Prasad [12]; Charan and Prasad [37] reported large values of activation energies in the range of 33.5-41.56 and 45.75 respectively for certain varieties of maize. These differences in activation energies could be attributed to varietal variations resulting from physiochemical and nutrient composition of the maize varieties.

#### 4. Conclusion

The water absorption kinetics of three maize varieties, *Akposoe*, *Abrohemaa* and *Omankwa* were studied using phenomenological models deduced from Fick's law of diffusion. The study found

that temperature and variety significantly influenced the water absorption rate and capacity of the kernels. The soaking duration could be shortened by raising the water temperature from 30°C to 60°C. Variations in rate of water absorption could be attributed to kernel characteristics such as physiochemical and nutritional composition of the varieties. The Fick's diffusion law adequately simulated and predicted the water absorption kinetics of the varieties. The effective moisture diffusivity values varied from  $3.66 - 4.74 \times 10^{-10} \text{ m}^2/\text{s}$  for *Abrohema*;  $3.19-4.33 \times 10^{-10} \text{ m}^2/\text{s}$  for *Omankwa*; and  $2.91-3.95 \times 10^{-10} \text{ m}^2/\text{s}$  for *Akposoe*. The temperature dependency of moisture diffusivity of the varieties were satisfactorily represented by the Arrhenius equation with the activation energies for *Abrohema* (8.31 kJ/mol), *Akposoe* (7.04 kJ/mol) and *Omankwa* (8.31kJ/mol).

## REFERENCES

- [1] Badu-Apraku, B. and Fakorede, M.A.B. 2017. Maize in Sub-Saharan Africa: importance and production constraints,” *Advances in Genetic Enhancement of Early and Extra-early Maize for Sub-Saharan Africa*, Cham, Switzerland: Springer International Publishing.
- [2] Govaerts, B.; Chávez, X.; Fernández, A.; Vega, D.; Vázquez, O.; Pérez, M.; Carvajal, A.; Ortega, P.; López, P.; Rodríguez, R.; Kruseman, G.; Donnet, L.; Palacios-Rojas, N. Verhulst, N.; Gardeazabal, A.; González, G.; Sánchez, K. C. and Rosado, L. G. 2019. Maize para México – plan Estratégico 2030. CIMMYT: México. <https://repository.cimmyt.org/xmlui/bitstream/handle/10883/20219/60937.pdf>
- [3] Ekpa, O.; Palacios-Rojas, N.; Kruseman, G.; Fogliano, V.; Linnemann, A. R. 2019 Sub-Saharan African maize-based foods: processing practices, challenges and opportunities Food Rev. Int. 35 (7):609-6039

- [4] **FAOSTAT 2016. Food Balance Sheet, Available:**  
<http://faostat.fao.org/site/345/default.aspx>
- [5] **Nuss, E. T.; Tanumihardjo, S. A. Maize: A. 2010. Paramount Staple Crop in the Context of Global Nutrition. Compr. Rev. Food Sci. Food Saf. 9(4):417–436. DOI: 10.1111/j.15414337.2010.00117x.**
- [6] World Bank. 2008. Maize Value Chain Study & Intervention Analysis in Ghana
- [7] Addo, A.; Bart-Plange, A. and Dzisi, K. 2006. Water Absorption Characteristics of Obatanpa and Mamaba Maize Hybrids (*Zea mays*). *International J. of Food Engineering* 2(3):1556-3758.
- [8] Demirhan, E. and Özbek, B. 2015 Modeling of the Water Uptake Process for Cowpea Seeds (*Vigna unguiculata* L.) under Common Treatment and Microwave Treatment. *Journal of Chemical Society of Pakistan*, 37, 1-10.
- [9] Shafaei, S.M. and Masoumi, A.A. 2014 Estimating Moisture Absorption Kinetics of Beans during Soaking Using Mathematical Models. *CIGR Journal*, 16, 231-237.
- [10] Kashaninejadl, M and Kashiri, M. 2007. Hydration kinetics and changes in some physical properties of wheat kernels. *J. of Iranian Food Science & Tech. Research* 45:58-59
- [11] Sfayhi-Terras, D.; Hadjyahia N. and Zarroug, Y. 2021. Effect of soaking time and temperature on water absorption capacity and dimensional changes of bulgur. *Journal of General Chemistry. Cereal Chem.* 98: 851– 857. <https://doi.org/10.1002/cche.10427>.
- [12] Verma, R. C. and Prasad, S. 1999. Kinetics of absorption of water by maize grains. *Journal of Food Engineering* 39:395–400.
- [13] Tang, J.; Sokhansanj, S. and Sosulski, F. W. 1994. Moisture-absorption characteristics of Laird lentils and hard shell seeds. *Cereal Chemistry* 71(5):423-428.
- [14] AOAC. (2003). *Official methods of analysis* (17<sup>th</sup> ed.). Arlington: Association of Official Analytical Chemists, Washington D.C.
- [15] Baryeh, E. A. 2001. Physical properties of bambara groundnuts. *Journal of Food Engineering* 47:321–326.
- [16] Hauhouot-O'Hare, M.; Criner, B.R.; Brusewitz, G.H. and Solie, J.B. 2000. Selected Physical Characteristics and Aerodynamic Properties of Cheat Seed for separation from Wheat. *CIGR Journal of Scientific Research and Development* 2:347-356.
- [17] Seyhan-Gurtas, F.; Ak, M. M. and Evranuz, E. O. 2001. Water diffusion coefficients of selected legumes grown in Turkey as affected by temperature and variety. *Turkish Journal of Agriculture* 25:297-304
- [18] Lu, R.; Siebenmorgen, T. J. and Archer, T. R. 1994. Absorption of water in long-grain rough rice during soaking. *Journal of Food Process Engineering* 17:141-154.
- [19] Gurses, L. 1981. Bazı Yerli Kurubaklagil Sebzelerinin Kimyasal Bileşimleri ve Bitlerin Konserveye Yaratılan Etkileri üzerine Araştırmalar. Ankara: Ankara Üniversitesi Yayınları, pp. 43-57.
- [20] Hsu, K. H.; Kim, C. J. and Wilson, L. A. 1983. Factors affecting water uptake of soybean during soaking. *Cereal Chemistry* 60:208-211.
- [21] Shafaei, S.M., Masoumi, A.A. and Roshan, H. 2014. Analysis of Water Absorption of Bean and Chickpea during Soaking Using Peleg Model. *Journal of the Saudi Society of Agricultural Sciences*, 15:135-144

- [22] Sopade, P.A.; Ajisegir, E.S, and Okonmah, G.N 1994. Modelling water absorption characteristics of some Nigerian varieties of cowpea during soaking. *Tropical Science* 34:297-305.
- [23] Addo, Ahmad and Bart-Plange, A. 2009. Kinetics of Water Sorption by Egusi Melon (*cucumeropsis edulis*) Seeds. *ARPN J. of Agricultural and Biological Sci.* 4(6):14-17.
- [24] Tunde-Akintunde, T.Y. 2010. Water absorption characteristics of Nigerian Acha (*Digitariaexilis*). *International Journal of Food Engineering*, 6 (5), art 11: 1-10. DOI: 10.2202/1556-3758.1650.
- [25] Agu, H.O, Badau M.H, & Abubakar U.M. 2013. Modeling the water absorption characteristics of various local pearl millet grains and hungry rice (*digitariaexilis*) during soaking using peleg's equation. *Focusing On Modern Food Industry*, 2 (4), 161– 69.
- [26] Udoro E.O.; Kehinde A.T.; Olasunkanmi, S.G.; & Charles, T.A. 2014. Studies on the physicochemical, functional and sensory properties of Gari processed from dried cassava chips. *Journal of Food Process Technology*, 5:293. doi:10.4172/2157-7110.1000293.
- [27] Tagawa, A.; Muramatsu, Y.; Nagasuna, T.; Yano, A.; Iimoto, M. and Murata. S. 2003. Water absorption characteristics of wheat and barley during soaking. *Transactions of the ASAE* 46(2):361-366.
- [28] Shittu, T. A.; Aminu, R. A. and Abulude, E. O. (2009). Functional effects of Xanthan gum on composite cassava–wheat dough and bread. *Food Hydrocolloid* 23:2254–2260.
- [29] Bandyopadhyay, S. and Roy, C.N. 1992. *Rice Processing Technology*. New Delhi: IBTT Publishing.
- [30] Abu-Ghannam, N. 1998. Modelling textural changes during the hydration process of red beans. *Journal of Food Engineering* 38:341–352.
- [31] Ituen, E. U. U.; Mittal, J. P. and Adeoti, J. S. 1985. Water absorption in cereal grains and its effects on their rupture stress. *J. of Food Process Engineering* 8:147-158.
- [32] Brennan, J.G.; Butters, J.R.; Cowell, N.D. and Lilly, A.E.V. 1990. *Dehydration in Food Engineering Operations*. London: Elsevier Applied Science.
- [33] Agarry, S.E.; Afolabi, T.J. and Akintunde, T.T.Y. 2014. Modelling the Water Absorption Characteristics of Different Maize (*Zea Mays L.*) Types during Soaking. *Journal of Food Processing & Technology*, 5:26. <https://doi.org/10.4172/2157-7110.1000326>.
- [34] Agidi, G.; Musa, S.D. and Chukwugoziem, I.J. 2009. Water Absorption Behaviour of Milled Parboiled Rice in Relationship with Variety and Some Basic Thermodynamic Properties of Steam. *AU J.T.* 13(2):101-106.
- [35] Webb, B.D. 1975. *Cooking, processing and milling qualities of rice*. Texas: Texas Agric. Experimental Station.
- [36] Idun-Acquah, N.N.; Addo A. and Bart-Plange, A. 2019. Effect of Soaking Temperature on Water Absorption Characteristics of Selected Ghanaian Cowpea Varieties. *Open Journal of Applied Sciences* 9: 736-748 <https://www.scirp.org/journal/ojapps>.
- [37] Charan, R. and Prasad, S. 1996. Moisture diffusion during hydration of maize. *Journal of Food Science and Technology* 33(5):383-388.