

### **Bioactivity of Leaf Extract of *Phyllanthus amarus* Against Fungal Pathogens Associated with Rot of Sweet Orange.**

#### **Abstract**

This study was carried out to determine disease incidence and control of fungal pathogens associated with rot of sweet orange using *Phyllanthus amarus* leaf extract. Forty sweet oranges were collected from the markets and taken to the laboratory. Isolation and identification of the fungi were carried out using standard procedures and the efficacy of the leaf extracts of *Phyllanthus amarus* was determined on the isolated fungi. The result of this work showed that four pathogens (*Aspergillus niger*, *Aspergillus flavus*, *Penicillium expansum* and *Rhizopus stolonifer*) were identified to be associated with the rot. The Main market had the highest incidence of 35%, the pathogenicity test carried out showed growth with similar characteristic features to the original diseased samples. The effect of the leaf extract of *Phyllanthus amarus* on the isolated fungi shows that the rate of inhibition increases with increase in concentration of the extract, *A. flavus* had the highest zone of inhibition (10.667) while *R. stolonifer* had the least among the pathogens (8.333). Sweet orange should be consumed shortly after harvesting or properly refrigerated and should be discarded if any alteration in color or taste of the fruit is noticed as this can be hazardous to human health.

Keywords: *Phyllanthus amarus* , laboratory, Orange, pathogens, fruit.

#### **INTRODUCTION**

Fruits are very important for the wellbeing of human due to their dietary and nutritional qualities. The consumption of fruit and vegetable products has drastically increased by more than 30% during the past few decades [1]. They are good sources of nutrients for growth, repair and control of body processes as most of them contain sugar, vitamins, mineral elements and small quantities of protein and oil [2]. *Citrus sinensis*, also known as sweet orange, is the most popular of the citrus fruits. It is widely cultivated in most regions of the world [3]. Oranges form a rich source of vitamin C, flavonoids, phenolic compounds and pectins. The main flavonoids found in citrus species are hesperidine, narirutin, naringin and eriocitrin [4]. Just one orange provides 116% of the daily requirement for vitamin C. Vitamin C is the primary water-soluble antioxidant, which prevents free radical generation in the body and damage to the tissues in the aqueous environment both inside and outside cells [5]. Drinking of orange juice without salt and sugar is associated with reduced severity of inflammatory conditions, like asthma, osteo-arthritis, and rheumatoid arthritis. Vitamin C is also necessary for the proper functioning of immune system. Vitamin C is good for preventing cold, cough and recurrent ear infections [6]. Studies have shown that oranges have been found to protect the moderate consumer against cardiovascular diseases [5], possess anti-carcinogenic properties, reduce the risk of kidney stones [7], possess anti-ulcer properties, antianxiety effect, anti-typhoid activity [8], antibacterial activity [5] and antifungal activities [9] amongst the many medicinal uses.

The improper handling, packaging, storage and transportation may result in decay and growth of microorganisms, which become activated because of the changing physiological state of the fruits and vegetables. Fruit, due to their low pH, higher moisture content and nutrient composition are very susceptible to attack by pathogenic fungi, which in addition to causing rots, may also make them unfit for consumption by producing mycotoxins [10]. The principle of spread of fungal infection in fruits supports that a single infected orange fruit can be the source of infection to other orange fruits during storage and on transit [11]. Soil-infesting fungi and bacteria that cause loss of fleshy tissue typically infect plants at the time of or just before harvesting. Infection may occur, however, during post-harvest handling or storage. Common air molds such as *Penicillium* species may gain entry into the susceptible tissue and cause loss during packaging. *Penicillium digitatum* and *Penicillium italicum* causes green and blue mold diseases respectively which are universal post-harvest diseases of citrus. The extensive spore production by these pathogens ensures its presence wherever fruit was handled, including field, packing house, equipment, de-greening, storage rooms, transit containers and market place [12].

Postharvest disease control methods must be environmentally friendly, safer for human health, economically viable, and able to extend shelf life. Biological control by natural compounds appears to be feasible and may present an alternative to synthetic fungicides. Using antagonistic microorganisms, compounds of natural origin and physical measures are some of the effective alternatives for biological control of postharvest diseases. Utilization of natural antimicrobial agents could represent a low-cost alternative therapy as microorganisms become resistant to conventional antimicrobial agents. Chemical studies of antimicrobial plant extracts could reveal new substances with potential usefulness as antimicrobial drugs or as models for the development of new drugs. Microorganism are known to destroy fruits, thereby reducing the quantity for consumption and the profit desirable from the fruit such as sweet orange. Fungal pathogens have become a serious problem in orange, due to this attack fungicides are used to control the pathogens. However, the leaves extract of *Phyllanthus amarus* have been reported to have an antimicrobial effects, it is good to test leaf efficacy thus providing a more alternative to the use of synthetic fungicides. The aim of the study is to determine the fungal pathogen associated with post-harvest rot of sweet orange in Mubi.

## **MATERIALS AND METHODS**

### **Sample collection**

Samples of spoilt Citrus fruit and healthy ones were collected from two (2) markets in Mubi (Mubi main market and Kasuwan Kuturu). The leaves of *Phyllanthus amarus* were obtained from Sabon Gari, Digil ward Mubi North in a sterilized polythene bag and taken to the herbarium unit of the Department of Botany, for identification and authentication.

### **Preparation of culture medium**

Sabourand Dextrose Agar (SDA) was used for the isolation of fungi from the orange fruit and for the preparation of pure culture. The medium was prepared from commercially produced dehydrated medium. 65g of SDA powder was dissolved in 1L of distilled water in a sterile conical flask covered with cotton wool and aluminum foil paper. It was mixed thoroughly and autoclaved at 121°C for 15minute under a pressure of 15 pounds. The medium was allowed to cool after autoclaving to 50°C; Chloramphenicol was added to inhibit the growth of bacteria and was then dispensed aseptically into sterile petri dishes.

### **Determination of disease incidence**

Oranges were collected at random in two markets in Mubi (Mubi main market and Kasuwan kuturu) market respectively. The incidence of disease was determined by counting the infected oranges using the following formula;

$$\text{Incidence} = \frac{\text{Total number of diseased plants}}{\text{Total number of plants observed}} \times 100$$

### **Isolation and identification of pathogen**

The sliced *Citrus cinensis* fruits prepared were placed on Sabourand Dextrose Agar (SDA) aseptically and incubated at 28°C for five days. A pure culture was obtained and maintained by sub-culturing each of the different colonies that emerged onto the SDA plates and incubated at 28°C for five days. As a control, healthy fruits were sterilized with 75% ethanol. The fruits were cut in segments with a sterile blade, placed on SDA and incubated at 28°C for five days. The fungal isolates were identified using cultural and morphological features such as colony growth pattern, conidial morphology and pigmentation as described by [13]. The technique of [14] was used for the identification of the isolated fungi with the use of safranin stain.

### **Pathogenicity test**

Pathogenicity test was carried out in order to ascertain whether the isolated fungi were really responsible for the spoilage of citrus fruits. Healthy fruits were sterilized with ethanol cylindrical plug tissues were cut out from the fruits using a sterilized 2mm sized cork borer. Agar plate containing a weak old fungal culture was aseptically placed in the holes; it was covered and sealed by means of petroleum jelly. The procedures were repeated separately across each of the fungal isolates. The inoculated samples and the control were placed in sterile polythene bags and were incubated in an oven for 5 days. The point of inoculation of each type of fungus were examined and recorded. The diameter of the rotten portion of the orange was measured. The fungi were then re-isolated from the inoculated fruits and compared with the initial isolates.

### **Preparation of leaf extract**

The preparation of the leaf extract was carried out by the procedures described by [15]. Dried leaves of *Phyllanthus amarus* were grinded into fine powder and sieved. The powder was dissolved in the solvent (ethanol) and shaken vigorously for 5-10 minutes and left for 24 hours after which the extract was filtered and dried under reduced pressure. The extract was prepared into four different concentrations ranging 50, 100, 200 and 400 mg/ml. The extract concentration was prepared by weighing 3.4g of the extract into 3ml of ethanol; a doubling dilution of the diluted extract was carried out into four different labeled bottles to obtain the concentrations.

### **Susceptibility testing**

This was carried out using agar well diffusion method. The standardized organisms were dropped on the freshly prepared SDA. Four wells were made on the inoculated SDA plate using a sterile cork borer of 3mm. The wells were properly labeled according to the different concentrations of the extract prepared. The punched wells were filled with 0.2ml of each of the extract (50, 100, 200 and 400 mg/ml), the plate were then allowed to stay on the

laboratory table for one hour for the extract to diffuse into the agar after which they were incubated at 28°C for 24 hours. The plates were observed for any evidence of inhibition which appeared as clear zones that was completely devoid of growth around the wells. The diameter of the clear zones was measured with a transparent ruler calibrated in millimeter (mm). Ketoconazole was used as positive control.

## RESULTS

### Isolation and identification of pathogens

Four fungal pathogens (*Aspergillus niger*, *Aspergillus flavus*, *Penicillium expansum* and *Rhizopus stolonifer*) were isolated and confirmed through pathogenicity test to be responsible for sweet orange fruit rot in the markets (plate i-iv).



Plate Ia: Pure culture of *A. flavus*

Plate Ib: Photomicrograph of *A. flavus*




Plate IIa: Pure culture of *P. expansum*



Plate IIb: Photomicrograph of *P.*




Plate IIIa: Pure culture of *A. niger*



Plate IIIb: Photomicrograph of *A. niger*




Plate IVa: Pure culture of *R. stonolifer*

Plate IVb: Photomicrograph of *R. stonolifer*

### Disease Incidence

The percentage (%) incidence of pathogens in the two markets surveyed shows that rot occurred in both market, whereby Mubi main market had the highest incidence with 35% and kasuwa kuturu with 25% (Table 1) respectively.



Table 1: Occurrence and Incidence of Fungal Pathogens.

Fungal isolates	Presence of isolates in each market	
	Mubi main market	Kasuwan kuturu
<i>Aspergillus niger</i>	+	+
<i>Aspergillus flavus</i>	+	-
<i>Penicillium expansum</i>	+	-
<i>Rhizopus stolonifer</i>	-	+
Incidence %	35	25

Key: + = Present, - = Absent

### Pathogenicity test

The pathogenicity test revealed that the pathogen isolated from the infected oranges were pathogenic (Table.2).The rate of decay caused by the isolated fungi *A. flavus* and *A. niger* to be most pathogenic with rapid disintegration of the healthy fruits. The pathogens having rot diameter of 53mm and 45mm respectively and shows similar growth characteristic features to the original diseased samples.

Table 2: Pathogenicity test

TEST ORGANISMS	DIAMETER OF ROT/ (mm)
<i>A. Niger</i>	45.00
<i>A. flavus</i>	53.00
<i>P. exapansum</i>	21.00
<i>R. stolonifer</i>	15.00

### Effect of Leaf Extract on the Isolated Fungi

The effect of the leaf extract of *Phyllanthus amarus* on the isolated fungi is given below (Table 3). There was a significant inhibition of growth of pathogens treated with *P. amarus* leaf extract. The inhibition diameter was best in *A. flavus* at the highest concentration (400mg/ml). The extract shows no inhibition at the least concentrations (50mg/ml) in all the isolated fungi. There was significant difference between the control and all other concentrations at probability level of  $p < 0.05$ .

**Table 3:** Effect of Leaf Extract of *Phyllanthus amarus* on the Isolated Fungi

(mg/ml)	Diameter Of Zone Of Inhibition (mm)			
	<i>A. niger</i>	<i>A. flavus</i>	<i>P. expansum</i>	<i>R. stolonifer</i>
<b>400</b>	9.000 <sup>d</sup>	10.667 <sup>c</sup>	9.333 <sup>c</sup>	8.333 <sup>c</sup>
<b>200</b>	6.667 <sup>c</sup>	9.333 <sup>c</sup>	8.333 <sup>c</sup>	7.333 <sup>c</sup>
<b>100</b>	4.000 <sup>b</sup>	5.000 <sup>b</sup>	5.000 <sup>b</sup>	4.000 <sup>b</sup>
<b>50</b>	0.00 <sup>a</sup>	0.000 <sup>a</sup>	0.000 <sup>a</sup>	0.000 <sup>a</sup>
<b>Control</b>	11.667 <sup>e</sup>	11.333 <sup>d</sup>	11.333 <sup>d</sup>	11.667 <sup>d</sup>

**NB:** Means with different superscript along the column are statistically significantly different at  $p < 0.05$ . **SE**  $\pm$  = Standard Error.

## DISCUSSION

Fruits and vegetables are very important and have high dietary and nutritional qualities. Consumption of fruit and vegetable products has dramatically increased by more than 30% during the past few decades [1]. However, microorganisms have been causing serious threats to agricultural produce. This study revealed the presence of *A. niger*, *A. flavus*, *P. expansum* and *R. stolonifer* and confirmed through pathogenicity to be responsible for the spoilage of *C. sinensis* (sweet orange) fruits sold in Mubi. This result agrees with the work of [16] where they reported *Aspergillus* spp, *Mucor* spp, *R. stolonifer* to be associated with rotten *Citrus* spp. It also corresponds with the work of [17] and [18] who also reported these pathogens to be causative agents of orange fruits rot. [19] also found out that *Aspergillus* was the common genus affecting different types of fruits, it appeared on 86.6% of orange fruits where *Aspergillus niger* and *Aspergillus flavus* were the dominant on the tested fruits of the fruit tested (Apple, Orange, Mango and Grape). [16] in their work reported fungal pathogens responsible for the spoilage of orange include *Aspergillus niger*, *Aspergillus flavus*, *Aspergillus fumigates*, *Aspergillus terreus*, *Mucor heimalis*, *Mucor racemosus*, *Absidia*, *Carmbifera*, *Rhizophus stolonifer* and *Fusarium oxysporum* are associated with *Citrus* species.

The percentage (%) incidence of pathogens in the two markets surveyed, shows that rot occurred in both markets, main market had the highest incidence with 35% and kasuwa kuturu with 25% whereas [20] reported kasuwa kuturu having highest with 58.3% and central market the least 33.3% this might be either due to the method of sample collection, weather and also duration in which the research was carried out.

The efficacy of the ethanolic leaf extract of *Phyllanthus amarus* on the isolated fungi shows that the rate of inhibition increases with increase in concentration of the extract. [21] reported that the leaf extract of *P. amarus* was effective on the growth inhibition of some fungal pathogens including *Aspergillus niger* and *A. flavus*. [22] also reported that the leaf

extract of *P. amarus* had a broad-spectrum activity against gram negative and gram positive bacteria as well as fungal species.

## CONCLUSION

It can be concluded that *A. niger*, *A. flavus*, *P. expansum* and *R. stolonifer* are pathogens causing rots in *Citrus* fruits. However, incidence of rot varies in the two markets. Ethanolic leaf extract of *P. amarus* was found to be effective in inhibiting the growth of the pathogen. There was increase in inhibition of the pathogen with the increase in concentration of the extract.

## RECOMMENDATION

Fruits and vegetables in general are known to be perishable and have a relatively short shelf-life causing them to deteriorate rapidly. Therefore, sweet orange should be consumed shortly after harvesting or properly refrigerated (or stored using other means) and should be discarded if any alteration in color or taste of the fruit is noticed as this can be hazardous to human health.

## REFERENCES

1. Barth, M., Hankison, T.R., Zhuang, H., Breidt, F. *Microbiological spoilage of fruits and vegetables*. In: Sperber W.H, Doyle M.P, editors. *Compendium of the Microbiological Spoilage of Foods and Beverages, Food Microbiology and Food Safety*. New York: C Springer Science Business Media, LLC. 2009; 135–183.
2. Zubair, N. A. Determination of microbial characteristics of selected fruits sold in major markets in Ilorin metropolis. *African Sciences*. 2009; **10**(2): 1595-6881.
3. Muhammed, N.O., Soji-Omoniwa, O., Usman, L.A., Omoniwa, B.P. (2013). Antihyperglycemic activity of leaf essential oil of *Citrus sinensis* (L.). Osbeck on Alloxan-Induced Diabetic Rats. *ARRB* **3**(4): 825-834)
4. Ghasemia, K.Y., Ebrahim-Zadeh, M.A., Samson, V. Antioxidant activity, phenol and flavonoid contents of 13 *Citrus* spp. Peels and tissues. *Journal of Pharmacology Science*. 2009; **22**:227-281
5. Milind, P. and Diev, B. Orange: range of benefit. *International Research Journal Pharmacy*. 2012;**3**(7): 59-63.
6. Guarnier, S., Riso, P., Porrini, M. Orange juice vs vitamin C: Effect of hydrogen peroxide-induced DNA damage in mono-nuclear blood cells. *Nutrition*. 2007; **97**: 639-643.
7. Honow, R., Laube, N., Schneider, A., Kessler, T., Hesse, I. Influence of grape fruit orange and apple juice consumption on urinary variables and risk of crystallization. *Br. Journal of Nutrition*. 2003; **90**: 295-300
8. Vivek, K.R., Nandini, S.S., Anitha, S. Anti-typhoid activity of aqueous extract of fruit peels *Citrus sinensis*. *IJPRD*. 2010; **2**: 217-219.
9. Neeta, S. and Abhishek, T. Effects of *Citrus sinensis* (L.) Osbeck epicarp essential oil on growth and morphogenesis of *Aspergillus niger* (L.). *African Journal of Microbiology Resources*. 2008; **163**:337-344.
10. Moss, M.O. Mycotoxin review. *Journal on Penicillium. Mycologist*. 2002; **16**:116-119.



11. Jay, J.M. Microbial Spoilage of Food. Modern Food Microbiology (4<sup>th</sup>ed.). Chapman and Hall Incorporated. New York. 2003; 187-195.
12. Ismail, M. and Zhang, J. Post-Harvest Citrus diseases and their control, outlooks on pest management. *Peouen*. 2004;15: 29-35.
13. Tafinta, I.Y., Shehu, K., Abdulganiyyu, H., Rabe, A.M., Usman, A. *Isolation and identification of fungi associated with the spoilage of sweet orange (Citrus sinensis) fruits in Sokoto State. Niger. Journal of Basic Applied Science*. 2013; **21**(3):193–196.
14. Oyeleke, A., Manga, S.B. *Essential of Laboratory Practice*. 3rd ed. Minna, Niger state, Nigeria: Tobest Publisher. 2008; 12–29.
15. Das, K., Tiwari, R. K. S., Shrivastava, D. K. Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends *Journal of Medicinal Plants Research*. 2010; **4**(2):104-111.
16. Nasiru, N., Ahmed, Y., Abel, D. Detecting seed borne fungi of soybean by different incubation methods. *Plant Pathology Journal*. 2015; **2**(2): 114-118.
17. Ifeanyi, A.E. Fungi associated with the deterioration of *Citrus sinensis*. B.Sc. Biology Usmanu Danfodiyo University Sokoto, Nigeria (Unpublished) 1995.
18. Bello, A.A. Microorganisms associated with the deterioration of some Citrus fruits. M.Sc. Botany Usmanu Danfodiyo University Sokoto, Nigeria (Unpublished) 2010.
19. Abdullahi, S.M., Musa, A.M., Yaro, A.H., Magaji, M.J, Sule, M.I., Sani, Y.M. Anticonvulsant screening of chloroform: soluble fraction of methanolic stem-bark extract of *Lonchocarpus sericeus*, Poir (Papillonaceae). *Biology Environment Journal Tropic*. 2013; **9**:8-12.
20. Usman, E.N. The effect of Aloe vera extract on orange fruit post-harvest rot agent in Mubi metropolis. *Unpublished B.Sc thesis*, Adamawa State University, Mubi, 2018.
21. Ilondu, E.M, Echigeme, M.V., Isitohan, F.O. Bioactive principles and antifungal activity of acetone extract fractions of three tropical spices against *Aspergillus niger* causing fruit rot of African pear (*Dacryodes edulis*). In: Book of Abstract and Programme, 40th Annual Conference of Nigerian Society for Plant Protection (NSPP) Abuja. 2015; 36.
22. Senjobi, B.A., Ogunkunle, A.O., Adekola, E.A. Effect of different land use types and their implications on land degradation and productivity in Ogun State. *Nigeria Journal of Agricultural Biotechnology and sustainable development*. 2016; **3**(1):7-18