

# **Original Research Article**

## **ANTIBIOTIC RESIDUES IN WASTE IMPACTED SURFACE WATERS FROM LAGOS, NIGERIA.**

---

**ABSTRACT** In the past few decades, there has been increasing interest and concern about the environmental presence of human and veterinary pharmaceutically active chemicals (PHACs). and the implication of such on ecosystem health at organismal and population levels This study was conceived, to identify and quantify the concentrations of an important sub-group i.e., the pharmaceuticals. Which have been shown to have ecological implications on various aquatic life-forms. In surface water bodies in Lagos Nigeria.

**Study design:** water specimens taken from six locations in the city of Lagos in Nigeria were analysed using SPE coupled with tandem LC/MS.

**Place and Duration of Study:** The project, split site collaboration. Spanned a period of nine months between 30.09.2013 to 30.04.2014. Analysis of water specimens was done at the IWW Water Centre. Mulheim, Germany.

**Methodology:** Analysis of water samples were carried out by IWW Water Centre, Mulheim, Germany according to published guidelines or validated in-house methods. Waters Acquity ultra performance liquid chromatography (UPLC-TQD/PDA and UPLC-TQS) (Milford) was used to carry out LC-MS/MS analyses. For GC-MS analyses an Agilent GC-MS system (6890 GC and 5973 MSD single quadrupole mass analyzer was used.

**Results:** In surface water samples, 12 pharmaceutical agents were detected. Of these Six antibiotics, Viz: Chloramphenicol, Erythromycin, Erythromycin-A dehydrate, Sulfadiazine, and Sulphaamethoxazole Trimetoprim were detected out of thirteen screened for, at environmentally significant concentrations.

**Conclusion:** This study has established the presence and locational prevalence of primary antibiotic and antibiotic degradates in surface waters in Lagos, Nigeria More studies are required to investigate the ecological implications of such presences.

### **Introduction**


An increasing number of pharmaceutical products are being used on a daily basis as the pharmaceutical industry continues to introduce new and enhanced products for the prophylaxis and active treatment of traditional infectious/transmissible, and increasingly important non communicable, diseases. These pharmaceuticals have been demonstrated to be incompletely removed from waste waters by current sewage treatment regimes, with some entirely unchanged. Antibiotics are among the most widely prescribed and used pharmaceutical classes in human and veterinary therapeutics, and non-prescription antibiotic use is widespread, especially in developing countries where these medicines are widely available for purchase and use without prescriptions (Okeke and Lamikanra, 1995; Van Duong *et al.*, 1997; Morgan *et al.*, 2011; Tamuno and Mohammed, 2011), due to a lack of regulation and monitoring for compliance. In these settings, antibiotics are seen as “cure alls” for virtually every real and imagined illness, and are used indiscriminately by individuals ignorant of the implications of such self-medication, especially the development of antibacterial resistance, and the associated human and environmental sequelae. Antibiotics have been described in wastewaters and sewage treatment influents and effluents

consequent on human and veterinary use by several investigators including, Hirsch *et al.*, (1999), Chang *et al.*, (2008), Kemper, (2008), and Gao *et al.*, (2012). In third world urban situations, where waste water treatment is almost completely absent, PhACs, including antibiotics, could potentially enter surface waters without any preliminary treatment, and the effects consequent on this fact, especially on non target species such as fish and other aquatic fauna, could potentially be even more profound than conjectured or realised at present. Several in vitro studies detail the acute and chronic effects of antibiotics on aquatic species. Indeed, the antibiotics are the singular most studied pharmaceutical class for acute toxicity effects on aquatic vertebrates and invertebrates (Wollenberger *et al.*, 2000; Le Bris and Pouliquen, 2004) with about 40 studied till date (Brausch *et al.*, 2011). Lagos, the commercial capital of Nigeria, with an estimated population of about twenty one million people, is one of the world's fastest growing metropolises, already gaining megacity status. Lagos is estimated to be one of the fastest growing megacities in the world. High population density clusters in largely unplanned urban and peri-urban locations, poverty and the overstretched public infrastructure and works, are largely descriptive of living conditions in Lagos. The generally poor sanitary conditions in most of the city and suburbs ,aggravated by the tropical climate characterized by seasonal high precipitation levels, and a lack of primary sewage treatment make wastewater impaction on surface waters a stark and ongoing reality. This study, which is consequent on an earlier investigation into pharmaceutical use and disposal patterns in households in Lagos, seeks to identify and quantify the levels of antibiotic residues in surface waters from six deliberately chosen locations within Lagos municipality and to compare detected concentrations with existing information from other studies for perspective.

## **Materials and Methods**

### **Sampling locations**

**Selection criteria for the locations surveyed included five intra- municipal locations Amuwo-Odofin, via Badagry, Isolo, Ojo, and Agbara.**



**Figure I: Sampling locations.**

500 ml. sub- surface water samples were collected in duplicate, on each collection date from each of the five locations. s into amber PET capped bottles. Suggested by the USGS publication National field manual for the collection of water-quality data (USGS, 2006). All locations were lotic water sites. On the spot water parameter estimation were done using a handheld Henna instruments<sup>®</sup> model HI-98129 meter (Table 1). Following collection, water bottles were placed in Styrofoam coolers with ice packs and transferred to the laboratory where kept at -20 degrees centigrade until analysed.

Table 1: Showing locational details of water collection points and water quality measurements of samples.

Sample ID	Location	GPS	Temp. (C)	DO (PPM)	Conductivity ( $\mu$ s)	pH
243598	River Owo	6°29'54.77"N; 3°06'11.98"E	26.4 °C	68	134	7.06
243600	Ojo	6°22'31.24"N;3°12'28.76"E	31.4 °C	2000	3999	7.83
243601	Liverpool	6°26'39.34"N;3°21'49.32"E	34.7 °C	64	3999	6.6

243597	Ajido	6°24'23.87"N;2°56'52.32"E	29°C	123	488	7.65
243602	Isolo	6°31'45.41"N;3°18'57.06"E	27.7°C	245	492	7.45
243599	Amuwo Odofin	6°28'47.5"N;3°17'47.5"E	28.2°C	2000	3999	7.6

Pre-processing of water samples for LC/MS

### Sample preparation and cleanup

An off-line Solid Phase Extraction (SPE): procedure was used for clean-up and pre-concentration of samples. All experiments were performed using automated sample preparation equipment, fitted with an external pump for dispensing samples through the SPE cartridges and with a switching valve for the selection of samples. Recovery studies were performed using surface water samples spiked to final concentrations of 0.1 and 1 µg/L of each pharmaceutical. The recoveries were achieved working in the neutral pH range using Oasis HLB 500 mg extraction cartridges. The conditions for Oasis HLB extraction were: SPE cartridges were conditioned with 6 mL methanol followed by 6 mL HPLC water at a flow-rate of 1 mL/min. One litre of filtered water sample was loaded at 60 mL/min. The cartridges were then washed to reduce highly polar interferences by running 1 mL 5% methanol in HPLC water through the cartridge. Elution of the analytes from the cartridge was done by first eluting with two 3 mL aliquots of methanol containing 5 mM Tetrabutylammonium chloride (TBACl). The eluted volume was then dried in a water bath at 30°C under a stream of nitrogen. The sample was reconstituted with 300 µL of a water-methanol solution (30:70, v/v). Finally, 20 µL were injected into the LC-MS-MS system.

### Analysis of Water Samples

LC/MS Analysis: of water samples were carried out by IWW Water Centre, Muelheim, Germany according to published guidelines or validated in-house methods (**Error! Reference source not found.**). Waters Acquity ultra performance liquid chromatography (UPLC-TQD/PDA and UPLC-TQS) coupled with an electrospray ionization tandem mass spectrometric system (Milford) was used to carry out LC-MS/MS analyses. For GC-MS analyses an Agilent GC-MS system (6890 GC and 5973 MSD single quadrupole mass analyzer) equipped with an automatic sampler (MPS 2) and Cooled Injection System (CIS 4, Gerstel) was used. Data acquisition, processing and evaluation were carried out using Agilent ChemStation software combined with Gerstel Maestro software package. The limit of detection (LOD) was defined by a signal to noise ratio of 3:1 for all applied methods. Tables 1 and 2 show the results of water sample analysis by location. Figure 1 shows the results of some simple statistical analyses on results from table 1 depicted as box and whisker plots.

## Results and discussion

In the present study 6 antibiotics falling into two bloc areas, the sulfa (3), macrolide (1), macrolide derivative (1) categories, and one unique entity (chloramphenicol), were determined and discovered at environmentally significant concentrations. The concentrations of the four sulfonamides, sulfamethoxazole, sulfadiazine, and trimethoprim, ranged from a minimum of 0.02 µg/L (Amuwo Odofin) to 1.5 µg/L in the sample from Isolo. Occurrence patterns were similar to earlier reports which indicate the sulfa antibiotics as very widespread in occurrence worldwide. Trimethoprim was detected in two samples (Amuwo Odofin and Isolo) at levels ranging from 0.12-0.40 µg/L. These figures are comparatively higher than recorded for sewage treatment facility (STF) influents and surface waters in several previous studies. Li and Zhang, (2011), reported concentrations of Trimethoprim in STF influents as ranging between 0.1-0.154 µg/L, and 0.0844-0.12 µg/L in effluents. Other studies include Karthikeyan and Meyer, (2006), 0.44 µg/L, in influents, and 0.39 µg/L in effluents. Gulkowskaa *et al.*, (2008) reported maximum concentrations in STF influents from five municipal STFs at 0.32 µg/L and Chang *et al.*, (2008) 0.00034 µg/L in river samples, and 0.42 µg/L in STF influents.

Sulfamethoxazole has been widely described in both influents and effluents of STFs all over the world, including the USA by several investigators including the USA (Karthikeyan and Meyer, 2006), and in other countries by investigators, including Batt *et al.*, 2007 (2.8 µg/L); Spongberg and Witter 2008 (0.2610 µg/L), Göbel *et al.*, 2005 (0.57 µg/L), Lindberg *et al.*, 2005 (0.674 µg/L), Clara *et al.*, 2005 (0.145 µg/L), and Choi *et al.*, 2008 (0.193 µg/L). The sulfamethoxazole concentrations detected in this study ranged from between a minimum of 0.09 µg/L and a maximum of 1.5 µg/L, exceeding the levels reported in the references above. Sulfamethoxazole is the sulfa antibacterial detected at the highest concentrations in environmental waters (Garcia-Garlan *et al.*, 2011).

Sulfadiazine (SDZ) is a sulfonamide widely used as a human and veterinary antibiotic to prevent and treat diarrhoea and other infectious diseases. Sulfadiazine has been reported in wastewaters at levels ranging between 5.10–5.15 µg/L (Peng *et al.*, 2006). Garcia-Galan *et al.*, (2011) reported recovery levels of sulfadiazine in surface waters at concentrations ranging from 0.00006-0.00022 µg/L. Our studies detected sulfadiazine at levels ranging between 0.02-0.04 µg/L.

Pharmaceutical Agent	Therapy Group	Analytical Methods		Water Samples from Nigeria					
		Method	LOD	AJIDO	RIVER OWO	AMUWO -Odofin	OJO	LIVERPOOL	ISOLO
			[µg/l]	[µg/l]	[µg/l]	[µg/l]	[µg/l]	[µg/l]	[µg/l]
Chloramphenicol	Antibiotics	LC-MS/MS	0,01	< 0,01	< 0,01	0,03	0,03	0,03	0,36
Chlortetracycline	Antibiotics	LC-MS/MS	0,05	< 0,05	< 0,05	< 0,05	< 0,05	< 0,05	< 0,05
Clarithromycin	Antibiotics	LC-MS/MS	0,05	< 0,05	< 0,05	< 0,05	< 0,05	< 0,05	< 0,05
Erythromycin-A dihydrate	Antibiotics	LC-MS/MS	0,06	< 0,06	< 0,06	0,12	< 0,06	< 0,06	0,48
Erythromycin	Antibiotics	LC-MS/MS	0,06	< 0,06	< 0,06	0,06	< 0,06	< 0,06	1,00
Oxytetracycline	Antibiotics	LC-MS/MS	0,05	< 0,05	< 0,05	< 0,05	< 0,05	< 0,05	< 0,05
Roxithromycin	Antibiotics	LC-MS/MS	0,02	< 0,02	< 0,02	< 0,02	< 0,02	< 0,02	< 0,02
Sulfadiazine	Antibiotics	LC-MS/MS	0,01	< 0,01	< 0,01	0,02	< 0,01	< 0,01	0,04
Sulfadimidine	Antibiotics	LC-MS/MS	0,01	< 0,01	< 0,01	< 0,01	< 0,01	< 0,01	< 0,01
Sulfamethoxazole	Antibiotics	LC-MS/MS	0,01	< 0,01	< 0,01	0,65	0,14	0,09	1,50

Tetracycline	Antibiotics	LC-MS/MS	0,05	< 0,05	< 0,05	< 0,05	< 0,05	< 0,05	< 0,05
Trimethoprim	Antibiotics	LC-MS/MS	0,01	< 0,01	< 0,01	0,12	< 0,01	< 0,01	0,40

Table 2: Detected antibiotics

**Table 3:** Showing high, low and mean concentrations of detected pharmaceuticals

Key: **AB:** antibiotic.

Pharmaceutical agent	Type	LOD	Low	High	Mean
Chloramphenicol	AB	0.01	0.03	0.06	0.045
Erythromycin	AB	0.06	0.06	1	0.53
Erythromycin-A dihydrate	AB	0.06	0.12	0.48	0.3
Sulfadiazine	AB	0.01	0.02	0.04	0.03
Sulfamethoxazole	AB	0.01	0.09	1.5	0.795
Trimetoprim	AB	0.01	0.12	0.4	0.26




Figure 2: Box plot showing the median (green), upper (purple) and lower (red) quartiles, and 95 % confidence intervals (error bars), of detected antibiotics across sampling locations.

Key: CPC: Chloramphenicol; ERY: Erythromycin; EAD: Erythromycin-A-dihydrate; SDZ: Sulfadiazine, SMX: Sulfamethoxazole; TMP: Trimethoprim

Discrete erythromycin was detected at a low of 0.06  $\mu\text{g/L}$  in the water sample from Amuwo Odofin , and at a high in the Isolo sample (1.0  $\mu\text{g/L}$  ). Erythromycin-A-dihydrate was detected in water samples from Amuwo Odofin and Isolo at reportable levels of 0.12 and 0.48  $\mu\text{g/L}$  respectively. This compound has been reported as commonly present in waste waters, STF influents and effluents in several other countries. Measured concentrations in other studies include 1.7  $\mu\text{g/L}$  (Kolpin *et al.*, 2002), 0.59-1.978



µg/L, and 0.43–2.054 µg/L respectively from two STFs in mainland China (Xu *et al.*, 2007), 0.226–1.537 µg/L, and 0.361–0.811 µg/L in STFs from Taiwan (Lin *et al.*, 2009). From the USA, Karthikeyan and Meyer, (2006) reported E-A-D concentrations at 0.06–0.19 µg/L. In the UK, Kasprzyk-Hordern, *et al.*, (2008), and Kasprzyk-Hordern *et al.*, (2009) reported 0.144–10.025 µg/L, and 0.23–2.841 µg/L. from two consecutive studies. The concentrations recorded in the present study fall within the lower value ranges recorded for earlier studies, with the concentrations recorded for Asian locations closest in comparison with present study results.

## Conclusion

In summary, this study has confirmed the presence of antibiotics in waste water impacted surface water bodies in Lagos, Nigeria. The findings from this investigation has also established that the concentrations of detected antibiotic moieties, and the locational frequency of detection are comparable to reports from other countries, especially from developing countries, from where , unfortunately, such reports are few, limiting further comparisons. Isolo and Amuwo Odofin were “hotspots” accounting for the highest number of detected antibiotic compounds, and highest total, and individual antibiotic residue concentrations. The latter findings are expressible as functions of very high populations in these locations, and proportionately higher levels of sewage generation compared to other less densely populated locations such as Badagry. In the case of the Isolo sampling location, other possible exacerbating factors apart from population, include the presence of a large municipal “general” hospital, and an active landfill used for the disposal of hospital and hazardous household wastes (among other more innocuous waste types), upstream of the sampling site. It has been documented that leachate plumes could extend considerable distances from landfill sites, and that even closed/inactive landfills could be sources of pharmaceutical water contamination for considerable periods (Slack *et al.*, 2005; Barnes *et al.*, 2007; Carrara *et al.*, 2008), and this fact could be responsible for the higher levels of pharmaceutical contamination of water, and conceivably other environmental matrices in Isolo. In the absence of more detailed hydrological data it can be reasonably accepted that leachates from the aforementioned landfill could modify the environmental prevalence and concentrations in impacted waters. Another probable exacerbating factor is the lack of municipal liquid sewage stream treatment and poor sanitation, especially in the more crowded and unplanned locations in the city. In locations such as Isolo where waste water impaction is quite high, and water flow rates slow, a possible scenario could be that PHAC concentrations in the water samples could be close to that in undiluted waste water and liquid sewage. Of course dilution in the definitely larger volume of the receiving waters is a factor which could reduce concentrations therein, but probably not by much. In conclusion, antibiotic profiles and presences in the studied locations are higher than in locations where there is effective wastewater and sewage treatment, a fact established through a perusal of extant literature and reviews. In addition, in the absence of reliable prescription, purchase, and usage statistics, these significant antibiotic presences cannot be precisely tied to usage (though as mentioned earlier, they are widely available and abused), but are probably more a consequence of poor sanitation and ineffective sewage treatment in the city of Lagos. Where waterworks do exist, the surface waters are used as water sources for recycling and domestic reuse especially for drinking. A possible scenario is that antibiotics are probably detectable in drinking water where available, and the associated sequelae on the consuming public can only be

conjectured at this point. More work is required to determine STF clearance, and concentrations of important PHACs in surface, ground and drinking waters, to move information and studies from the “snapshot” establishment of presence status to the level of sustained observation and monitoring.

#### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

#### **References**

- Okeke IN, Lamikanra A. Quality and bioavailability of tetracycline capsules in a Nigerian semi-urban community. *International Journal of Antimicrobial Agents*. 1995 Jul 1;5(4):245-50.
- Duong DV, Binns CW, Le TV. Availability of antibiotics as over-the-counter drugs in pharmacies: a threat to public health in Vietnam. *Tropical medicine & international health*. 1997 Dec;2(12):1133-9.
- Tamuno I, Mohammed SI. Self-medication with antibiotics amongst students of a Nigerian Tertiary Institution. *J Basic Appl Sci Res*. 2011;1(10):1319-26.
- Hirsch R, Ternes T, Haberer K, Kratz KL. Occurrence of antibiotics in the aquatic environment. *Science of the Total environment*. 1999 Jan 12;225(1-2):109-18.
- Chang SJ, Hsueh TJ, Chen IC, Huang BR. Highly sensitive ZnO nanowire CO sensors with the adsorption of Au nanoparticles. *Nanotechnology*. 2008 Mar 25;19(17):175502.
- Kemper N. Veterinary antibiotics in the aquatic and terrestrial environment. *Ecological indicators*. 2008 Jan 1;8(1):1-3.
- Gao H, Liu Z, Song L, Guo W, Gao W, Ci L, Rao A, Quan W, Vajtai R, Ajayan PM. Synthesis of S-doped graphene by liquid precursor. *Nanotechnology*. 2012 Jun 19;23(27):275605.
- Wollenberger L, Halling-Sørensen B, Kusk KO. Acute and chronic toxicity of veterinary antibiotics to *Daphnia magna*. *Chemosphere*. 2000 Apr 1;40(7):723-30.
- Le Bris H, Pouliquen H. Experimental study on the bioaccumulation of oxytetracycline and oxolinic acid by the blue mussel (*Mytilus edulis*). An evaluation of its ability to bio-monitor antibiotics in the marine environment. *Marine Pollution Bulletin*. 2004 Mar 1;48(5-6):434-40.
- Brausch JM, Rand GM. A review of personal care products in the aquatic environment: environmental concentrations and toxicity. *Chemosphere*. 2011 Mar 1;82(11):1518-32.

Karthikeyan KG, Meyer MT. Occurrence of antibiotics in wastewater treatment facilities in Wisconsin, USA. *Science of the total environment*. 2006 May 15;361(1-3):196-207.

Gulkowskaa A, Leunga HW, Soa MK, Taniyasub S, Yamashitab N, Yeunga LWY, Richardsona BJ, Leic AP, Giesya JP, Lam PKS (2008) Removal of antibiotics from wastewater by sewage treatment facilities in Hong Kong and Shenzhen, China. *Water Res* 42:395–403

Batt AL, Kim S, Aga DS. Comparison of the occurrence of antibiotics in four full-scale wastewater treatment plants with varying designs and operations. *Chemosphere*. 2007 Jun 1;68(3):428-35.

Spongberg AL, Witter JD. Pharmaceutical compounds in the wastewater process stream in Northwest Ohio. *Science of the total environment*. 2008 Jul 1;397(1-3):148-57.

Göbel A, Thomsen A, Mc Ardell CS, Joss A, Giger W. Occurrence and sorption behavior of sulfonamides, macrolides, and trimethoprim in activated sludge treatment. *Environmental science & technology*. 2005 Jun 1;39(11):3981-9.

Lindberg RH, Wennberg P, Johansson MI, Tysklind M, Andersson BA. Screening of human antibiotic substances and determination of weekly mass flows in five sewage treatment plants in Sweden. *Environmental science & technology*. 2005 May 15;39(10):3421-9.

Clara M, Strenn B, Gans O, Martinez E, Kreuzinger N, Kroiss H. Removal of selected pharmaceuticals, fragrances and endocrine disrupting compounds in a membrane bioreactor and conventional wastewater treatment plants. *Water research*. 2005 Nov 1;39(19):4797-807.

Peng J, Zhou MF, Hu R, Shen N, Yuan S, Bi X, Du A, Qu W. Precise molybdenite Re–Os and mica Ar–Ar dating of the Mesozoic Yaogangxian tungsten deposit, central Nanling district, South China. *Mineralium Deposita*. 2006 Oct;41(7):661-9.

Kolpin DW, Furlong ET, Meyer MT, Thurman EM, Zaugg SD, Barber LB, Buxton HT. Pharmaceuticals, hormones, and other organic wastewater contaminants in US streams, 1999– 2000: A national reconnaissance. *Environmental science & technology*. 2002 Mar 15;36(6):1202-11.

Xu W, Zhang G, Li X, Zou S, Li P, Hu Z, Li J. Occurrence and elimination of antibiotics at four sewage treatment plants in the Pearl River Delta (PRD), South China. *Water research*. 2007 Nov 1;41(19):4526-34.

Meyer KE. Asian management research needs more self-confidence. *Asia Pacific journal of management*. 2006 Jun;23(2):119-37.

Kasprzyk-Hordern B, Dinsdale RM, Guwy AJ. The occurrence of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs in surface water in South Wales, UK. *Water research*. 2008 Jul 1;42(13):3498-518.

Kasprzyk-Hordern B, Dinsdale RM, Guwy AJ. The removal of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs during wastewater treatment and its impact on the quality of receiving waters. *Water research*. 2009 Feb 1;43(2):363-80.

Barnes JW, Radebaugh J, Brown RH, Wall S, Soderblom L, Lunine J, Burr D, Sotin C, Le Mouélic S, Rodriguez S, Buratti BJ. Near-infrared spectral mapping of Titan's mountains and channels. *Journal of Geophysical Research: Planets*. 2007 Nov;112(E11).

Carrara A, Crosta G, Frattini P. Comparing models of debris-flow susceptibility in the alpine environment. *Geomorphology*. 2008 Feb 15;94(3-4):353-78.

UNDER PEER REVIEW