

Review Article

Mangrove-Epiphytic Plants in Selected Mangrove Rehabilitation Areas of Davao Occidental, The Philippines

ABSTRACT

The study was conducted in the mangrove rehabilitation sites of Davao Occidental, Philippines particularly Sitio Agdao of Brgy. Tubalan, Sitio Baybay of Brgy. Buhangin and Sitio Lahusan of Brgy. Fishing Village of Malita, Davao Occidental. The study was conducted from October to December 2020. Preliminary stations were established in the conduct of the study. Station 1 in Sitio Agdao, Station 2 in Sitio Baybay and the Station 3 in Sitio Lahusan. This site is considered to be a slightly undisturbed mangrove forest dominated by species of mangrove trees.

The free exploration method was carried out first before the conduct of the study. Samples consisted of host plants and their associates (epiphytes) are collected for proper identification. Three study areas were assessed by the number of individual species to calculate their relative abundance and density.

A total of 7 species identified individually among of the three-study areas. The study also shows that species of mangrove epiphytes in the study area composed of 3 to 5 species per station and exhibit 1 to 3 identified species per plot and individual species per station revealed that *Davallia denticulata* and *Aglaomorpha quercifolia* were the most common species of mangrove epiphytic plants in almost all three study areas. However, Lahusan sampling station occurred numerous quantity of Mangrove epiphytic plants species among 3 identified study areas including *Imperata cylindrica* that cannot be found in Agdao and Baybay Stations.

Statistically, no significant difference was observed among study areas.

Keywords: Mangroves, Epiphytic Plants, Environment, Rehabilitation Areas

1. INTRODUCTION

Epiphyte is one of the subdivisions of the Raunkier system. The term epiphytic derives from the greek “epi” – (meaning ‘upon’) and “phyton” (meaning ‘plant’). Epiphytic plants are sometimes called “air plants” because they do not root in soil. However, there are many aquatic species of algae that are epiphytes on other aquatic plants (seaweeds or aquatic angiosperms) (Everhart *et al.*, 2009).

Most studies describing the flora of mangroves assess the diversity and community structure of trees but do not record the presence of epiphytes (Ashton and Macintosh, 2002). Although epiphytes are very abundant in dense mangroves, few studies have assessed the richness of epiphytic species or specifically of bromeliads (Robertson and Platt, 2001). To understand the distribution patterns of epiphytes and to preserve them, it is important to know not only the diversity, but also the plant host relationships (Magalhães and Lopes, 2015).

Due to the lack of studies and information about epiphytic plants in mangrove forests, and considering that the diversity of these sensitive species can be a tool for mangrove conservation, this study aims to assess the diversity of epiphytic plants in a subtropical mangrove, evaluating their distribution and relationship with their host trees.

The study was conducted to determine and assess the common mangrove-epiphytic plants in selected mangrove rehabilitation sites of Davao Occidental. This study aimed to: (1) Identify the common mangrove-epiphytic plants in the study areas; (2) Determine the density and abundance of mangrove-epiphytic plants in the study areas; (3) Identify the mangroves that serve as host for these epiphytes; and, (4) Measure the prevailing levels of the physico-chemical parameters in each sampling station such as water salinity, temperature and pH.

2. MATERIAL AND METHODS

2.1 Research Locale

The study was conducted in the identified mangrove rehabilitation sites of Davao Occidental, particularly Sitio Agdao of Brgy. Tubalan, Sitio Baybay of Brgy. Buhangin and Sitio Lahusan of Brgy. Fishing Village, Malita, Davao Occidental having a map coordinates a Latitude: $6^{\circ} 20' 59.99''$ N and Longitude: $125^{\circ} 31' 59.99''$ E (www.google.com/map).



Figure 1. The Map of Tubalan Cove, Malita, Davao Occidental showing the location of the study sites.

2.2 Research Design and Establishment of Sampling Sites

The study took place during a period of 2 months and was conducted in the coastline of Tubalan Cove in Malita, Davao Occidental where the selected mangrove rehabilitation sites is situated. Three stations were established in the conduct of the study. Station 1 in Sitio Agdao, Station 2 in Sitio Baybay and the Station 3 in Sitio Lahusan. The distance between the stations was approximately 5 km. This site was considered to be a slightly undisturbed mangrove forest dominated by species of mangrove trees. The area is regularly inundated by a normal high tide.

2.3 Data Gathering Procedure

The free exploration method was carried out first before the conduct of the study. Research samples were collected from an area of 300m². These samples consisted of host plants and their associates (epiphytes) which are attached to the body of the hosts. The next process involved identification of the types of the host plants, the types of their associates, and the zones (parts of the host's body) on which the epiphytes grow. Each mangrove tree was surveyed in an appropriate way. In the case of epiphyte occurrence on higher sprays, it is necessary to climb the tree for the purpose of determining the epiphyte species.

The three parts (zones) of the host plants that are attached to the associated plants. The three zones consisted of zone 1 (lower part), zone 2 (middle part), and zone 3 (upper part), as illustrated in Figure 1.




Figure 2. The Attached Zones (Usman *et al.*, 2019)

2.4 Collection of Mangrove Epiphytes and Sampling Design

Epiphytes were collected randomly from the three stations. The samples were placed into ziploc polyethylene bags, which were labeled and transported back to the laboratory for proper identification and further processing.

The Transect Plot Method was employed in the conduct of the study. Three sampling areas were established in the study. In each area, there were three sampling plots established with 10m x10m in dimension. Each plot was regularly spaced from other plots at 10m. Each area has a distance of 50m. Transects were plotted randomly in each sampling station.

2.5 Physico-Chemical Parameters

The Water salinity was determined using an Atago Refractometer. Water Sample from the sampling stations was placed in the slide cover for the convenient reading and recording of data.

Water temperature was determined using a field thermometer. The apparatus was placed into the water for about 5 minutes to ensure proper reading and recording of data.

The pH scale measures the acidity and alkalinity of the water in the sampling stations. It was measured using the HANNA pH meter. About 50ml of sea water sample was collected for pH determination.

2.6 Statistical Tools and Analysis

The Analysis of variance was used to analyze the data using Statistical Packages for Social Sciences (SPSS). Other gathered data were analyzed using the following:

Species Density. All individual mangroves-epiphytic plants species that were found in each quadrat was counted per species and density and calculated with the formula of Odum, 1971.

$$\text{Density} = \frac{\text{No. of Individual of the same species}}{\text{Area (m}^2\text{)}}$$

Relative Abundance. Relative abundance of each mangroves-epiphytic plants species was estimated based on the formula of Odum, 1971.

$$\text{Relative Abundance} = \frac{\text{No. of Individual of a Species}}{\text{Total No. of Individual of all Species}} \times 100$$

3. RESULTS AND DISCUSSION

3.1 Species of Mangrove-Epiphytic Plants

A total of 6 mangrove-epiphytic plants were identified and distributed among study areas namely ¹Rabbits Foot Fern (*Davalia denticulata*), ²Oakleaf Fern (*Aglaomorpha quercifolia*), ³Bird's Nest Fern (*Asplenium nidus*), ⁴Java Fern (*Microsorium pteropus*), ⁵Golden Anubias (*Anubias barteri*), and ⁶Kugon Grass (*Imperata cylindrica*) (Table 1). These were identified using available field guide and other references, and with consultation from technical experts.

The physico-chemical parameters of the study areas were also measured and noted for further analysis. Salinity shows slightly differences among study areas. Bagumbayan is slightly higher than Baybay and Agdao stations with general average of 34 ppt. Temperature ranges from 29°C to 30°C which makes an average of 30°C. pH also exhibits slightly difference ranges from 6.8 to 7.2. The reading of salinity and pH shows little significance in the growth of mangrove epiphytes since most of the species of epiphytes occur near the trunk area. Thus, the measurement of temperature shows tolerable limit for epiphytes.

Table 1. Identified Species in the Davao Occidental

Scientific Name	English Name
<i>Davalia denticulata</i>	Rabbit Foot Fern
<i>Aglaomorpha quercifolia</i>	Oakleaf Fern
<i>Asplenium nidus</i>	Bird's Nest Fern
<i>Microsorium pteropus</i>	Java Fern
<i>Anubias barteri</i>	Golden Anubias
<i>Imperata cylindrica</i>	Kugon Grass

It was also noticed that matured mangrove trees (approximately 2 meters height starting from the base) serves as common host for most mangrove-epiphytic plants. Probable reason is that these mangrove-epiphytic plants attain favourable light conditions enough to grow under the dense canopy of mangrove trees. These epiphytic-plants do not acquire water and minerals from their host (tree) (Yong J.W *et al.*, 2014). Further, it was also observed that most mangrove-epiphytic plants were found mostly in the middle zone (trunk) of the trees.

3.2 Relative Abundance of Mangrove-Epiphytic Plants

3.2.1 Sitio Agdao

Results of the study shows that species of mangrove epiphytes in the study area composed of 3 to 5 species per station and exhibit 1 to 3 identified species per plot. The species of *Davalia denticulata* and *Aglaomorpha quercifolia* were the most common species of mangrove-epiphytic plants in the Agdao Station with relative abundance higher than 50% followed by *Asplenium nidus* in 33.33%.



Plate 1. *Davalia denticulata*



Plate 2. *Aglaomorpha quercifolia*

3.2.2 Sitio Baybay

Baybay station exhibited 3 mangrove-epiphytic plants species namely *Anubias barteri*, *Microsorium pteropus* and *Asplenium nidus*. The most common species that can be found in all plots was *Anubias barteri* with relative abundance of 46%.



Plate 3. *Anubias barteri*

3.2.3 Sitio Lahusan

Among 3 stations, Sitio Lahusan has the most numbered of mangrove-epiphytic plants with 4 species identified (*Imperata cylindrica*, *Aglaomorpha quercifolia*, *Microsorium pteropus*,

and *Davalia denticulata*). Of these mentioned species, *Aglaomorpha quercifold* still exhibited as the most abundant with 54%. In this station, the species of *Imprerata cycindrica* can be noticed, in which this cannot be found in other 2 stations.



Plate 4. *Aglaomorpha quercifold*

3.3 Density of Mangrove-Epiphytic Plants

The Agdao station shows much higher density of 0.01 to 0.04 species per 100m² with *Davalia denticulata* and *Aglaomorpha quercifold* as the most popular species followed by *Asplenium midus*.

Sitio Baybay station's exhibited 0.02 to 0.03 species per 100m² of density with *Microsorium pteropus*, *Anubias barteri* and *Asplenium midus* as the most commonly found mangrove-epiphytic plants.

The case of Lahusan station revealed species density with 0.01 to 0.02 species per 100m². This station was observed the presence of *Imprerata cycindrica* that cannot be seen in other two stations.

Table 2. Density of Mangrove-Epiphytic Plants among stations

Station	Mangrove-Epiphytic Plants	Density (100m ²)
Sitio Agdao, Brgy. Tubalan, Malita, Davao Occidental	<i>Davalia denticulate</i> <i>Aglaomorpha quercifold</i> <i>Asplenium midus</i>	0.01 – 0.04
Sitio Baybay, Brgy. Tubalan, Malita, Davao Occidental	<i>Microsorium pteropus</i> <i>Anubias barteri</i> <i>Asplenium midus</i>	0.02-0.03
Sitio Lahusan, Brgy. Fishing Village, Malita, Davao Occidental	<i>Imprerata cycindrica</i> <i>Aglaomorpha quercifold</i> <i>Microsorium pteropus</i> <i>Davalia denticulata</i>	0.01-0.02

3.4 Identified Mangroves as Host for Epiphytic Plants

There are 2 mangrove species that serves as host of epiphytic plants namely *Sonneratia sp.* and *Avicennia sp.* (Table 3). Most of the identified mangrove-epiphytic plants attached to *Sonneratia* species. The species of *Anubias bacteri* and *Imprerata cycindrica* are thriving in the body of *Avicennia* species.

Table 3. Identified Mangroves as host of epiphytic plants.

Station	Mangrove-Epiphytic Plants	Mangroves
Sitio Agdao, Brgy. Tubalan, Malita, Davao Occidental	<i>Davalia denticulata</i> <i>Aglaomorpha quercifolia</i> <i>Asplenium midus</i>	<i>Sonneratia sp.</i> <i>Sonneratia sp.</i> <i>Sonneratia sp.</i>
Sitio Baybay, Brgy. Tubalan, Malita, Davao Occidental	<i>Microsorium pteropus</i> <i>Anubias barteri</i> <i>Asplenium midus</i>	<i>Sonneratia sp.</i> <i>Avicennia sp.</i> <i>Sonneratia sp.</i>
Sitio Lahusan, Brgy. Fishing Village, Malita, Davao Occidental	<i>Imperata cylindrica</i> <i>Aglaomorpha quercifolia</i> <i>Microsorium pteropus</i> <i>Davalia denticulata</i>	<i>Avicennia sp.</i> <i>Sonneratia sp.</i> <i>Sonneratia sp.</i> <i>Sonneratia sp.</i>

These mangrove-epiphytic plants mostly can be found in the Zone 2 (Middle Part) of the mangrove trees.

Lahusan station with the most numbered mangrove-epiphytic plant species revealed that these plants can be found in the middle part (100%) of the tree. Sitio Agdao with 83.33%, and Sitio Baybay (66.66%), are observed to be found at the middle area of the mangrove trees. Others are located in zone 3 (upper part of the mangrove trees) (Table 4).

Table 4. Zonation for Mangrove-Epiphytic Plants.

Station	Total No. of Samples	Zone 1 (lower part near the roots)	Zone 2 (middle part of the body near trunk)	Zone 3 (upper part near twigs and leaves)
Sitio Agdao, Brgy. Tubalan, Malita, Davao Occidental	6	-	83.33%	16.66%
Sitio Baybay, Brgy. Tubalan, Malita, Davao Occidental	6	-	66.66%	33.33%
Sitio Lahusan, Brgy. Fishing Village, Malita, Davao Occidental	7	-	100%	-

3.5 Physico-Chemical Parameters

The physico-chemical parameters of the study areas were also measured and noted for further analysis (Table 5).

Salinity shows slightly differences among study areas. Bagumbayan is slightly higher than Baybay and Agdao stations with general average of 34 ppt.

Temperature ranges from 29⁰C to 30⁰C which makes an average of 30⁰C. pH also exhibits slightly difference ranges from 6.8 to 7.2.

Many epiphytes require high exposure and others and cannot endure either as much sun or the associated aridity. Water deficit and high light intensity might hinder the growth of epiphytes and any state of environmental change, photosynthesis and growth processes are likely be affected (Hietz and Briones, 2008).

Table 5. The physico-chemical parameter of the study area.

Station	Salinity	Temperature	pH
Sitio Agdao, Brgy. Tubalan, Malita, Davao Occidental	33ppt	30.0 degree celsius	6.8
Sitio Baybay, Brgy. Tubalan, Malita, Davao Occidental	35ppt	29.9 degree celsius	7.2
Sitio Lahusan, Brgy. Fishing Village, Malita, Davao Occidental	34ppt	30.6 degree celsius	6.9

4. CONCLUSION

There were 7 species were identified among study areas individually among of the three-study area. The study also shows that species of mangrove epiphytes in the study area composed of 3 to 5 species per station and exhibit 1 to 3 identified species per plot.

Individual species per station revealed that *Davalia denticulata* and *Aglaomorpha quercifolia* were the most common species of mangrove epiphytic plants in almost all three study areas.

The results of the study and statistical analysis in regards with the relative abundance and relative density of Mangrove epiphytic plants species in the study areas exhibits no significant difference.

In all stations most epiphytic plants can be seen in the middle part of the mangrove the body near trunk.

Physico-chemical parameters shows having slightly differences among study areas.

ACKNOWLEDGEMENTS

The authors are very grateful to the Southern Philippines Agri-Business and Marine and Aquatic School of Technology (SPAMAST) in completing this study. We thank the Faculty Members and Staff of SPAMAST: Dr. Marlyn B. Llamag, Dr. Ruth S. Lucero, Dr. Pedro M.

Avenido, Dr. Dario R. Morastil, Prof. Edgar Alla P. Firman and all SPAMAST Faculty and Staff for their participation and assistance during the conduct of this study.

COMPETING INTERESTS

The authors have declared that no competing interests exist.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Andrade, J. L., Cach-Perez, M.J., Chilpa-Galvan, N., Tamayo-Chim Ma., Orellana R. And Reyes C. (2013). *Climatic and structural factors influencing epiphytic bromeliad community assemblage along a gradient of water-limited environments in the Yucatan Peninsula, Mexico*. Trop ConservSci 6: 283-302.
2. Ashton, E., Macintosh D. (2002). *A review of mangrove biodiversity conservation and management*. Centre for Tropical Ecosystems Research, University of Aarhus, Denmark.
3. Benzing, D.H. (1990). *Vascular epiphytes: general biology and related biota*. Cambridge: Cambridge University Press, 354 p.
4. Calumpong, H.P., Maypa, A.P. & Magbanua, M. (1987). *Population and algininate yield and quality assessment of four Sargassum species in Negros Island, central Philippines*. Hydrobiologia 398, 211–215 (1999).
5. Everhart, S., Ely J., Keller, H. (2009). *Evaluation of tree canopy epiphytes and bark characteristics associated with the presence of Corticolous myxomycetes*. P. 509-517.
6. Giri, C., Ochieng, E., Tieszen, L.L., Zhu, Z., Singh, A., Loveland, T., Masek, J. & Duke, N. (2010). *Status and distribution of mangrove forests of the world using earth observation satellite data (PDF)*. Global Ecology and Biogeography. 20 (1): 154–159. doi:10.1111/j.1466-8238.2010.00584.x. Retrieved 2012-02-08
7. Greenway, M. (1995). *Trophic relationships of macrofauna within a Jamaican seagrass meadow and the role of theechinoid Lytechinus variegatus (Lamarck)*. Bulletin of Marine Science, 56(3), 719–736.
8. Hassan R., Scholes R., Ash N. (2005). *Millenium Ecosystem Assessment: Ecosystems and Human Wellbeing*, Appendix A, Volume 1, Current State and Trends. Island Press, Washington.
9. Hill, M. (1973). Diversity and evenness: a unifying notation and its consequences. Ecology, 54, 427–432
10. Hogan, M. (2010). Fern. *Encyclopedia of Earth*. National Council for Science and the Environment. Archived November 9, 2011, at Wayback Machines.

11. Magalhaes, J.L. And Lopes, M. (2015). *Species richness and abundance of low-trunk herb epiphytes in relation to host tree size and bark type, eastern Amazonia*. Rev Árvore 39: 457- 466.
12. Naidoo Y., Steinke T.D, Mann F.D, Bhat And Gairolas. (2008). *Epiphytic organisms on the pneumatophores of the mangrove Avicennia marina: occurrence and possible function*. African Journal of Plant Science Vol. 1 (2), pp. 012-015
13. Pacyao, J.P.R, Genciano, V.M.P. 2018. *Management Strategies Employed under PNAP Mangrove Rehabilitation Project in Davao del Sur, Philippines*. International Journal of Current Research, 10, (7), 71081-71094.
14. Pacyao, J.P.R, Barail, S. 2020. *Anthropogenic Activities inside the Mangrove Conservation and Rehabilitation Area: A case of Davao Occidental, Philippines*. International Journal of Fisheries and Aquatic Research. 8 (5):294-298.
15. Pacyao, J.P.R, Macadog, H. O. 2018. *Secondary Productivity of the Philippine National Aquasilviculture Program (PNAP): Mangrove Rehabilitation Project Brgy. Bagumbayan, Malalag, Davao del Sur, Philippines*. International Journal of Fisheries and Aquatic Research. ISSN 2456 – 7248. Volume 3; Issue 3; July 2018. Page No. 38-41.
16. Pacyao, J.P.R., and Llamag, M.B. 2018. *Success Indicator of the Philippine National Aquasilviculture Program (PNAP) Mangrove Rehabilitation Project in Davao del Sur, Southern Philippines*. Open Science Journal Volume 3 (1).
17. Pawlowicz, R. (2013). *Key Physical Variables in the Ocean: Temperature, Salinity, and Density*. Nature Education Knowledge. 4 (4): 13.
18. Roberston K.M And Platt W.J. (2001). *Effects of multiple disturbances (fire and hurricane) on epiphyte community dynamics in a subtropical forest, Florida, USA*. Biotropica 33: 573-582.
19. Rodriguez, C. And Stoner A. (1990). *The epiphyte community of mangrove roots in tropical estuary; distribution and biomass*. Aquatic Botany. 36:117-126.
20. Seher, D. (2015). *Assessment of Water Quality Using Physico-chemical Parameters of Camligoze Dam Lake in Sivas, Turkey*. Ecologia, 5:1-7.
21. Pratana Bayu A., Alhamd L., Rahajoe. J.R. (2012). *Asosiasi Dan Karakterisasi Tegakan Pada Hutan Rawa Gambut Di Hampangen Kalimantan Tengah*.