

Carp Fattening and Socio-economic Status of Farmers of Dinajpur District, Bangladesh: An Expedient Overview

Abstract

A Survey work was designed to explore the present status of carp fattening and socio-economic standing of the fish farmers of three upazillas of Dinajpur district such as Dinajpur Sadar, Parbatipur, and Birampur, and was continued until June 2021 from July 2020. Several methodological tools, including participatory rural appraisal (PRA) and primarily a questionnaire survey were used to collect data from 59 fish farmers. The majority of the farmers were between the ages of 41–50, where 45% of them had four to five family members, representing 71.19% nuclear and 28.81% joint families. Large number of the farmers (28.81%) could only sign their names. Approximately 81.35% were Muslims and 18.65% belonged to the Hindus where near about half of the farmers' (47.46%) annual income were in between BDT 1,00,001-2,0,000. *Paka* sanitary was utilized by 49.15% respondents. All the farmers drank from tube wells and had access to electricity. The typical pond size was 0.31-40 ha (27.12%), with an average depth of 1.7 m. Farmers owned 59.33% of the ponds, on the other hand, 40.67% were leased ponds and 79.66% had the provision of groundwater for cultivation; however, 50.84% exchanged water whenever required. About 57.63% prepared the ponds and applied 123.5 kg/ha of salt, 12.35-44.5 kg/ha of TSP, and 12.35-75.25 kg/ha of urea. To boost carp growth, 83.08% employed availed growth promoters. Indian major carp and exotic carp were considered for this purpose and have had only fish lice as a constraint throughout the nine to ten months of cultivation period. Commercial feeds were utilized by 40.68% farmers and 81.35% maintained a feeding frequency of two times a day. Majority of them (72.89%) sold live fishes. Lack of scientific knowledge, shortage of high-quality seeds and feeds, lack of funds, and lack of marketing facilities were found to be the main obstructions.

Keywords: Aquaculture, Carp fattening, Survey, Dinajpur district, Bangladesh

1. Introduction

Fish is the second most valuable agricultural produce in Bangladesh, and its production contributes to the livelihoods and employment of millions of people all over the globe. Bangladesh ranked 3rd in inland open water capture production and 5th in world aquaculture production [1]. GDP growth rate in the fisheries sector is 5.74%. The fisheries sector contributes 3.57% to the national GDP and 26.50% to the agricultural GDP of the country. As a result of the implementation of fisheries-friendly activities of the government and the provision of demand-based and appropriate technical services at the farmer and entrepreneurial levels, fish production has increased to 45.03 lakh MT in the Fiscal Year (FY) 2019-20, which is 27% more than the total production (35.48 lakh MT) in FY 2013-14. It mentioned that the total fish production in the country was 7.54 lakh MT in FY 1983-84. Fish production has increased more than six times in 37 years [2]. Fish and fisheries have been an indispensable part of the life and livelihood of the country's people since immemorial time [3]. Pond aquaculture represents huge potentiality accounting for 44.43% of the total recorded production and 57.70% of the area under culture and it has the potential to increase further [4]. There are approximately 47.12 million ha inland closed water bodies in Bangladesh, of which 3.98 million ha ponds are suitable for fish culture [5]. Therefore, the country has huge potential for freshwater aquaculture; this potential cannot be fully utilized for various reasons [3].

The potentiality of carp polyculture is beyond the reach to be sustainable due to climate change which makes the fisheries and aquaculture sectors of Bangladesh vulnerable to environmental degradation in various magnitudes, such as groundwater sinking and

contamination through pollutants etc. [6]. Fisher folks are considered one of the most backward sections in our society. Information on the socio-economic framework of the fish farmers forms a good base for planning and development of the economically backward sector. The lack of adequate and authentic information on the socio-economic conditions of the target population is one of the serious impediments to the successful implementation of the developmental program [7]. Dominating species for pond aquaculture are Indian major carps and exotic carps [8].

Dinajpur district is situated at a higher level from the Bay of Bengal and the dry part of Bangladesh [9] where the total fish production in pond culture was 48170 MT [10]. The farmers of Dinajpur are facing problems of scarcity of water throughout the year except the monsoon season [9]. Fattening is a popular technique to increase biomass in animal rearing and this practice is often followed for the production of beef cattle [11] and crab [12] in Bangladesh. Stocking of overwintered and larger size fish species in carp fattening is considered beneficial to mitigate the fish culture problem of lower water columns under drought-prone Barind areas, because overwintering is a proven technique to obtain the fast growth of fish [13] [14] and larger stocking size under lower density can help to get maximum fish biomass within a shorter period of time [15]. There are some research efforts to mitigate the low alkalinity and high turbidity problems and to use the larger stocking size of fish than traditional practice [16] to increase fish production in carp polyculture ponds under the Barind area.

Meanwhile, some fishermen improved the traditional culture system into carp fattening technology. This approach has enhanced output than the traditional culture system. The success of such aquaculture largely depends on the extension activities provided by different Government and Non-Government Organizations.

2. MATERIALS AND METHODS

2.1 Study period

The survey was conducted for a period of one year from July 2020 to June 2021. Frequent field visits were conducted to collect the data from the fish farmers of the selected upazilas of Dinajpur district.

2.2 Selection of the study area

The study was conducted in three upazilas of Dinajpur district, namely Dinajpur Sadar, Parbatipur, and Birampur. The study area is located in between 25°10' and 26°04' north latitudes and in between 88°23' and 89°18' east longitudes (Figure 1).




Figure 1. Map showing selected upazilas in Dinajpur district

2.3 Collection of data

2.3.1 Questionnaire survey

The present study collected data through the questionnaire interview and participatory rural appraisal (PRA) from fish farmers. For the questionnaire interview, a set of preliminary questionnaires was prepared. For the interview, a simple random sampling method was followed. A total of 59 farmers were interviewed. Before the field survey, background information on the number, location and distribution of fish farms and aquaculture activities was collected.

The questionnaire was divided into several sections. The first section focused on farmers personal information, the second section on-farm information, the third one on culture-related information, the fourth on feed and feeding, and the final section focused on cost analysis.

During the survey, questions were asked systematically and explanations were made wherever necessary. The information supplied by the selected farmers was recorded directly on the interview schedules. To minimize errors, data were collected in local units. These were subsequently converted into appropriate units.

2.3.2 Focus Group Discussion (FGD)

During this study, FGD was used to acquire particular significant issues such as pond preparation, enhanced natural feed, use fertilizer, fishing systems, management of the feeding system in the pond and also feeds used as well as marketing related information of carp fattening, socio-economic conditions of farmers, etc. Each FGD session was conducted including 8-10 fish farmers and overall two sessions were conducted in each upazila.

2.3.3 Cross-Check Interview

It was essential to check the information to justify the collected data after completing it through questionnaire interviews and FGDs. In some cases, cross-check interviews were conducted with the key person in the selected areas, such as Upazila Fisheries Officer (UFO), School teachers, local leaders. NGO workers wherever information was opposing or demanded additional measurement.

2.4 Analysis of data

After collection, data were coded and entered into a computer for analysis. All possible errors and inconsistencies were eradicated for verification of the collected data. The tabular description procedure was mainly used to analyze the collected data. This tabular technique was applied to analyze data using simple statistical tools like averages and percentages. Then the collected data were analyzed, preparing tables and graphs with MS-Excel (Microsoft Excel 2016)

3. RESULTS AND DISCUSSION

3.1 Personal Information of Farmers

This section deals with the farmer's personal information, i.e. age group, sex, religious status, educational status, marital status, size and types of family, annual income, health facilities, sanitary facilities, organizational membership, etc. A total of 59 respondents participated in the survey from three upazilas; Dinajpur Sadar, Parbatipur, and Birampur (Table 1).

Table 1. Number and percentage of respondents participated in the survey

Name of the study area	Respondents (No.)	Respondents (%)
Dinajpur Sadar	16	27.12
Parbatipur	25	42.38
Birampur	18	30.5

3.1.1 Age Group

The present study showed that majority of the farmers (35.60%) was between 41 to 50 years of age group; whereas in Sirajgonj district, the farmers were a bit younger (18 to 43 years) than the present study according to Rana [17]. Middle-aged people are generally active, and they are the most productive group in terms of adopting new technologies and making quick decisions (Figure 2).




Figure 2. Age distribution of the selected fish farmers in the study area

3.1.2 Gender

In the survey, among the selected farmers, 89.83% were male and 10.17% were female. This value represents that a deficient number of women participate in this technology of carp fattening (Figure 3).




Figure 3. Gender distribution of farmers

3.1.3 Religious Status

From the present survey, it was found that 81.35% of fish farmers were Muslims and remaining 18.65 % were Hindus (Figure 4). The findings resemble to the status in Gazipur district [18].




Figure 4. Religious status of the farmers

3.1.4 Educational Status

Education has a significant impact on the modernization of farm business operations since it provides a person with up-to-date information about current procedures as well as technological advances in various production processes. The respondents were classified into five categories. It was found that 28.81% can sign names only, 18.65% can't write and read (Figure 5). The findings made clear that educational status of the representative farmers in this area was not that much improved [19].




Figure 5. Educational status of the fish farmers in the study area

3.1.6 Family Type

From the study, two types of the family had seen, joint and nuclear. The nuclear family generally presents two-generation and in this survey area, where Figure 6 shows that 71.19% of families were nuclear and the rest, 28.81% families were joint which is similar to the findings of Alam [20].




Figure 6. Family type of fish farmers in the selected area

3.1.7 Family Size

The family sizes of the fish farmer were divided into four categories according to the number of the family member. The highest percentage was obtained in the 4 to 5 members family (42.38%) and lowest was above 7 members family which connects to the findings of Ali et al [19]. (Figure 7)




Figure 7. Family size of farmers in the study area

3.1.8 Annual Income of Farmers

The selected fish farmers were grouped into five categories based on the level of their annual income. The second category had the highest proportion (47.46%) of farmers while the lowest proportions of farmers (3.39%) were in the fifth category (Table. 2). According to Zafar *et al.* [9], 46% of farmers' annual income was BDT 1.51-1.8 lac, this relates to the current study.

Table 2. Number and percentage of annual income of the selected fish farmers in the study area

Income Level (BDT)	Respondents (No.)	Respondents (%)
Below-1,00,001	14	23.72
1,00,001-2,00,000	28	47.46
2,00,001-3,00,000	9	15.26
3,00,001-4,00,000	6	10.17
Above 4,00,0000	2	3.39

3.1.9 Drinking Water Facilities

The provision of clean and safe drinking water is regarded as one of the most valuable aspects of a society. The study showed that 100% of fish farmers household used tube-wells for drinking water [21].

3.1.10 Sanitary Facilities

It was observed that all respondents had sanitary facilities. From the survey, it was found that 49.15% of the farmers were used *paka* sanitary, whereas 6.78% of them used *kacha* (Figure 8). The fish farmers' sanitary conditions were better than those of the rice-fish farmers in Mymensingh district [22].



Figure 8. Sanitary facilities of the selected fish farmers in the study area

3.1.11 Electricity Facilities

In this digital era, one of the visions of the Bangladesh Government is to provide electricity to every household. All the respondents of the surveyed area had electricity facilities whereas Ali *et al.*[19] found it to be only 62%.

3.1.12 Organizational Membership

From the present study, it was found that 47.446% farmers were member of organization such as MBSK, GBK and 52.54% were not involved with any organization (Table 3). These findings were more or less similar to the findings of Ali *et al.* [19].

Table 3. Number and percentage of organizational membership of selected farmers

Organization Name	Respondents (No.)	Respondents (%)
MBSK	8	13.56
GBK	20	33.9
NO	31	52.54

3.2 Farm Related Information

3.2.1 Size of Pond

The size of the ponds is an important factor manipulating the use of inputs in the fish pond. The sample ponds were grouped into five categories depending upon their different sizes in the surveyed area. The ponds (27.12%) were dominated by those with a water spread area of 0.31-0.40 ha, (Figure 9). As per Saha *et al.* [23] the size of the ponds ranged from 0.05 to 0.15 ha which is lesser than the present reports.




Figure 9. Pond size of the selected fish farmers in the study area

3.2.2 Pond Ownership

According to the study, 59.33% farmers owned their ponds, while 40.67% leased them. And there was 93.22% single ownership and 6.78% multi-ownership (Figure 10). This statement is confirmed by Pravakar *et al.* [24].




Figure 10. A graphical presentation of pond ownership of farmers

3.2.3 Average Depth

It was observed that the average water depth of ponds were 1.7 m, whereas 50.85% were within 1.82-2.13 m and 49.15% ponds were 1.21-1.52 m (Figure 11). This statement was affirmed by Jahan *et al.* [25].




Figure 11. Average depth of ponds in the selected areas

3.2.4 Water Source

The water capacity of the pond is meager. In the selected study area, 79.66% of farmers used ground water and the rest of 20.34% used surface water whereas Ahmed [26] found it to be 55 % only (Figure 12).




Figure 12. A graphical presentation of water source of farmers

3.2.5 Water Exchange

It was observed that, 45.76% of the farmers do not exchange water. Among them, 50.84% exchange water when required, 1.69% weakly and 1.69% on monthly (Figure 13). These findings were a bit different from that of Ali *et al.* [27].




Figure 13. Water exchange by fish farmers of the selected areas

3.3 Culture Related Information

3.3.1 Pond Preparation

The selected farmers followed semi-intensive culture system for carp fattening. Before culture, 57.63% of farmers prepared their ponds. Among them, 26.47% farmer prepared dike, 97% cleaning weed, liming and rotenone used 94.12% and 52.95%. Very few farmers, about 14.70%, were drying their pond before culture where 42.37% did not prepare their pond before culture (Table 4). The guidelines for good aquaculture practice aspects were followed after [3].

Table 4. Preparation of pond before carp fattening

Respondents (%)		
Yes (57.63)		No (42.37)
Dike	26.47	
Weed cleaning	97	
Liming	94.12	
Rotenone	52.95	
Drying	14.70	

3.3.2 Use of Lime

All the participants of the study area used lime. The amount of lime varies with the duration and condition of water in the pond (Table 5).

Table 5. Use of lime in carp fattening ponds

Amount (kg/ha)	Interval (days)	Respondents (No.)	Respondents (%)
123.5	30	26	44.07
61.75	15	14	23.74
49.4	15	10	16.94
Required amount	When required	9	15.25

3.3.3 Salt

Salt is used to treat bacterial gill infections, controlling many external parasites, protects from fungus spores in water and relieve stress during handling and transport. In the winter season, they used it. 123.5 kg/ha salt used one-month interval.

3.3.4 TSP

TSP was used in 12.35-44.5 kg/ha in seven days intervals. Keep the TSP into the water with an oilcake and use urea before serving where Hossain *et al.* [28] stated it to be 635 kg/ha/month) to enhance the natural feed.

3.3.5 Urea

To increase pond carrying capacity, off-farm inputs such as chemical fertilizers and supplementary feeds are required. So that vast amounts of nitrogen (urea) are used to increase the growth of fish during the short period. The farmers in the selected study areas used urea 12.35-75.25 kg/ha in a seven days interval.

3.3.6 Growth Promotor

Growth promoters were chemical and biological substances which were added to fish's food with the aim to improve the growth of fishes. G.P.A, Aquamix, Cemovit, at a dose of 7-8 g/kg feed and Silver meal 2 ml/kg was used by the farmers, 40.16% was used Silver meal, GPA - 13.60%, Aquamix - 23.72% and Chemovit - 5.08%. And 16.92% used no growth promotor (Figure 14).




Figure 14. Growth promotor used by farmers of the selected areas

3.3.7 Species

Species selection for carp fattening based on diversified feeding habits, including surface, column and bottom feeder fish species. Catla (*Catla catla*), mrigal (*Cirrhinus cirrhosis*), rui (*Labeo rohita*), grass carp (*Ctenopharyngodon idella*), bighead (*Hypophthalmichthys nobilis*), silver carp (*Hypophthalmichthys molitrix*), mirror carp (*Cyprinus carpio*) were considered for carp fattening. With these species, some farmers culture tilapia (*Oreochromis niloticus*), pangas (*Pangasius pangasius*) as well as shing (*Heteropneustes fossilis*) which is encouraged by Bhanu *et al.* [29] and Azad *et al.* [30]

3.3.8 Density

Farmers in the study area maintained fish density by utilizing sufficient natural food and allowing fish to move freely. Fish were stocked based on feeding habits and the objective of proper feed use (Table 6).

Table 6. Stocking density of fish in the selected study area

Species	Feeding habit	Density/ha
Catla	Surface feeder	247-494
Silver	Surface feeder	124-247
Grass carp	Surface, column and marginal	247-494
Bighead	Surface feeder	124-247
Rui	Column feeder	494-980
Mrigal	Bottom feeder	494-1100
Mirror carp	Bottom feeder	247-770

3.3.9 Weight of Species

Farmers considered a variety weight of fish seed for culture. It is believed that large size seeds grow up more rapidly than others (Table 7).

Table 7. Weight of fish species considered for culture

Fish species	Weight (g)
Rui	250-500
Mrigal	250-500
Catla	500-1000
Silver	350-500
Bighead	350-500
Grass carp	350-1000
Mirror carp	250-350

3.3.10 Price of Seed

The price of carp seed varied from farmer to farmer. Generally, it fluctuates due to availability, season and location (Table 8).

Table 8. The price of fish seeds used for carp fattening

Species	Price of seed (Tk/kg)
Rui	175-200
Mrigal	175-200
Catla	175-200
Silver	90-110
Bighead	90-110
Grass carp	170-200
Mirror carp	150-180

3.3.11 Disease

The survey found that among 59 respondents, 51 respondents faced the lice problem. Few farmers, only four, said about bacterial and fungal disease problems. For bacterial disease, they use Chemistpro, Pond Care (S.K.F.). Thiovit (Syngenta) was used for fungal infection. After one month of stocking, they used medicine at 500ml/ha at a 15 days interval to control lice. A significant portion of the study areas farmers at 55.93% used Killer Argulouse (Anjum), Deltrix (Fishtech) used 25.42% and 6.78% Sumithion (Figure 15).




Figure 15. Medicines used for control of fish lice by farmers

3.3.12 Problems

Most of the farmers of the survey areas were facing technical and social problems. The main problems are lack of capital, high price of quality fish seeds and feeds, less protein in feed, poor technical knowledge, lack of social awareness etc. It was observed that 89.83% of farmers of the study areas were at problem. Among them, 66.03% faced financial and 50.85% social issues where 20.75% were in technical problems. Only 10.17% had no problem (Figure 16). Similar issues were also confronted by Mohsin [31], Habib *et al.* [32] and Saha [33] as well.




Figure 16. A graphical presentation of problems faced by farmers

3.4 Feed and Feeding Information

This section deals with the findings related to the feeds found in the study. The selected characteristics were fish feed used by farmers, feed types, feeding frequencies, feed application methods, feed storage facility, name of feed companies, etc.

3.4.1 Use of Feed by Farmers

In this study area, all the respondents used different types of feed in the ponds which connects to findings of Islam *et al.* [34].

3.4.2 Feed Types

Farmers of the study areas used ready or commercial; farm made and commercial+farm made feed. Among them, 40.68% used commercial feed, which indicates the maximum number of farmers depend on commercial feed. Only 20.34% of farmers use commercial +farm made feed (Figure 17).




Figure 17. Feeds used by farmers of the selected areas

3.4.3 Commercial Feed Company

A total of forty out of 59 people used commercial feed from various companies. The most common commercial feeds were Aci godrej, Quality, Narish, Tongwei, Meghna, Purobi and Aman. Aci godrej used by 55% farmers (Table 9).

Table 9. Feed from different companies used by farmers

Companies Name	Respondents (No.)	Respondents (%)
Aci godrej	22	55
Quality	8	20
Narish	4	10
Tongwei	2	5
Meghna	2	5
Purobi	1	2.5
Aman	1	2.5

3.4.4 Feeding Frequency

Farmers were applied feed in the morning, noon and afternoon. It was found that 81.35% of the farmers supplied feed twice a day, morning and afternoon. On the other hand, 18.65% farmers supplied feed thrice daily, in the morning, noon and the evening which resembles to the findings of the other researchers [35] (Figure 18)




Figure 18. A graphical presentation of feeding frequency

3.4.5 Feeding Method

It was found that in the selected study area, all the farmers applied feed manually.

3.4.6 Use of Vehicle

In the selected study area, it was found that 33.89% of farmers used vehicles to apply feeds to fish and 90% used nauka (boat) and the rest used other types. The significant portion, which was about 66.11% had no vehicles (Figure 19).




Figure 19. Vehicle use by farmers of the selected study area

3.4.7 Storage Facilities of Feed

It was found that 30.51, - 37.29 and 10.17% of the farmers had *paka*, semi-*paka*, and *kacha* feed storage facilities and the rest of 22.03% farmers had no storage facility which resembles many other research works [25] (Figure 20).




Figure 20. Storage facilities of feed in the selected study area

3.4.8 Marketing Live Fish

In the study areas, 72.89% of farmers were marketing fish as live. About 27.11% were not marketing live fish (Figure 21).




Figure 21. Marketing of live fish by fish farmers of the selected are

4. CONCLUSION

Carp polyculture is an important fish production technology because of the suitability of carps for stocking with other aquaculture species and high market demand as well as price. Among the polyculture systems, carp fattening is a relatively new and the latest technology having very high productivity. Due to different problems like lack of capital, high price of inputs, poor technical knowledge, social problems etc the culture method is not so popular and successful so far in the study areas. In spite of different problems and constraints, carp fattening is a potential system offering opportunities to increase fish production as well as income of the aquafarmers because of the high price of large fish produced by applying the technology. Therefore, the government should take the necessary steps to overcome the drawbacks and hindrances to the sustainability of the fish-producing technology that can improve the livelihood of the poor farmers, and play significant roles in the economy of the country.

CONSENT

According to the international standard, participants written consent has been collected and preserved by the author(s).

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Shamsuzzaman MM, Mozumder MMH, Mitu SJ, Ahamad AF, Bhyuian MS. The economic contribution of fish and fish trade in Bangladesh. *Aquaculture and Fisheries*. 2020; 5(4):174-181.

2. Ahmed F, Monsur A, Eyasmin S, Mitu MP, Rahman T. Bangladesh Economic Review 2021. Finance Division, Ministry of Finance Government of the People's Republic of Bangladesh, Bangladesh Secretariat, Dhaka-1000, Bangladesh. 2021.
3. DoF. National fish week compendium (in Bengali). Department of Fisheries, Ministry of Fisheries and Livestock, Bangladesh. 2012.
4. DoF. National Fish Week 2019 Compendium (in Bengali). Department of Fisheries, Ministry of Fisheries and Livestock, Bangladesh. 2019.
5. BBS (Bangladesh Bureau of Statistics). Statistical Yearbook of Bangladesh. Bangladesh Bureau of Statistics. Statistics Division, Ministry of Planning. 2020.
6. DoF. National Fish Week 2017 Compendium (in Bengali). Department of Fisheries, Ministry of Fisheries and Livestock, Bangladesh. 2017.
7. Ellis F. Rural livelihoods and diversity in developing countries. Oxford University Press. 2000.
8. Hasan MR, Ahmed GU. Issues in carp hatcheries and nurseries in Bangladesh, with special reference to health management. Food and Agriculture Organization Fisheries Technical Paper. 2002; 147-164.
9. Zafar MA, Mia MF, Parvez I, Kibria ASM, Roy KC. Present status of aquaculture practices in some selected areas of Dinajpur district. Research in Agriculture Livestock and Fisheries. 2020; 7(2): 341-349.
10. DoF. National Fish Week 2021 Compendium (in Bengali). Department of Fisheries, Ministry of Fisheries and Livestock, Bangladesh. 2021.
11. Sarma PK, Raha SK, Jorgensen H. An economic analysis of beef cattle fattening in selected areas of Pabna and Sirajgonj districts. Journal of the Bangladesh Agricultural University. 2014; 12(1): 127-134.
12. Ferdoushi Z. Crab fattening in Bangladesh: a socio-economic perspective. Journal of Environmental Science and Natural Resources. 2013; 6(1): 145-152.
13. Alam MJ, Kohinoor AHM, Islam MS, Mazid MA. Polyculture of carps using overwintered fingerlings under different stocking densities. Bangladesh Journal of Fisheries Research. 2002; 6(2): 117-124.
14. Jobling M. Are compensatory growth and catch-up growth two sides of the same coin. Aquaculture International. 2010; 18(4): 501-510.
15. Grover JH, Islam MA, Shah WA, Rana MAH, Chowdhury HA. Training manual for extension personnel on low cost environment friendly sustainable aquaculture practices. ICLARM, Dhaka, Bangladesh. 2000.

16. Talukder MGS, Hossain MA, Mohsin ABM, Khan RI. Performances of bottom dwelling carps in polyculture ponds under drought prone barind area of Bangladesh. *Journal of Aquaculture and Marine Biology*. 2018; 7(1): 00178.
17. Rana MS. An economic analysis of pond fish culture in some selected areas of Sirajgonj district. MS Thesis, Department of Agricultural Economics, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh. 1996.
18. Rahman MM. Socio-economic aspects of carp culture development in Gazipur, Bangladesh. MS Thesis, Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh. 2003.
19. Ali MH, Hossain MD, Hasan ANGM, Bashir MA. Assessment of the livelihood status of the fish farmers in some selected areas of Bagmara upazilla under Rajshahi district. *Journal of the Bangladesh Agricultural University*. 2008; 6(2): 367-374.
20. Alam MS, Flowra FA, Salam MA, Kabir AKMA, Ali H. Fishing gears, fish marketing and livelihood status of the poor fisherman around the Basantpur beel at Lalpur Upazila. 2009; *J. Agrofor. Environ.* 3 (1): 173-177.
21. Asif A, Habib MAB. Socio-economic condition of fish farmers of Jhikargachha upazila in Jessore district, Bangladesh. *Asian Journal of Medical and Biological Research*. 2017; 3(4): 462-475.
22. Podder, M. 2005. Socio-economic aspects of rice-fish culture development in some selected areas of Mymensingh district. MS Thesis. Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh. 50 pp.
23. Saha NC, Islam MS, Saha JK, Modak PC. Economics of pond fish production in some selected areas of Bangladesh. *Bangladesh Journal of Aquaculture*. 1995; 17: 13-18.
24. Pravakar P, Sarker BS, Rahman M, Hossain MB. Present status of fish farming and livelihood of fish farmers in Shahrasti upazila of Chandpur district, Bangladesh. *American-Eurasian Journal of Agricultural and Environmental Science*. 2013; 13(3): 391-97.
25. Jahan H, Parvez I, Kibria ASM. Current aquaculture practices in Dinajpur District: Special emphasis on fish feeds. *International Journal of Fisheries and Aquatic Research*. 2018; 3(1): 35-42.

26. Ahmed, J. 2010. Attitude of farmers towards the effect of pond ownership on fish production. M. S. Thesis. Department of Agricultural Extension, Bangladesh Agricultural University, Mymensingh.
27. Ali MM, Hossain MB, Rahman M, Rahman S. Post-stocking management practices by the pond fish farmers in Barisal district, Bangladesh. *Global Veterinaria*. 2014; 13(2): 196-201.
28. Hossain ME, Khan MA, Saha SM, Dey MM. Economic assessment of freshwater carp polyculture in Bangladesh: Profit sensitivity, economies of scale and liquidity. *Aquaculture*. 2022; 548: 737552.
29. Bhanu Prakash CH, Khairnar SO, Mandal A, Kumar A, Kumar B. Composite fish farming: A review on economic enterprise for rural empowerment and livelihood generation. *International Journal of Fisheries and Aquatic Studies*. 2018; 6(4):545-550.
30. Azad MAK, Rahman MR, Rahman Z, Kader MA, Haque MM, Alam MJ. Polyculture of carp, tilapia and pangas using low cost inputs. *Pakistan Journal of Biological Sciences*. 2004; 7(11): 1918-1926.
31. Mohsin ABM, Islam MN, Hossain MA, Galib SM.. Constraints and prospects of carp production in Rajshahi and Natore districts, Bangladesh. *University Journal of Zoology, Rajshahi University*. 2012; 31: 69-72.
32. Habib MAB, Rashiduzzaman M, Molla AR, Hasan MR, Begum A. A survey on socio-economic and technical aspects of pond fish culture in Bangladesh. In *Socio-economics of Aquaculture, Proceedings of International Symposium*. 1994; 93: 265-274.
33. Saha MK. A study on fish production technology in North-west Bangladesh. MS Thesis, Department of Aquaculture, Bangladesh Agricultural University, Mymensingh. 2003.
34. Islam FK, Asif AA, Ahmed M, Islam MS, Sarker B, Zafar MA, Rahman M. Performances of resource poor households in aquaculture practices in sadar upazila, Meherpur, Bangladesh. *International Journal of Fisheries and Aquatic Studies*. 2017; 5(6): 281-288.
35. Rahman MH, Zafar MA, Hossain MA, Kibria ASM. Present status of integrated aquaculture in some selected areas of Nilphamari district in Bangladesh. *International Journal of Fisheries and Aquatic Studies*. 2018; 6(2): 290-295.