

## **Interest of percutaneous vertebroplasty in the management of spinal metastases**

### **Abstract**

Vertebroplasty techniques, especially kyphoplasty, are becoming an increasingly important treatment option for spinal metastases. Recent studies have shown the effect of vertebroplasty on postoperative pain of metastatic origin without prolonged follow-up

The main objective of our work is to confirm the short-term effect of vertebroplasty in the treatment of pain related to vertebral metastases and to study the effect on vertebral stability.

We conducted a single-center retrospective study over a period of 4 years of 30 patients with spinal metastases without neurological signs, a total of 46 vertebroplasty.

The mean age of our patients was 55.13 years. The average preoperative visual analgesia scale (VAS) was 8.46 and reached an average of 2.73 maintained until the last recoil of 2 years postoperatively, attesting to the significant decrease in spinal pain ( $p < 0.05$ ).

Beck's index went from a mean of 0.58 preoperatively to a mean of 0.84 postoperatively, with a significant difference ( $p < 0.05$ ).

Angular kyphosis decreased from a mean of  $17.8^\circ$  preoperatively to  $7.3^\circ$  postoperatively at the last recoil with a significant difference ( $p < 0.05$ ).

Three cases of polymethacrylate (PMMA) cement leakage were noted, with no neurological compromise.

Spinal vertebroplasty techniques are reliable, effective and reproducible in the management of vertebral metastatic lesions, in its two main components, analgesia and stabilization.

**Key words: metastasis - spine - vertebroplasty**

## **Introduction:**

Metastatic localizations are mainly observed in lungs; liver and bone [1]. When it comes to bony metastasis the spine is most frequently involved [2]. These metastatic locations manifest essentially with invalidating pain deteriorating the quality of life. Pathological fractures associated with neurological compromise may also occur [3].

Vertebroplasty is a minimally invasive treatment consisting in injecting the vertebral body with Poly-Methyl-Methacrylate (PMMA) obtaining pain relief and restoration of vertebral height at the same time [4-5].

The aim of this study was to evaluate the outcome of vertebroplasty in the treatment of metastatic spine lesions: Pain and mechanical stability wise.

## **Methods:**

We lead a monocentric retrospective study over a period of 4 years including 30 patients with vertebral metastasis. Patients included presented severe back pain resistant to traditional therapies. Patients with neurological compromise were excluded from this study because spinal decompression surgery was also indicated. Diagnostic criteria for spinal metastasis were based on clinical examination and neuro-imaging findings. In some cases a biopsy was underwent to confirm the diagnosis. The procedure was lead under general anesthesia. Patient in a prone position. Injection of cement was realised under fluoroscopy until the lesion was fully opacified or if the cement reached the posterior wall. In case of loss of vertebral height we proceeded with a kyphoplasty: A technique allowing the restoration of vertebral height by dilating the vertebral body via a balloon catheter. All patients received post-operative pain relief and neurological surveillance was lead for 12 hours. Physical exam was conducted pre operatively; at 1 Month, 6 months 1 year and 2 years post-surgery. Post-operative evaluation was based on the visual analogue scale (VAS). Radiological findings collected and evaluated were the followings: location of the lesion; number of lesions; type of lesion (lytic, sclerotic or mixed); vertebral height through the Beck index and the local kyphotic deformity [6]. Spinal stability was analysed via the Kostuik score pre and post operatively. A score of three or more is in favour of instability while a score of less than 3 is in favour of spinal stability [7].

The data collected was analysed using the SPSS (The Statistical Package for the Social Sciences) software.

## **Results:**

A total of 46 vertebroplasty were performed in the period between 2015 and 2018 on 30 patients. The Patient group was composed of 12 males and 18 females and the mean age was 55.13 with extremes from 20 to 66. Primary cancers were Multiple myeloma (n=12) lung cancer (n=8), breast cancer (n=6) renal carcinoma (n=2) and

papillary thyroid carcinoma (n=2). The average number of lesions per patient was 1.53 with an extreme up to 3 levels. Among our patients lesions were found in all regions of the spine mainly the thoracolumbar spine (Table 1)(Figure 1). No complications related to anesthesia or to the procedure were noted. None of our patients had a local recurrence. The mean pre op VAS was at 8.46 with a mean of 2.73 in the post-operative evaluation. This difference was found to be statistically significant. All of our patients stopped taking group 3 analgesics for their back pain. Three of our patients presented with cement leakage with no neurological compromise (figure 2).

Post-operative imaging showed an 80% restoration of the anterior wall height in all of our patients ; the beek index going from a mean of 0.58 to a mean of 0.84 post operatively which is statistically significant with a  $p < 0.05$ .

Local kyphosis went from a mean of 17.8 to a mean of 7.3 degrees; this difference was also found to be statistically significant with a  $p < 0.05$  in favour of the effect of vertebroplasty in reducing the vertebral deformity. (Table 2)

The post-operative evaluation according to the Kostuik score showed a score of less than 3 in 91% of the vertebrae compared to 82 % with a score of 3 or more in the preoperative evaluation.

## **Discussion:**

Management of spinal metastasis gained a lot of interest with the extension of life expectancy of cancer patients.

The Main symptom of spinal metastases is a debilitating back pain with a high functional impact and resistance to the usual analgesic treatments.

In other cases; the lesions could be asymptomatic and being discovered only at the stage of complications such as spinal cord compression or pathological fractures [8] .

Non-surgical treatments seem to have very limited results and radiotherapy although it showed short term success when it comes to improving the quality of life, it has many side effects and cannot prevent pathological vertebral fractures with risk of spinal cord compression [9].

Vertebroplasty techniques which have recently gained interest in treatment of secondary lesions seem to be a reliable surgical alternative in lowering pain levels due to spinal metastasis: the reduction of the VAS between the pre and post-operative assessment in our study correlated significantly with the published data [10]. In terms of histological findings an antitumoral effect of PMMA has been proposed: This effect is related to the heat production during the polymerization process destroying the local micro circulation. This exothermic reaction may also be involved in the intra osseous nociceptive neural ablation explaining in part the analgesic effect of cementoplasty [11]. In radiological findings: the kyphoplasty technique allows a restitution of the vertebral height and reduction of the vertebral kyphotic deformity [12].

PMMA cement leakage is a considerable complication especially if the vertebral wall was disrupted [13]. A meticulous technique may prevent this complication.

**Conclusion:**

Vertebroplasty techniques seem to be reliable, reproducible and effective when it comes to the management of pain and instability related to spinal metastasis. A possible antitumor effect reinforces the interest in these techniques. This rising interest is also explained by the low incidence of complications related to vertebroplasty.

**CONSENT**

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

**ETHICAL APPROVAL**

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

UNDER PEER REVIEW

### Tables and figures List:

Table 1: Result analysis of 30 patients with spinal metastasis treated with vertebroplasty

<b>Male</b>	<b>36%</b>
<b>Female</b>	54%
<b>Mean Age</b>	55.13
<b>Age</b>	
<b>&lt;30</b>	1
<b>30-39</b>	2
<b>40-49</b>	14
<b>50-59</b>	9
<b>60-69</b>	4
<b>Primary cancer</b>	
<b>Myeloma</b>	12
<b>Pulmonary Carcinoma</b>	8
<b>Breast papillary carcinoma</b>	6
<b>Renal carcinoma</b>	2
<b>Thyroid papillary carcinoma</b>	2
<b>Number of levels treated</b>	46
<b>Cervical</b>	1
<b>Dorsal</b>	21
<b>Lumbar</b>	22
<b>Sacral (Figure 2)</b>	2
<b>Number of levels per patient</b>	
<b>1</b>	4
<b>2</b>	10
<b>3</b>	6
<b>4</b>	1
<b>Type of lesion</b>	
<b>Lytic</b>	41
<b>Sclerotic</b>	0
<b>Mixed</b>	5

Table 2: Comparison of pre and post-operative results.

	Pre-operative mean	Post-operative mean	The chi square test
<b>VAS</b>	8.46	2.73	$p < 0.05$
<b>Beck Index</b>	0.58	0.84	$P < 0.05$
<b>Angular kyphosis</b>	17.8 degrees	7.3 degrees	$P < 0.05$

Figure 1: Clinical case showing a vertebroplasty for a vertebral metastasis at L1 level





Figure 2: Clinical case showing a sacroplasty for a vertebral metastasis with cement leakage with no neurological compromise



**COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers

of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

### References:

1. Qian Z, Sun Z, Yang H, Gu Y, Chen K, Wu G. Kyphoplasty for the treatment of malignant vertebral compression fractures caused by metastases. *J Clin Neurosci*. 2011;18(6):763-7.
2. Mesfin A, Buchowski JM, Gokaslan ZL, Bird JE. Management of Metastatic Cervical Spine Tumors. *J Am Acad Orthop Surg*. 2015;23:38-46.
3. Dalbayrak S, Önen MR, Yılmaz M, Naderi S. Clinical and radiographic results of balloon kyphoplasty for treatment of vertebral body metastases and multiple myelomas. *J Clin Neurosci*. 2010;17:219-24.
4. Tseng Y-Y, Lo Y-L, Chen L-H, Lai P-L, Yang S-T. Percutaneous polymethylmethacrylate vertebroplasty in the treatment of pain induced by metastatic spine tumor. *Surg Neurol*. 2008;70:78-83.
5. Wardak Z, Bland R, Ahn C, Xie X-J, Chason D, Morrill K, et al. A Phase 2 Clinical Trial of SABR Followed by Immediate Vertebroplasty for Spine Metastases. *Int J Radiat Oncol*. 2019;104:83-9.
6. Afathi M, Mansouri N, Farah K, Benichoux V, Blondel B, Fuentes S. Use of Cement-Augmented Percutaneous Pedicular Screws in the Management of Multifocal Tumoral Spinal Fractures. *Asian Spine J*. 2019;13:305-12.
7. Amoretti N, Diego P, Amélie P, Andreani O, Foti P, Schmid-Antomarchi H, et al. Percutaneous vertebroplasty in tumoral spinal fractures with posterior vertebral wall involvement: Feasibility and safety. *Eur J Radiol*. 2018;104:38-42.
8. Ahn H, Mousavi P, Roth S, et al. Stability of the metastatic spine pre and post vertebroplasty. *J Spinal Disord Tech*. 2006;19:178-82.
9. Jawad MS, Fahim DK, Gerszten PC, Flickinger JC, Sahgal A, Grills IS, et al. Vertebral compression fractures after stereotactic body radiation therapy: a large multi-institutional multinational evaluation. *J Neurosurg Spine*. 2016;24:928-36.
10. Chi JH, Gokaslan ZL. Vertebroplasty and kyphoplasty for spinal metastases. *Curr Opin Support Palliat Care*. 2008;2:9-13.



11. Yang HL, Sun ZY, Wu GZ, et al. Do vertebroplasty and kyphoplasty have an antitumoral effect? *Med Hypotheses*. 2011;76:145-6.
12. Zou J, Mei X, Gan M, et al. Is kyphoplasty reliable for osteoporotic vertebral compression fracture with vertebral wall deficiency? *Injury*. 2010;41:360-4.
13. Chen JF, Lee ST. Percutaneous vertebroplasty for treatment of thoracolumbar spine bursting fracture. *Surg Neurol*. 2004;62:494-500

UNDER PEER REVIEW