

Original Research Article

A case series of cyclic vomiting syndrome in children at pediatric gastroenterology clinic in Benghazi medical

ABSTRACT

Background: Cyclic vomiting syndrome (CVS) is a functional disorder characterized by repeated episodes of sudden onset of intense vomiting that is followed by symptom-free periods. The diagnosis is based on the fulfillment of Rome IV criteria. The Cyclic vomiting syndrome shares similar features to migraine headaches. It is under-recognized and often misdiagnosed with significant delays in therapy.

Aims: To evaluate the clinical characteristic of pediatric patients diagnosed with cyclic vomiting syndrome in our institution and their response to treatment.

Patients and Methods: A prospective case series of 13 pediatric patients with cyclic vomiting syndrome seen between period of November 2018 to November 2020 at pediatric gastroenterology outpatient's clinic in Benghazi medical center. All relevant data including; age and sex, age of presentation, duration of symptoms, age at diagnosis, presence of aura or prodromal symptoms, patient or family history of headache, treatment and their response to treatment were collected.

Results: The mean age for diagnosis was 112.2 ± 37.7 months. 46.2% of patients had recognizable prodromal symptoms. Patients with CVS had intense nausea and persistent vomiting that requiring hospitalization. In 44.4% of patients had stopped the episodes as a response to propranolol.

Conclusion: Cyclic Vomiting Syndrome (CVS) is a functional, an under-recognized, and misdiagnosed episodic. The illness is characterized by stereotypical pattern of vomiting leading to frequent hospitalizations.

Keywords: Cyclic vomiting syndrome, Rome IV criteria, stereotypic vomiting, children.

1. INTRODUCTION

Cyclic vomiting syndrome (CVS) is an under-recognized and frequently misdiagnosed episodic illness. It is a chronic functional illness that can occur in both children and adults. And it is characterized by sudden episodes of intense nausea and vomiting which last for hours to days' alternate with symptom-free periods which last for weeks to months [1,2]. The symptoms of this illness can lead to repeated emergency visits, frequent hospitalizations, school absenteeism, and decreased quality of life. It is often misdiagnosed due to a lack of diagnostic tests and the unappreciated pattern of recurrence. The incidence of CVS is unknown to adult patients but is cited as 3%–14% [1] and affects between 0.04% – 2% of

pediatric patient populations [3, 4]. The recent population-based study reported a prevalence of up to 6.1% in children under 2 years old [5]. There is a slight predominance of girls over boys (86 VS 57%) [6] with a mean age of onset of 5 years[7] and the mean age at diagnosis is 8.2 to 9.5 years[8].

The etiology and pathogenesis of CVS are poorly understood, but it shares similar features with migraine headaches, it may be considered as a migraine variant. The International Classification of Headache Disorder (ICHD III beta) [9] considers CVS as a pediatric migraine variant. The Rome IV Criteria included CVS among the functional gastrointestinal illnesses [10-12]. Other several theories are under investigation, including autonomic [13] and hypothalamic-pituitary-adrenal axis hyper-reactivity[14], and mitochondrial dysfunction[15-17].

CVS is characterized by recurrent, stereotypic episodes of intense nausea and vomiting for a period ranging from hours to days that are separated by symptom-free intervals which last weeks to months [18]. 'Stereotypic' episode means that each episode is similar within individuals as to the time of onset, intensity, duration, frequency, and associated symptoms and signs. CVS typically has four phases: prodromal phase (pre-emetic), emetic phase, recovery phase, and asymptomatic or inter-episodic phase. Several patients have an aura in the prodromal phase lasting for several minutes and sometimes last up to hours, frequently consisting of abdominal pain, lethargy, sweating, or salivation. The emetic phase usually consists of intense nausea associated with frequent vomiting that tends to start in the early morning [19,20]. In the recovery phase, vomiting stops and appetite improves, and activity recovers then patients return to baseline health between episodes.

A number of CVS patients can identify the triggering conditions that can trigger the onset of the cycle of vomiting episodes. Triggers include negative and positive psychological stress or physical stress (infections, lack of sleep) the onset of menses which is also known as (catamenial CVS), and some food items (e.g. chocolate, cheese, monosodium glutamate) [19,21-23]

The diagnosis of CVS should be made after excluding other causes of vomiting and the typical clinical presentation as described by the Rome IV criteria. Rome IV criteria define CVS as stereotypic episodes of vomiting with the following characteristics: 1) presence of at least two acute episodes of vomiting in the preceding six months, 2) each episode of vomiting persisting for less than one week, 3) at least one week of an interval between episodes of vomiting, and 4) an absence of vomiting between episodes [24,25].

There is no curative treatment for CVS. Treatment goals are to reduce the severity and frequency of episodes and to improve the quality of life. Treatment includes identification and avoids triggers. During acute episodes, supportive therapy including IV rehydration and high-dose ondansetron is also recommended with psychological support to the patient and family. [24,25]

The NASPGHAN consensus statement recommendations of prophylactic therapy include the use of cyproheptadine (a first-generation antihistamine) or amitriptyline a tricyclic antidepressant as the first line. Propranolol (a beta-blocker) is used as the second line of prophylactic treatment. In patients with infrequent attacks, triptans should be considered as an abortive agent for those more than 12 years. [26, 27]

This study described the clinical characteristic of pediatric patients diagnosed with cyclic vomiting syndrome in Benghazi medical center and their response to treatment.

2. METHODOLOGY

A prospective case series study was conducted on patients with cyclic vomiting syndrome seen at the pediatric gastroenterology outpatient clinic in Benghazi medical center, during the period of November 2018 to November 2020. 13 pediatric patients fulfilling the Rome IV criteria were included.

2.1 Data Collection

Data was collected via the form created by the researcher. 13 of the patient medical records were reviewed and recorded data on the form. The form contained demographic data such as age and gender. In addition; clinical data such as laboratory tests were collected.) All relevant data including age and sex, age of presentation, duration of symptoms, age at diagnosis, presence of aura or prodromal symptoms, patient or family history of headache, treatment, and their response to treatment were collected.

2.2 Data Analysis

It was executed using the Statistics Package Social Science (SPSS) program version 20. Continuous variables were presented as means, median, standard deviation, and range. Categorical variables were reported with the use of proportions and percentages. Also, column charts and pie charts were used to describe and compare variables.


3. RESULTS

A total of 13 patients were diagnosed with CVS during the study period. Two third of them were male (61.2%). Almost all of the patients were older than 5 years in age. The mean age at onset of symptoms was 86.6 ± 38.9 months, whereas the mean age for diagnosis was 112.2 ± 37.7 months; as shown in Table 1. The duration of delay in diagnosing CVS mean 25.6 ± 17.7 months. 46.2% of patient reported to have recognizable prodromal symptoms which was abdominal pain in 7 patients (53.8%), headache with abdominal pain reported by 5 patients (38.5%) and excessive salivation reported by one patient (7.7%); as shown in Table 2.

Table 1. Characteristics of patients

Variable		No.	%
Gender	Male	8	61.5%
	Female	5	38.5%
Family History Migraine	Yes	8	61.5%
	No	5	38.5%
Age in months			
Mean \pm SD		118 \pm 34.1	
Median		120	
Minimum–maximum		66-163	
Age at onset of symptoms (months)			
Mean \pm SD		86.6 \pm 38.9	
Median		81	
Minimum–maximum		36-144	
Age at diagnosis (months)			
Mean \pm SD		112.2 \pm 37.7	
Median		120	
Minimum–maximum		63-163	
Number of previous admissions			
Mean \pm SD		8.5 \pm 12.3	
Median		3	
Minimum–maximum		0-40	

Positive family history of migraine in 8 patients (61.5%) and negative in 5 patients (38.5%). As Shown in Table 1 and Figure 1.

**Fig. 1. Family History Migraine**

The episodes of CVS were stereotypic among all patients with vomiting episodes for each patient last the same duration every time and each episode recurred at same interval each time in the same patient. Among our patients, interval between episodes of cyclic vomiting was 8.3 ± 5.7 months. The mean duration of each episode was 4.7 ± 1.3 days. As shown in Table 2.

Patients with CVS had intense nausea and persistent vomiting that requiring hospitalization, among our patient mean of admission to the hospital was 8.5 ± 12.3 times; as shown in Table 1. Four patients (30.7%) had infrequent episodes of CVS and they received supportive management only and they didn't receive any prophylactic treatment. Nine patients (69.2%) had too frequent CVS episodes, a prophylactic treatment with propranolol have been started for them. Only one patient (11.1%) didn't respond the prophylactic treatment with propranolol and shifted to topiramate, 4 patients (44.4%) the episodes stopped and 4 patients (44.4%) the frequency of the cycles reduced. As shown in Table 2 and Figure 2.

Table 2. Clinic data of patients

Variable	No.	%
Prodrome symptoms		
Yes	6	46.2%
No	7	53.8%
Symptoms during prodrome		
Salivation	1	7.7%
Abdominal pain	7	53.8%
Abdominal pain and headache	5	38.5%
Prophylactic		
No treatment	4	30.7%
Propranolol	9	69.2%
Response to treatment for 9 patients		
Cycles stopped	4	44.4%
Cycles frequency decreased	4	44.4%
Not respond to propranolol (other treatment)	1	11.2%
delay DX (months)		
Mean \pm SD	25.6 ± 17.7	
Median	24	
Minimum–maximum	2-48	
Duration of episode (days)		
Mean \pm SD	4.7 ± 1.3	
Median	5	
Minimum–maximum	3-7	
Inter-episode interval (months)		
Mean \pm SD	8.3 ± 5.7	
Median	8	
Minimum–maximum	2-24	




Fig. 2. Response to propranolol treatment

4. DISCUSSION

A total of 13 patients were diagnosed with CVS during the study period. Two third of them were male (61.2%) which is opposite to other reports in other studies that showed female were more affected than male [27,28,29]. Almost all of the patients in our study were older than 5 years in age which is consistent with other studies in different population [28,29,30].

The mean age at onset of symptoms was 86.6 ± 38.9 months (7.2 ± 3.2 years) which was similar to that reported by Khalil AF and Liao KY et al, [27, 29] whereas the mean age for diagnosis was 112.2 ± 37.7 months (9.3 ± 3.1 years). The mean interval from onset of symptoms to the accurate diagnosis was 25.6 ± 17.7 months (2.1 ± 1.4 years), which was comparable with that reported by Liao KY et al. [29]

46.2% of patient reported to have recognizable prodromal symptoms which was more than that reported in patients of other studies (25–38%), [31, 28] And less than reported by Khalil AF (two third of patients). [27]

Prodromal symptoms that reported were abdominal pain in 7 patients (53.8%), headache with abdominal pain reported by 5 patients (38.5%) and excessive salivation reported by one patient (7.7%).

Positive family history of migraine in 8 patients (61.5%) which seemed higher than that reported by Ertekin V et al and Ziqing Ye et al (29.2% and 11.9%). [28, 30]

The episodes of CVS were stereotypic among all patients with vomiting episodes for each patient last the same duration every time and each episode recurred at same interval each

time in the same patient. Among our patients, interval between episodes of cyclic vomiting was 8.3 ± 5.7 months. The duration of attack ranged from 3 to 7 days (median: 5 days) which was similar to that reported by Ertekin V et al. [28]

Patients with CVS had intense nausea and persistent vomiting that requiring hospitalization, among our patient mean of admission to the hospital was 8.5 ± 12.3 times. 4 patients (30.7%) had infrequent episodes of CVS and they received supportive management only and they didn't receive any prophylactic treatment. Nine patients (69.2%) had too frequent CVS episodes, a prophylactic treatment with propranolol have been started for them. Only one patient (11.1%) didn't respond the prophylactic treatment with propranolol and shifted to topiramate, 4 patients (44.4%) the episodes stopped and 4 patients (44.4%) the frequency of the cycles reduced. Khalil AF reported that the response rate to propranolol was 90.1% and similarly Haghighat et al in a large series reported that the response rate to propranolol was 92% without significant adverse effects. [27, 31,32].

5. CONCLUSION

Cyclic Vomiting Syndrome (CVS) is a functional, an under-recognized and misdiagnosed episodic illness characterized by stereotypical pattern of vomiting leading to frequent hospitalizations, multiple comorbidities, and poor quality of life. The treatment of CVS remains challenging. Pediatricians should be aware to this under-recognized illness and consider it in the differential diagnosis of recurrent episodic vomiting in children.

CONSENT

All respondents' parents who agreed to participate in the study were required to initialize the informed consent form.

ETHICAL APPROVAL

The principle of Medical Research Ethics guidelines from the Faculty of Medicine, University of Benghazi, were observed during this study

REFERENCES

1. Stanghellini V, Chan FK, Hasler WL, et al. Gastroduodenal disorders. *Gastroenterology*. 2016;150(6):1380–1392.
2. Li BU, Misiewicz L. Cyclic vomiting syndrome: a brain-gut disorder. *Gastroenterol Clin N Am*. 2003;32(3):997–1019.
3. Li BUK Cyclic vomiting syndrome and abdominal migraine. *Int Semin Pediatr Gastroenterol*. 2000; 9:1–9.
4. Abu-Arafeh I, Russel G. Cyclical vomiting syndrome in children: a population based study. *J Pediatr Gastroenterol Nutr*. 1995; 21:454–8.
5. Chogle A, Velasco-Benitez CA, Koppen IJ, Moreno JE, Ramírez Hernández CR, Saps M. A population-based study on the epidemiology of functional gastrointestinal disorders in young children. *J Pediatr*. 2016; 179:139–43.
6. Lin YP, Ni YH, Weng WC, Lee WT. Cyclic vomiting syndrome and migraine in children. *J Formos Med Assoc*. 2011;110(6):382–7

7. Li BU, Murray RD, Heitlinger LA, Robbins JL, Hayes JR. Heterogeneity of diagnoses presenting as cyclic vomiting. *Pediatrics*. 1998;102(3 Pt 1):583–7.
8. Lagman-Bartolome AM, Lay C. Pediatric migraine variants: a review of epidemiology, diagnosis, treatment, and outcome. *Curr Neurol Neurosci Rep*. 2015;15(6):34.
9. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia*. (2013) 33:629–808.
10. Benninga MA, Faure C, Hyman PE, St James Roberts I, Schechter NL, Nurko S. Childhood functional gastrointestinal disorders: neonate/toddler. *Gastroenterology*. (2016) 130:1519–26. doi: 10.1053/j.gastro.2005.11.065
11. Hyams JS, Di Lorenzo C, Saps M, Shulman RJ, Staiano A, van Tilburg M. Functional disorders: children and adolescents. *Gastroenterology*. (2016) 150:1456–68.
12. Doi.Brezin F, Wiedemann A, Feillet F. Cyclic vomiting syndrome in children. *Arch Pediatr*. (2017) 24:1129–36. doi: 10.1016/j.arcped.2017.08.010
13. Rashed H, Abell T L, Familoni B O. et al Autonomic function in cyclic vomiting syndrome and classic migraine. *Dig Dis Sci* 199944(8 Suppl)74S–78S.
14. Tache Y. Cyclic vomiting syndrome: the corticotropin-releasing-factor hypothesis. *Dig Dis Sci* 199944(8 Suppl)79S–86S.
15. Rinaldo P. Mitochondrial fatty acid oxidation disorders and cyclic vomiting syndrome. *Dig Dis Sci* 199944(8 Suppl)97S–102S.
16. Boles R G, Williams J C. Mitochondrial disease and cyclic vomiting syndrome. *Dig Dis Sci* 199944(8 Suppl)103S–107S.
17. Boles R G, Chun N, Senadheera D. et al Cyclic vomiting syndrome and mitochondrial DNA mutations. *Lancet* 1997350(9087)1299–1300.
18. Gee S. On fitful or recurrent vomiting. *St Bartholomew's Hospital Rep*. 1882; 18:1–6.
19. Fleisher DR, Gornowicz B, Adams K, Burch R, Feldman EJ. Cyclic vomiting syndrome in 41 adults: the illness, the patients, and problems of management. *BMC Med*. 2005; 3:20.
20. Levinthal DJ, Bielefeldt K. Adult cyclical vomiting syndrome: a disorder of allostatic regulation? *Exp Brain Res*. 2014;232(8):2541–2547.
21. Prakash C, Staiano A, Rothbaum RJ, Clouse RE. Similarities in cyclic vomiting syndrome across age groups. *Am J Gastroenterol* 2001; 96(3): 684-688
22. Pareek N, Fleisher DR, Abell T. Cyclic vomiting syndrome: what a gastroenterologist needs to know. *Am J Gastroenterol* 2007; 102(12): 2832-2840.
23. Chepyala P, Svoboda RP, Olden KW. Treatment of cyclic vomiting syndrome. *Curr treat options gastroenterol* 2007; 10(4): 273-282.
24. Zeevenhooven J, Koppen IJ, Benninga MA: The new Rome IV criteria for functional gastrointestinal disorders in infants and toddlers. *Pediatr Gastroenterol Hepatol Nutr*. 2017, 20:1-13. 10.5223/pghn.2017.20.1.1
25. Li BUK, Lefevre F, Chelimsky GG, Boles RG, Nelson SP, Lewis DW et al. North American Society for Pediatric Gastroenterology, Hepatology, and nutrition consensus statement on the diagnosis and management of cyclic vomiting syndrome. *J Pediatr Gastroenterol Nutr* 2008; 47:379–393
26. Donnet A, Redon S. Cyclic vomiting syndrome in children. *Curr Pain Headache Rep* 2018; 22:3
27. Khalil AF. Cyclic vomiting syndrome in children: a single-center experience. *Alex J Pediatr* 2020; 33:64-73
28. Ertekin V, Selimoğlu MA, Altınkaynak S. Prevalence of cyclic vomiting syndrome in a sample of Turkish school children in an urban area. *J Clin Gastroenterol* 2006; 40:896–898

29. Liao KY, Chang FY, Wu LT, Wu TC. Cyclic vomiting syndrome in Taiwanese children. *J Formos Med Assoc* 2011; 110:14–18
30. Ye Z, Xue A, Huang Y, Wu Q. Children with cyclic vomiting syndrome: phenotypes, disease burden and mitochondrial DNA analysis. *BMC Gastroenterol* 2018; 18:104
31. Haghighat M, Rafie SM, Dehghani SM, Fallahi GH, Nejabat M. Cyclic vomiting syndrome in children: experience with 181 cases from southern Iran. *World J Gastroenterol*. 2007;13(12):1833-1836.
32. Hilly M, Adams ML, Nelson SC. A study of digit fusion in the mouse embryo. *Clin Exp Allergy*. 2002;32(4):489-98.

UNDER PEER REVIEW