

# PHYTOCHEMICAL SCREENING AND *IN-VITRO* ANTIOXIDANT AND CYTOTOXIC EFFECTS OF *IPOMOEA BILOBA*

## ABSTRACT

*Ipomoea biloba* is a medicinal plant belonging to the Convolvulaceae family. It is an aquatic perennial runner plant used as a medical herb for various diseases such as asthma and rheumatism and dried leaves are used to apply for burns. In the present study, the Methanol extract of *Ipomoea biloba* leaves was experimented to evaluate the phytochemical properties, *invitro* antioxidant activity and *invitro* anticancer assay on MCF7 cell line. The leaves of *Ipomoea biloba* extract used in an analysis which shows the ability to produced estrogen in the form of estradiol via estrogen receptor in the cell cytoplasm. Phytochemical screening showed the presence of carbohydrate, amino acids, alkaloids, saponin, steroids, terpenoids and phenols. The antioxidant activity of Methanol extract indicates the significant antioxidant content and it's compared with standard ascorbic acid. The results of this study highlight the interest of *Ipomoea biloba* extract for the isolation of anticancer molecules.

**Key words:** Phytochemicals, Antioxidants, *Ipomoea biloba*, Cytotoxicity, Anticancer

## INTRODUCTION

*Ipomoea biloba* is a semi-aquatic tropical plant. *Ipomoea biloba* was mainly located on the west coast of India, bounded by Arabian Sea. The plant growth was extensive on the sand dunes near the shore “cyanodon dactylon” a gramiane member was also found growing luxuriously Arabian Sea (22). In the West Indies, a weak leaf decoction is a remedy for asthma and rheumatism. It is also drunk daily in the last month of pregnancy to promote an easy delivery (2).

Phytochemicals are chemical compounds formed during the plants normal metabolic process. Phytochemicals could also exhibit other bioactivities such as antimutagenic, anticancer, antioxidant, anticarcinogenic and anti- inflammatory properties (9). These chemicals are often referred to as “secondary metabolites (15). Most of the phytochemicals from plant

source such as phenolics and flavonoids have been reported to have positive impact on health and cancer prevention (13). High content of phenolic and flavonoids in medicinal plants have been associated with their antioxidant activities that plays a role in the prevention of the development of age-related disease particularly cause by oxidative stress(3). The plant derived natural antioxidants which are in the form of raw extracts constituents are very efficient to block the process of oxidation by neutralizing free radicals (21). A free radical is an individual molecule with more than one unpaired electrons. Free radicals lead to antioxidant shield ingestion, which can cause cell function interference and oxidative problems for the membranes (1). Antioxidants showed a substantial role in safeguarding the body against reactive oxygen damage (5).

Cancer is a group of diseases that cause cells in the body to change and grow out of control. Cancer begins with a genetic defect. Human's genetic factors, meaning genes, are located within the cell structures called chromosomes. Cancer develops when mutations take place in genes that control a cell's normal functions but which are simply damaged (6). According to the report of the International Agency for Research on Cancer of the World Health Organization published in 2014, the global incidence of cancer has been approximately 14 million new cases and is projected to register 19.3 million in 2025 (12). The most type cancer found is breast cancer. More than 3.1 million US women with a history of breast cancer were alive on January 1, 2014. While some are being undergoing treatment. In 2015, approximately 40,290 women are expected to die from breast cancer (19). Plants have contributed lot of medicinal compounds being used today to treat diseases like cancer, Hormonal imbalances, jaundice, diabetes, inflammation etc., (17). Treatment with medicinal plants is considered very safe as there is no or minimal side effects (11).

## **MATERIALS AND METHODS**

### **Plant Collection**

*Ipomoea biloba* were collected in and around Salem District, Tamilnadu, India was identified and confirmed. In Tamil it was known as "Hadapan kodi".

### **Preparation of Plant Extracts**

Fresh leaves were collected from the plants, washed and shade dried and powered. The powder was extracted using Methanol solvent by soxhlet apparatus. The residue was filtered and the solvent were evaporated under reduced pressure and stored for further studies. The extract was used for the determination of phytochemical constituents, *In-vitro* antioxidant and for *In-vitro* anticancer studies.

### **Phytochemical analysis**

The plant extract was assessed for the existence of carbohydrates, Protein, Aminoacids, Alkaloids, Flavonoids, Saponin, Steriods, Terpenoids and Phenols by the phytochemical analysis (screening) using typical standard methods .

### **Antioxidant Activity**

#### **Dpph Radical Scavenging Activity**

DPPH radical scavenging activity was carried out by the method of Molyneux 2004 (16). To 1 ml of 100 µm DPPH solution in Methanol, equal volume of the test sample in Methanol of different concentration was added and incubated in dark for 30 minutes. The change in coloration was observed in terms of absorbance using a spectrophotometer at 514 nm. 1 ml of Methanol instead of test sample was added to the control tube. Different concentration of ascorbic acid was used as reference compound. Percentage of inhibition was calculated from the equation:

$$[(\text{Absorbance of control} - \text{Absorbance of test}) / \text{Absorbance of control}] \times 100$$
. IC<sub>50</sub> value was calculated using Graph pad prism 5.0].

#### **Hydroxy Radical Scavenging Activity**

The hydroxyl radical scavenging activity of the test sample was estimated according to the method of Halliwell *et al.*,1992.(10) The hydroxyl radical was generated by a fenton-type reaction. The reaction mixture contained 2.0 ml of sample in varied 0.1 ml concentration to which, 0.1 ml EDTA (1mM), FeCl<sub>3</sub> (10mM) mixture, H<sub>2</sub>O<sub>2</sub> (10mM), 0.36 ml deoxyribose(10mM), 0.33 ml phosphate buffer (50mM, pH 7.4) and 0.1 ml of ascorbic acid (1mM) was added in sequence.

The mixture was incubated at 37°C for 1 hr. To this mixture was added 1.0 ml each of TCA (10%) and TBA (0.67%) and kept in boiling water bath for 20 minutes. The colour

developed was read at 532 nm. The control tube contains phosphate buffer, instead of sample. Different concentration of ascorbic acid was used as reference compound.

### **MTT Assay for Cell Cytotoxicity**

#### **Principle**

MTT (3-(4, 5 dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide) assay, is based on the ability of a mitochondrial dehydrogenase enzyme of viable cells to cleave the tetrazolium rings of the pale yellow MTT and form a dark blue colored formazan crystals which is largely impermeable to cell membranes, thus resulting in its accumulation within healthy cells. Solubilization of cells by the addition of detergents (DMSO) results in the liberation of crystals which are solubilized. The number of surviving cells is directly proportional to the level of formazan product created. The color can be quantified using a multi-well plate reader.

#### **Materials Required**

DMEM medium, Fetal Bovine Serum (FBS) and antibiotic solution were from Gibco (USA), DMSO (Dimethyl sulfoxide) and MTT (3-(4,5 dimethylthiazol-2-yl)-2,5-diphenyl tetrazoliumbromide) (5 mg/ml) were from Sigma, (USA), 1X PBS was from Himedia, (India). 96 well tissue culture plate and wash beaker were from Tarson (India).

### **PROCEDURE**

#### **Cell culture**

MCF-7 (Human breast cancer cells) cell line were cultured in liquid medium (DMEM) supplemented 10% Fetal Bovine Serum (FBS), 100 u/ml penicillin and 100 µg/ml streptomycin, and maintained under an atmosphere of 5% CO<sub>2</sub> at 37°C.

#### **MTT Assay**

The plant extract (*Ipomoea biloba*) was tested for *in vitro* cytotoxicity, using MCF-7 cells by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Briefly, the

cultured MCF-7 cells were harvested by trypsinization, pooled in a 15 ml tube. Then, the cells were plated at a density of  $1 \times 10^5$  cells/ml cells/well (200  $\mu$ L) into 96-well tissue culture plate in DMEM medium containing 10 % FBS and 1% antibiotic solution for 24-48 hour at 37°C. The wells were washed with sterile PBS and treated with various concentrations of the plant extract in a serum free DMEM medium. Each sample was replicated three times and the cells were incubated at 37°C in a humidified 5% CO<sub>2</sub> incubator for 24 h. After the incubation period, MTT (20  $\mu$ L of 5 mg/ml) was added into each well and the cells incubated for another 2-4 h until purple precipitates were clearly visible under an inverted microscope. Finally, the medium together with MTT (220  $\mu$ L) were aspirated off the wells and washed with 1X PBS (200  $\mu$ L). Furthermore, to dissolve formazan crystals, DMSO (100  $\mu$ L) was added and the plate was shaken for 5 min. The absorbance for each well was measured at 570 nm using a micro plate reader (Thermo Fisher Scientific, USA) and the percentage cell viability and IC<sub>50</sub> value was calculated using Graph Pad Prism 6.0 software (USA).

## RESULTS AND DISCUSSION

### Preliminary Phytochemical Screening

The results of preliminary qualitative phytochemical analysis on leaves of Methanol solvent extract of *Ipomoea biloba* were showed in (Table 1).

**Table 1: Preliminary phytochemical analysis of Methanol extract of *Ipomoea biloba***

S.NO	PHYTOCHEMICAL COMPOUNDS	PLANT EXTRACT
1.	Carbohydrate	+
2.	Protein	-
3.	Amino acids	+
4.	Flavanoids	-
5.	Alkaloids	+
6.	Saponin	+
7.	Steriods	+
8.	Terpenoids	+
9.	Phenols	+

Analysis of the plant extracts revealed the presence of phytochemicals such as phenols, tannins, flavonoids, saponins, glycosides, steroids, terpenoids, and alkaloids (4). Plant derived natural products such as flavonoids, terpenoids and steroids etc. pharmacological properties including antioxidant and anticancer activity. They possess biological properties such as antiapoptosis, antiaging, anticarcinogen, antiinflammation, antiatherosclerosis, cardiovascular protection and improvement of endothelial function, as well as inhibition of angiogenesis activities (18). Saponins have the property of precipitating and coagulating red blood cells. Cholesterol binding properties and bitterness (7).


Different concentrations ranging from 25-200 µg/ml of the Methanol extract of leaves of *Ipomoea biloba* were tested for their antioxidant activity in different *in-vitro* models. The percentage of inhibition was observed and found that the free radicals were scavenged by the test compounds in a concentration dependent up to the given concentration in all the models.

#### **DPPH Radical Scavenging Activity**

The activity of DPPH radical scavenging of the leaves extract was presented in Figure 1. The percentage of inhibition in DPPH in different concentration like respectively where as the percentage inhibition of ascorbic acid in concentration like 25, 50,75,100,200 µg/ml 32,45,55,68,72 respectively whereas the percentage inhibition of ascorbic acid in concentration like 25, 50,75,100, 200µg/ml were found to be 15, 20, 25, 42,54 respectively. The IC 50 values for DPPH scavenging activity for Methanol extract of leaves of *Ipomoea biloba* and ascorbic acid were 0.51 µg/ml and 0.93µg/ml respectively.

% scavenging =  $\frac{[\text{Absorbance of control} - \text{Absorbance of test sample}]}{\text{Absorbance of control}} \times 100$


**Figure 1: DPPH Radical Scavenging Activity of Methanol extract of leaves of *Ipomoea biloba***



### Hydroxyl Radical Scavenging Activity

The hydroxyl radical scavenging activity of plant extract was presented in Figure 2. Hydroxyl radicals were scavenging in different concentration like 25, 50, 75, 100, 150 µg/ml were observed in 12, 16, 22, 32, 42 respectively where as the percentage 10, 18, 27, 34, 45 inhibition of ascorbic acid in concentration like 25, 50, 75, 100, 150 µg/ml were found to be respectively. The IC<sub>50</sub> values for hydroxyl radical scavenging activity Methanol extract of leaves of *Ipomoea biloba* and ascorbic acid were 0.99 µg/ml and 0.96 µg/ml respectively. Values are the average of *In-vitro* experiments and represented as mean  $\pm$  standard deviation.

**Figure 2: Hydroxyl Radical Scavenging Activity of Methanol extract of leaves of *Ipomoea biloba***




There are studies have been carried out to evaluate the antioxidant activity of *Ipomoea biloba* species using DPPH assay and reported that, particularly *Ipomoea biloba* exhibited higher antioxidant activity .Whereby the present study proof that, the leaves extract of *Ipomoea biloba* has the potential compound(s) react as antioxidant which is suitable to develop a drugs for the prevention of human disease related to free radical mechanism (20).




## MTT Assay for Cell Cytotoxicity

Figure 3: Cell Viability (%)




IC<sub>50</sub> Value of tested sample: 80.70 µg/ml


**Figure 4: Images of control cells and Plant extract treated cells**



**Control cells**




**Plant extract 500 µg/ml**




**Plant extract 100 µg/ml**




**Plant extract 80 µg/ml**



**Plant extract 40 µg/ml**



**Plant extract 20 µg/ml**



**Plant extract 10 µg/ml**

MTT (3-(4, 5 dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide) assay, is based on the ability of a mitochondrial dehydrogenase enzyme of viable cells to cleave the tetrazolium rings of the pale yellow MTT and form a dark blue colored formazan crystals which is largely impermeable to cell membranes, thus resulting in its accumulation within healthy cells (Figure 3). Solubilization of cells by the addition of detergents (DMSO) results in the liberation of crystals which are solubilized. The number of surviving cells is directly proportional to the level of formazan product created. The color can be quantified using a multi-well plate reader (Figure 4).

MCF-7 cells are useful for *in vitro* breast cancer studies as a result of the cell line retaining several ideal characteristics particular to the mammary epithelium. These include the ability for MCF-7 cells to process estrogen in the form of estradiol via estrogen receptors in the cell cytoplasm. This results in the MCF-7 cell line being an estrogen receptor (ER) positive cell line. MCF-7 is also progesterone receptor positive and HER2 negative (8). *In vitro* anticancer studies has suggested that an anticancer effect of *Ipomoea biloba* extracts is possibly due to inhibition of DNA replication in cancer cell lines. It also reported anticancer property of *Ipomoea biloba* (14).

## CONCLUSION

The results obtained in this study thus suggest the identified phytochemical compounds, antioxidant properties and anti breast cancer activity may be the bioactive constituents of *Ipomoea biloba* are proving to be an increasingly valuable reservoir of bioactive compounds of substantial medicinal merit. This results in the MCF-7 cell line being an estrogen receptor (ER) positive cell line. MCF-7 is also progesterone receptor positive and HER2 negative.

Hence, the above plant extract could be explored for its highest therapeutic efficacy by pharmaceutical companies in order to develop safe drugs for various ailments.

**NOTE:**

*The study highlights the efficacy of "herb" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.*

## LITERATURE CITED

1. Andallu B, Varadacharyulu NC. 2003. Antioxidant role of mulberry (*Morus indica* L. cv. Anantha) leaves in streptozotocin-diabetic rats. *Clinica Chimica Acta* 338(1/2): 3-10.
2. Anna L.H. Dibiyantoro & G.H. Schmelzer, 3 March 2019, Plant Resources of South-East Asia.
3. Azwanida NN .(Azwanida, Med Aromat plants 2015, 4:3). A review on the extraction Methods US Medicinal plants, principle, strength and limitation
4. Babbar, N.; Oberoi, H.S.; Sandhu, S.K.; Bhargav, V.K. Influence of different solvents in extraction of phenolic compounds from vegetable residues and their evaluation as natural sources of antioxidants. *J. Food Sci. Technol.* 2014, 51, 2568–2575.
5. Baynes JW. 1991. Role of oxidative stress in development of complications in diabetes. *Diabetes* 40(4): 405-412.
6. Brown AM, Jeltsch JM, Roberts M, and Chambon P (2013) , Activation of pS2 gene transcription is a primary response to estrogen in the human breast cancer cell line MCF-7, 81 (20) 6344-6348.
7. Diaz-Gerevini, G.T.; Repossi, G.; Dain, A.; Tarres, M.C.; Das, U.N.; Eynard, A.R. Beneficial action of resveratrol: How and why? *Nutrition* 2016, 32, 174–178. [CrossRef] [PubMed] 109.
8. Fanelli, Alex (2016). "MCF-7 Cells: human breast adenocarcinoma cell line". Retrieved 3 December 2017.

9. Fung TT, Chiuve SE, Willett WC, Hankinson SE, Hu FB, Holmes MD. Intake of specific fruits and vegetables in relation to risk of estrogen receptor-negative breast cancer among postmenopausal women. *Breast Cancer Res Treat*. 2013
10. Halliwell *et al.*,1992. Antioxidants: The Basics-what they are and how to Evaluate them Advances in Pharmacology Volume 38, 1996, Pages 3-20
11. Hamta, A, Shariatzadeh, SMA, Soleimani Mehranjani, SMA, Fallah Huseini, H, Hosseinabadi, F. The cytotoxic effects of Glycyrrhiza glabra L. root extract on 4T1 cell line derived from BALB/c mice mammary tumors. *J Med Plants*. 2014;2:92–103.
12. Joensuu, Heikki; Jyrkkö, Sirkku; Kellokumpu-Lehtinen, Pirkko-Liisa; Kouri, Mauri; Roberts, Peter J. & Teppo, Lyly (toim.) (2013) *Syöpätaudit*. Helsinki: Kustannus Oy Duodecim.
13. Kerry Bone, Sion Mills -2013 Principles of Herbal Pharmacology. In principle and Practice of Phytotherapy 2013. DOI: <http://dx.doi.org/0.2013/b978>.
14. Manas Kotepui Diet and risk of breast cancer *Contemp Oncol (Pozn)*. 2016; 20(1): 13–19. Published online 2016 Mar 16. doi: 10.5114/wo.2014.40560 PMCID: PMC4829739 PMID: 27095934.
15. Mithraja Mj, Jounson M, Mahesh M, Miller Paul Z and Jeeva S (2012). Inter specific variation studies on the phyto- constituents of Christella and Adiantum using photochemical methods. *Asian Pacific Journal of Tropical Bio medici Azwanida NN* A review on the extraction Methods US Medicinal plants, principle, strength and limitation.(Azwanida, Med Aromat plants 2015, 4:3)ne,2(S1):S40-S45.
16. Molyneux, P. (2004) The Use of Stable Free Radical Diphenylpicrylhydrazyl (DPPH) for Estimating Antioxidant Activity. *Songklanakarin Journal of Science and Technology*, 26, 211-219.
17. Padmanadhan and Jungle, Das, 2008 Ajit Kumar Das, B K Dutta & GD Sharma Department of botany, SS college
18. Pu, F.; Ren, X.L.; Zhang, X.P. Phenolic compounds and antioxidant activity in fruits of six Diospyros kaki genotypes. *Eur. Food Res. Technol*. 2013, 237, 923–932.
19. Rick alteri, MD., Tracie Bertaut 2015 Marc.Lippman Breast cancer Research and Treatment Journey. Springer.com. ISSN : 0167-6806 Journal no. 10549

20. Sangetha S, Zuraini Z, Sasidharan S, Suriani S (2008). Free radical scavenging activity of *Cassia spectabilis* and *Cassia fistula*. Int. J. Nat. Eng. Sci., 2: 111-112.
21. Sharareh Rezaeian Almos Environ Impacts of antioxidants on Hydroxyl Radical Production from Individual and mixed Transition Metals in a surrogate Lung fluid(HHS public Access) 2015. Nov.45(40): 75555-7562 DOI: 10,1015
22. Yik Ling Chew, Elaine Wan Ling Chan, Pei Ling Tan, Yau Yan Lim, Johnson stanslas, JooKheng Goh. 2011. Assessment of phytochemical content, polyphenolic composition, antioxidant and antibacterial activity of Leguminosae medicinal plants in Peninsular Malaysia. Chew BMC Complementary and Alternative medicine 11:12.