

Properties and Effect of Fresh Concentrated Extract of Garlic on Different Bacteria and Fungi

ABSTRACT

Aims: Study some characteristics of fresh concentrated extract of garlic (FCEG) and analyze its effect on the growth of different bacteria and fungi.

Place and Duration of Study: Sample: Faculty of Chemical Sciences. Autonomous University of San Luís Potosí, S.L.P., between march and November 2020.

Methodology: To obtain the FCEG, 15 heads of raw garlic, previously peeled, were ground in a mortar and the suspension obtained was filtered through gauze, pressing to obtain a greater amount of filtrate, and kept covered at 4°C. For the study of some of their characteristics, the yeast *Candida albicans* was used. Petri dishes containing Sabouraud dextrose agar were inoculated with 1×10^6 yeasts/mL and 50 µL of FCEG, and were incubated at 28°C for 5 days, comparing growth with respect to a control without FCEG (all inhibition experiments described below were performed with the same protocol), while for the study of antifungal properties analyzed the effect of the extract on different strains of bacteria, yeasts, and fungi.

Results: Some of the factors that modify antifungal activity of FCEG are dilution, incubation temperature, protein concentration, half-life, capping and uncapping of tubes, and treatment with activated carbon. Regarding the growth inhibition analyses, it was found that all the species analyzed were susceptible to FCEG, among which the following stand out: the fungi *Histoplasma capsulatum*, the yeasts *Cryptococcus neoformans* and *C. albicans*, as well as different species of Dermatophytes, *Aspergillus*, and bacteria. **Conclusion:** FCEG shows a good antimicrobial effect against a wide variety of bacteria, yeast, and fungal species, which makes its application in medical therapy and agriculture possible, as well as being cheap, easy to obtain and does not cause side effects, although more studies are required for its therapeutic application.

Keywords: Garlic, properties, antimicrobial effect, bacteria, yeast, fungi

1. INTRODUCTION

Antimicrobial resistance (AMR) occurs in viruses, bacteria, fungi, and parasites as a manifestation inevitability of its capacities to evolve. The decreased effectiveness of antimicrobials existing is a consequence of the very complex interaction between natural selection, environment, and patterns of use of these drugs. This has given result that AMR has become a problem of global public health: as the variety of existing pathogens that affect humans and animals have developed various mechanisms to defend against available antimicrobials [1]. AMR is acquired through genes that code for resistance to antibiotics or through mutation or alteration of some gene. These mutations can be intrinsic or spread through communities microbial cells vertically (by inheritance) or in horizontally (by using elements mobile genes and extrachromosomal plasmids). This genetic diversity among populations microbial combined with a generation time very quickly, particularly among bacteria, provides microorganisms with a response extraordinary adaptation to the selective pressure of antimicrobials [2]. Therefore, are required approaches multidisciplinary and multisectoral like One Health or other related concepts such as Planetary Health or Eco Health that involve the disciplines of human, animal, and environmental health in a context of synergy of capabilities and experience. Also, a wire is required driver who can amalgamate the efforts of the different disciplines with an intersectoral and inter-disciplinary approach to optimize these sources allocated to AMR control, improve epidemiological surveillance, implement measures mitigation of its effects, as well as protection for the population at higher risk of acquiring multidrug-resistant infections [1, 3].

Other lines of research in antimicrobial antibiotics have focused on the use of some natural or modified plant extracts, and other natural compounds, with highly satisfactory results, such as: the effect of essential oils of oregano (*Oreganum compactum*), thyme (*Thymus vulgaris*), and rosemary (*Rosmarinus officinalis*) against *Aspergillus niger* [4], the study of the fungi toxic effects of *Allium sativum* (L) and *Ocimum gratissimum* (L.) using aqueous extraction methods on six fungal pathogens [5], a bioactive compound (trichodermin) isolated from the culture extracts of the fungus *Trichoderma brevicompactum*, and has a marked inhibitory activity on *Rhizoctonia solani* and *Botrytis cinerea* [6], the antileishmanial activity of a mixture of *Tridax procumbens* and *A. sativum* in mice [7], the bacterial effect of the extract of *Coriandrum sativum* on bacteria gastrointestinal pathogens [8], the antibacterial activity, of six essential oils and five ethanolic extracts from different Colombian plants, against Gram positive and Gram negative bacteria [9], the antifungal activity against *Cryptococcus neoformans* strains of *Euphorbia tirucalli* L. [10], and the effect of different aqueous extracts of plants on the in vitro mycelial growth of phytopathogenic fungi [11].

Another line of research in antimicrobial antibiotics has focused on the use of garlic, to which it is common to attribute a wide variety of properties, among which are: antifungal and antileishmanial activity [4, 7], the treatment of infections that does not affect the central nervous system [12], the in vitro anticoagulant effect of garlic extract on the blood coagulation cascade [13], evaluate the relative antidiabetic potential of garlic aqueous [14], and other medicinal properties such as: thorax, asthma, and skin diseases, blood disorder, hypertension, heart attack, antioxidant, etc., [15]. Regarding the antimicrobial antifungal properties of garlic, there are a wide variety of studies, among which are: the antifungal activity of some essential oils and their major phenolic components against *Aspergillus niger* [4], the in vitro effects of garlic (*A. sativum* L.) on fungal pathogens isolated from rotted Cassava roots [5], their antimycotic activity against *Trichophyton rubrum* [16, 17], *Alternaria tenuissima* [18], some Dermatophytes and *Candida albicans* [19, 20], *Histoplasma capsulatum* var. *capsulatum* [21], *Moniliophthora roreri* [22], *Aspergillus parasiticus* y *A. niger* [23], and the effect of garlic in the treatment of Saprolegniasis in rainbow trout (*Oncorhynchus mykiss*). Saprolegnia was long considered a fungus, but recently by consensus a new kingdom was created, thus separating it from the fungi kingdom and inserting it into the new kingdom: the Chromista Kingdom, which is much closer to algae than to fungi [24].

Too was studied their effect antimicrobial like against *Escherichia coli*, *Staphylococcus aureus*, and *A. niger* [25, 26, 27, and 28], *Streptococcus mutans* [29], molds, yeasts, coliforms, and *E. coli* [30] on the Gram positive bacteria *Bacillus subtilis* and *Staphylococcus epidermidis* and other Gram negative bacteria *E. coli* and *Shigella dysenteriae* [31]. The information consulted supports the health benefits associated with garlic consumption. The crushing of garlic bulbs allows obtaining alliin, which turns into allicin due to enzymatic oxidation. This compound has a fundamental role in the garlic's medicinal properties, and the garlic plant's activity depends on its ability to produce allicin. Preclinical studies have shown the improvement of the immune system, and specific proteins associated with its immunostimulating effect have been identified. In clinical studies, supplements with garlic derivatives have been administered to patients, and a decrease in the incidence of influenza and acute respiratory diseases has been observed [32]. But, there are few reports of its effect against environmental contaminating fungi, and of some of its properties such as: its half-life, storage temperature, the effect of different dilutions, and the minimum inhibitory concentration of the FCEG. Therefore, the objective of this work was to analyze some characteristics of FCEG, and their effect on the growth of different bacteria and fungi, in order, for in the future, obtain a product that is competitive with the common antimycotics used in the market.

2. MATERIAL AND METHODS

2.1 Obtaining of fresh extract of garlic (FCEG)

To obtain the FCEG, 15 heads of raw garlic (from the Republic market in the City of San Luis Potosí), previously peeled, are placed in an extractor and ground to obtain a garlic suspension, which is filtered through gauze, pressing the extract for better performance. Subsequently, it is transferred to an amber bottle and stored at 4°C.

2.2 Analysis of the properties of FCEG

For the study of the properties of the FCEG, the dimorphic fungus *C. albicans* was used, due to its ease in quantifying the number of yeasts, and its rapid growth in the medium used: Sabouraud Dextrose Agar (SDA).

2.2.1 Effect of different FCEG dilutions on the growth of *C. albicans*

100 µL of the FCEG are taken, and diluted with different volumes of sterile saline solution (SS) at 0.85% (w/v), according to the following Table:

Table 1. Dilution of the FCEG with saline solution

Dilution	FCEG	Saline solution
$\frac{\mu\text{L}}{\mu\text{L}}$	(μL)=	
Concentrate	50	0
1:10	100	900
1:100	10	990
1:1000	10	9990

Subsequently, aliquots of 50 µL are taken from each of the dilutions to be analyzed, and the antifungal effect is analyzed at 1×10^6 yeast/mL of *C. albicans*, incubating at 28°C, for 5 days hours in SDA, comparing growth with respect to a control without FCEG.

2.2.2 Effect of different FCEG concentrations on the growth of *C. albicans*

Different FCEG concentrations are taken (0-50 µL=0-1.0 mg/mL of protein) and diluted with different volumes of sterile saline solution (SS) at 0.85% (w/v) and were added 1×10^6 yeast/mL of *C. albicans*, incubating at 28°C, for 5 days in SDA, comparing growth with respect to a control without FCEG.

2.2.3 Effect of FCEG (50 µL= 1.0 mg/mL of protein) on the growth of different *C. albicans* concentrations

To different *C. albicans* concentrations (1×10^6 to 10×10^6 yeast/mL) spread in Petri dishes, they were added with 50 µL of FCEG, incubating at 28°C, for 5 days in SDA, comparing growth with respect to a control without FCEG.

2.2.4 Half-life of antifungal activity of FCEG to 4°C

The FCEG in covered containers was incubated at 4°C for 60 days, to determine the time in which it loses its antifungal properties, it was analyzed by sowing every 5 days 1×10^6 yeast/mL of *C. albicans* in SDA in the presence of 50 µL of FCEG, incubating at 28°C, for 5 days, comparing growth with respect to a control without FCEG.

2.2.5 Effect of tube capping on the antifungal activity of FCEG at 4°C and 28°C

The FCEG is aliquoted by duplicate in volumes of 5 mL in 8 mL test tubes and incubated uncovered at 4°C and 28°C for 24 hours, taking aliquots of 50 µL from each tube every 4 hours, and added to Petri dishes containing 1×10^6 yeast/mL of *C. albicans*, to determine the effect antifungal seeded in SDA at 28°C for 5 days.

2.2.6 Effect of the temperature on the antifungal activity of FCEG

The FCEG is aliquoted by duplicate in volumes of 5 mL in 8 mL test tubes and incubated at 60°C for 60 minutes, taking aliquots of 50 µL from each tube every 20 minutes, and added to Petri dishes containing 1×10^6 yeast/mL of *C. albicans*, to determine the effect antifungal seeded in SDA at 28°C for 5 days.

2.2.7 Effect of adsorption in activated carbon on the antifungal activity of FCEG

5 g of activated carbon (CAGR, ground and grain) were placed in 125 mL Erlenmeyer flasks, and 20 mL of FCEG were added, incubating them at 28°C for 6 days under constant agitation (100 rpm). Subsequently, aliquots of 50 µL were taken every 24 hours, and added to Petri dishes containing 1×10^6 yeast/mL of *C. albicans*, to determine the effect antifungal seeded in SDA at 28°C for 5 days.

2.3 Effect of FCEG on Lee medium filamentation for *Candida albicans*

Resuspend a pellet of a young colony suspected of *C. albicans* in 0.5 mL of Lee's medium and 50 µL of FCEG, were incubate at 37°C for 2½ to 3 hours. A fresh examination is performed to observe the development of the germ tube. This test is positive for *C. albicans* and *Candida dubliniensis* if a germ tube of approximately 5 to 15 µm long from the yeast, should not present constriction at the point of origin. The germ tube is like a hand mirror. After 3 hours of incubation, all *Candida* species can form tubes germ cells, except *Candida glabrata* [33].

2.4 Determination of the antifungal effect of FCEG on the growth of different fungi

Samples of the different species of fungi (obtained from Laboratory of Experimental Mycology/FCQ/UASLP) to be analyzed were taken and resuspended in 1 mL of SS. Subsequently, 50 µL aliquots of each suspension of the different fungi were taken and planted in petri dishes containing SDA, and 50 µL of the FCEG were added, spreading with a glass rod in the form of a triangle, and they were incubated at 28°C for 7 days, and the growth was compared with the controls of the different fungi seeded without FCEG.

2.5 Determination of the antimicrobial effect of FCEG on the growth of different bacteria

Samples of the different species of bacteria (obtained from Laboratory of Microbiology/FCQ/UASLP) to be analyzed were taken and resuspended in 10 mL of McFarland solution. Subsequently, 50 µL aliquots of each suspension of the different bacteria were taken and planted in petri dishes containing SDA, and 50 µL of the FCEG were added, spreading with a glass rod in the form of a triangle, and they were incubated at 35°C for 24-48 hours, and the growth based on the optimum density obtained from McFarland turbidity standard was compared with the controls of the different bacteria seeded without FCEG [34].

2.6 Determination of Protein

This was determined by the method of Lowry et al. (1951), using bovine serum albumin as standard [35].

3. RESULTS AND DISCUSSION

3.1 Analysis of some properties of FCEG

In this work, some properties of the antifungal activity of FCEG were analyzed, such as: half-life, temperature, effect of different dilutions, minimum inhibitory concentration of FCEG, etc., using the fungus like yeast *C. albicans* as standard, obtaining the following results: Regarding the effect of different dilutions of FCEG, it was found that any dilution with SS, inhibits its antifungal properties, which does not occur when analyzed without previously diluting it (fig. 1). On the other hand, the minimum inhibitory concentration of FCEG was between 40 and 50 µL (0.8 and 1 mg/mL of protein) (fig. 2), while the concentration of 50 µL used similarly inhibited the growth of different concentrations of *C. albicans* (1×10^6 to 10×10^6 yeast/mL), since no growth was observed at 1×10^6 , 2×10^6 , and 3×10^6 yeasts/mL, and only little growth was observed at the other concentrations (fig. 3). In relation to the half-life of the activity of the FCEG, it was found that it is completely lost between 55 and 60 days after obtaining it at 4°C in covered containers, although the loss of activity begins at 36 days (fig 4), while in uncovered containers, both at 4°C as at 28°C, activity is lost at 24 hours of incubation (fig 5), as well as at 60°C at 60 minutes, both in covered and uncovered containers (fig 6). Likewise, when trying to partially purify the FCEG with activated carbon (ground and in grain), it was observed that this treatment also eliminates the antifungal activity of the extract, since at 76 hours, yeast growth is observed in the cultures, being this 100% at 6 days of incubation (fig 7), and finally, the effect of FCEG on germ tube induction in *C. albicans* was analyzed, finding that it significantly inhibits germ tube formation (fig 8).

In relation to these parameters, there are few related reports, like: for evaluate the antimould activity of oregano, thyme, rosemary, and clove essential oils and some of their main constituents: eugenol, carvacrol and thymol against *A. niger*.

UNDER PEER REVIEW

and all the investigated agents showed no inhibitory effect on this fungus, which grow at concentrations lower than 10% (v/v) [4], for the lyophilized extract of garlic diluted to 10%, the minimum inhibitory concentration was 500 µg/mL for *T. mentagrophytes*, *T. rubrum* and *Microsporum canis* [19], the antifungal activities of fresh garlic (*A. sativum*) and ginger (*Z. officinale*) on the growth of three known pathogenic fungi were investigated. The test organisms were *Aspergillus* spp *Penicillium* spp and *C. albicans*. Under the conditions analyzed the more concentrated extracts (100 g/50 mL) inhibited the fungi growth more than the less concentrated (100 g/100 mL) extracts where moderate inhibition was observed. The actions of these extracts in these organisms were more on garlic. Between the two extracts used, the best in terms of inhibition is garlic followed by ginger [20]. The ajoene inhibits the development of *H. capsulatum* in mycelial phase at concentrations of 1.25 to 4.3 g/mL [21]. Too was evaluate the effect of the garlic crude extract on the in vitro development of *A. parasiticus* and *A. niger* to control the occurrence of these fungi on crops and other foods. The effect of the extract on the development of the fungi was evaluated by measuring of inhibition zones, puncture, colony forming units (CFU) and determination of minimum inhibitory concentration (MIC). The interaction of the garlic crude extract with the fungi was observed using visible light microscopy. The garlic crude extract induced inhibition zones of 12 mm in *A. parasiticus* and 15.5 mm for *A. niger*, it also inhibited growth in 13 and 46.8% respectively. The CMI for *A. parasiticus* was found to be the 1:2 dilution (50 µL of crude extract) and the 1:32 dilution (3.12 µL of crude extract) for *A. niger*. The treatment also inhibited the production of mycelium and sporulation of the two fungi species [23]. On the other hand, was analyzed the in vitro sensitivity of an isolate of *T. rubrum* and an isolation of *T. mentagrophytes* against ajoene, finding that this compound was able to inhibit the growth of both isolates showing a minimum inhibitory concentration (MIC) of 60 µg/mL and a concentration minimum fungicide (CMF) of 75 µg/mL [36]. The extract concentration also affects the antimicrobial activity, *Allium* extract concentrated (100%) have more lethal effect, followed by 75, 50, and 25% the least effective, obtained from McFarland turbidity methodology, and *Pseudomonas aeruginosa* showed resistance against extracts of both *A. sativum* and *Allium tuberosum*, with no zone of inhibition. The highest antimicrobial activity of *A. tuberosum* was noticed against *S. aureus* and *B. subtilis* with 43.9 and 40.7-mm zone of inhibition using 100% extract, respectively, followed by 75, 50, and 25%. The least antimicrobial activity was noticed for *Enterococcus faecalis* with zone of inhibition of 20.07 mm in 100% extract, whereas *P. aeruginosa* showed resistance against all concentrations evaluated [37]. Bernaldez and Vicencio (2021), determined the antibacterial activity of soap from garlic extract using the paper-disc method and Kirby-Bauer antibacterial sensitivity test against *S. aureus* and *E. coli*, and to determine the physical properties of garlic soap and the presence of saponin through phytochemical screening. Garlic soap showed antibacterial activity against *E. coli* and *S. aureus*. Mean zone of inhibition was numerically higher in plate extract obtained using garlic soap (14.70 mm-18 mm), compared to commercial soap [38], and was studied the antibacterial activity of the ethanolic extracts of *A. sativum* "Garlic" on *S. aureus* ATCC 25923 cultures. Ethyl alcohol of 96° was used for the extraction of the metabolites of the plants and concentrations at 100%, 75% and 50% were prepared, and the antibacterial activity was demonstrated by the method of disc diffusion in agar or Kirby Bauer, the ethanolic extracts of the *A. sativum* "Garlic" leaf presented inhibition halos of 20,433 mm, 17,126 and 10,659 for the concentrations of 100%, 75% and 50% respectively [39]. With respect to stability, we found that it is completely lost between 55 and 60 days after obtaining it at 4°C in covered containers, and evaluating the suitability of five different garlic cultivars for the processing of unsalted garlic paste, chopped fried garlic, and fried sliced garlic, the concentration of allicin in the products was evaluated immediately after processing and at 45-day intervals during 180 days of storage, and the amount of allicin lost during the process to obtain paste for the different varieties was less than 9.5%, and it reached a maximum loss of 22% for the commercial varieties during storage (180 days) [40].

This microorganism is a dimorphic fungus, which is in the form of yeast when it is in the saprophytic state, while in the parasitic state, it forms filaments (hyphae and pseudohyphae) of variable length. This capacity is related to the pathogenicity of this yeast fungus, in addition to various metabolic products, as well as some components of the cell wall are involved in the mechanisms of pathogenicity by this species [41]. The transition from yeast to hyphae is one of the virulence attributes that enable to *C. albicans* to invade tissues. It has been found that the growth of filamentous form has advantages over yeast in penetrating the cell or tissue, and although the hypha may be suitable to open the gap between tissue barriers, thanks because its tip is the site of secretion of enzymes capable of degrading proteins, lipids, and other cellular components, it facilitates their infiltration into solid and woven substrates [42]. There are three findings that support the hypothesis of that filamentation is required for virulence by this fungus: 1. Filament formation is stimulated at 37°C in the presence of serum, with neutral pH. 2. The newly formed filaments (called tubes germ cells) are more adherent to cells mammals than yeasts and adherence is the requirement for tissue penetration. 3. Yeasts captured by macrophages produce filaments and can lyse them, therefore, the formation of filaments is a form to evade host defense mechanisms [43, 44]. And finally, in this work, the FCEG significantly inhibits germ tube induction in *C. albicans*, which can be very important and effective for the treatment of diseases related to *C. albicans*, but more studies are required to determine the role of germination inhibition, in addition to the fact that in the literature we did not find related reports.




Fig. 1. Effect of different FCEG dilutions on the growth of *C. albicans*. 1×10^6 yeast/mL. 28°C. 5 days of incubation. *+ 2.5 cms, ++ 5.0 cms, +++ 7.5 cms, ++++ 10 cms.




Fig. 2. Effect of different FCEG concentrations (mg/mL) on the growth of *C. albicans*. 1×10^6 yeast/mL. 28°C. 5 days of incubation. *+ 2.5 cms, ++ 5.0 cms, +++ 7.5 cms, ++++ 10 cms.




Fig. 3. Effect of FCEG (1 mg/mL) on the growth of different *C. albicans* concentrations 28°C.

5 days of incubation. 1,10 controls without FCEG. *+ 2.5 cms, ++ 5.0 cms,

UNDER PEER REVIEW

+++ 7.5 cms, ++++ 10 cms.




Fig. 4. Half-life of antifungal activity of FCEG to 4°C in 60 days. *C. albicans* (1×10^6 yeast/mL). 28°C. 5 days of incubation. *+ 2.5 cms, ++ 5.0 cms, +++ 7.5 cms, ++++ 10 cms.




Fig. 5. Effect of tube capping on the antifungal activity of FCEG at 4°C and 28°C. *C. albicans* (1×10^6 yeast/mL) 28°C. 5 days of incubation. *+ 2.5 cms, ++ 5.0 cms, +++ 7.5 cms, ++++ 10 cms.




Fig. 6. Effect of tube capping on the antifungal activity of FCEG at 4°C and 28°C. *C. albicans* (1×10^6 yeast/mL) 28°C. 5 days of incubation. *+ 2.5 cms, ++ 5.0 cms, +++ 7.5 cms, ++++ 10 cms.




Fig. 7. Effect of adsorption in activated carbon (5 g) on the antifungal activity of FCEG (20 mL). *C. albicans* (1×10^6 yeast/mL) 28°C. 5 days of incubation. 100 rpm. *+ 2.5 cms, ++ 5.0 cms, +++ 7.5 cms, ++++ 10 cms.




Fig. 8. Effect of FCEG (50 μ L) on the filamentation in *C. albicans* in Lee medium.

3.2 Analysis of the antifungal and antibacterial effect of FCEG on the growth of different fungi.

A.- Control time 0

B.- Positive Control 2½ hours

C.- Problem with FCEG (50 µL)

On the other hand, when analyzing the antifungal effect of FCEG on the growth of different species of fungi and bacteria, it was found that it inhibits completely the growth of all the fungi and bacteria analyzed (fig 9 -16), which is like with different reports in the literature, in which the antimicrobial effect of garlic is analyzed under different conditions: such for *Fusarium oxysporum*, it was shown that *A. sativum* was effective at all the extract concentrations tested (25, 50, 75, and 100 g in 100 mL of sterile distilled water), with inhibitions ranging from 41.1 to 76.6% [5], 1% of *A. sativum* extract concentration of 0.0004 mg/mL, inhibit the growth of *T. rubrum* [16], for the ethanolic extract from *A. sativum* bulbs had an antifungal effect on crops of the same dermatophyte at different concentrations (%) [17], the essential oil of garlic for the control of *A. tenuissima*, in which, radial growth in vitro and biomass of this fungus in presence of the garlic essential oil (1000 ppm), the radial growth and biomass production were inhibited 100 and 86.20%, respectively, it compared with control PDA-Tween 80 without oil [18], for lyophilizate garlic, in *C. albicans*, it showed bigger inhibition diameter among 4000 to 5000 µg, a minimum inhibitory concentration of 2500 µg/mL and fungicidal action of 5000 µg [19], and for some dermatophytes [20], also, the inhibitory effect of ajoene on the mycelial phase of *H. capsulatum* [21]. Too, the garlic-chili hydrolyzates extracted from maceration and/or Soxhlet extraction with ethanol as solvent, show an effective inhibition on pathogenic fungi like the *Monilia* assuming a similar behavior on it, as well as the hydrolysates of oregano that manage to reach an inhibition percentage of 75.2% at a concentration minimum of 40% for approximately 12 days [22], and the effect against *A. parasiticus* and *A. niger* [23], and Saprolegniasis in rainbow trout (*Oncorhynchus mykiss*) [24] was evaluated, observing that the inhibition is very efficient, and their antimicrobial effect against *E. coli*, *S. aureus*, and *A. niger* [25,

26,27, and 28], *S. mutans* [29], molds, yeasts, coliforms, and *E. coli* [30] on the Gram positive bacteria *B. subtilis* and *S. epidermidis* and other Gram negative bacteria *E. coli* and *S. dysinteriae* [31].



Fig. 9. Effect of FCEG on the growth of different fungal strains. 1×10^6 cells/mL. 28°C. 7-15 days of incubation. *+ 2.5 cms, ++ 5.0 cms, +++ 7.5 cms, ++++ 10 cms.




Fig. 10. Effect of FCEG on the growth of different bacteria and yeast strains, by optimum density obtained from McFarland turbidity standard. 1×10^6 cells/mL. 35°C. 1-2 days of incubation. (For *C. albicans* 28°C and 5 days of incubation).




Fig. 11. Effect of different dilutions of FCEG in sterile saline solution on the growth of *C. albicans*. (1×10^6 yeast/mL. 28°C . 5 days of incubation).

- 1.- Control with FCEG
- 2.- *C. albicans* + FCEG 1.10 (v/v)
- 3.- *C. albicans* + FCEG 1.100 (v/v)
- 4.- *C. albicans* + FCEG 1.1000 (v/v)




Fig. 12. Effect of FCEG in sterile saline solution on the growth of different concentrations of *C. albicans*. (50 μL . 28°C . 5 days of incubation).

- 1.- *C. albicans* 1×10^6 yeast/mL without FCEG
- 2.- *C. albicans* 10×10^6 yeast/mL without FCEG
- 3.- *C. albicans* 5×10^6 yeast/mL with FCEG
- 4.- *C. albicans* 6×10^6 yeast/mL with FCEG
- 5.- *C. albicans* 7×10^6 yeast/mL with FCEG
- 6.- *C. albicans* 8×10^6 yeast/mL with FCEG




Fig. 13. Effect of different concentrations of FCEG on the growth of *C. albicans*. (1 x10⁶ yeast/mL. 28°C. 5 days of incubation).

- 1.- *C. albicans* 1 x10⁶ yeast/mL without FCEG
- 2.- *C. albicans* 10 x10⁶ yeast/mL with 50 µL of FCEG
- 3.- *C. albicans* 1 x10⁶ yeast/mL with 10 µL of FCEG
- 4.- *C. albicans* 1 x10⁶ yeast/mL with 20 µL of FCEG
- 5.- *C. albicans* 1 x10⁶ yeast/mL with 30 µL of FCEG
- 6.- *C. albicans* 1 x10⁶ yeast/mL with 40 µL of FCEG




Fig. 14. Effect of FCEG on the growth of *H. capsulatum*. (28°C. 15 days of incubation).

- 1.- *H. capsulatum* without FCEG
- 2.- *H. capsulatum* with 50 µL of FCEG
- 3.- *H. capsulatum* with 50 µL of FCEG




Fig. 15. Effect of FCEG on the growth of *Wangiella dermatitidis* (28°C. 5 days of incubation).

- 1.- *Wangiella dermatitidis* without FCEG**
- 2.- *Wangiella dermatitidis* with 50 μ L of FCEG**




Fig. 16. Effect of FCEG on the growth of *Cladosporium carrionii* (28°C. 5 days of incubation).

- 1.- *Cladosporium carrionii* without FCEG**
- 2.- *Cladosporium carrionii* with 50 μ L of FCEG**

4. CONCLUSION

Actually, the pharmaceutical industry has largely focused on high-throughput biochemical screening programmes for the discovery and development of new drugs, the use of natural products for medicinal and antimicrobial purposes is an ancient practice [15, 45], and the use of garlic, to which it is common to attribute a wide variety of properties, among which are: antifungal and antileishmanial activity [4, 7], but, until 1944 when Cavallito and Bailey [46] isolated and described the properties of allicin, the compound responsible for garlic's characteristic pungent odor, that researchers gained a clearer insight into the chemical wonder carefully packaged by nature in the composite bulbs of this edible *Allium*, and thus began decades of extensive research on allicin, "the heart of garlic" [47], and the ajoene [(E, Z)

4,5,9-tritriadodeca-1,6,11-triene-9-oxide] is an organosulfur compound derived from garlic, from which systematically

UNDER PEER REVIEW

demonstrated its potential inhibitory activity in vitro and in vivo on most of the fungi that cause human mycoses this compound acts selectively on the plasma membrane by inhibiting the synthesis of phosphatidylcholine, with accumulation of phosphatidylethanolamine and, consequently, causing cell death [15, 48, and 49], and the effects of garlic oil were studied with *Penicillium funiculosum* as a model strain. Results showed that the minimum fungicidal concentrations (MFCs, v/v) were 0.125 and 0.0313 % in agar medium and broth medium, respectively, suggesting that the garlic oil had a strong antifungal activity. The main ingredients of garlic oil were identified as sulfides, mainly including disulfides (36 %), trisulfides (32 %) and monosulfides (29 %) by gas chromatograph-mass spectrometer (GC/MS), which were estimated as the dominant antifungal factors [50]. Too, was studied their powder has a good adsorption capacity for mercury [51], and in this work we analyze some characteristics of FCEG, and their effect on the growth of different bacteria and fungi, with the next conclusions:

- 1.- The dilution of the FCEG with sterile saline solution 0.85% (w/v) inhibits its antifungal properties.
- 2.- The different concentrations of FCEG analyzed have the same inhibitory.
- 3.- As the concentration of *C. albicans* yeasts increases, the effectiveness of the FCEG partially decreases.
- 4.- FCEG has a half-life between 45-60 days at 4°C, while in uncovered containers it loses its activity at 24 hours, both at 4°C and 28°C.
- 5.- FCEG loses its antifungal activity at 60 minutes to 60°C.
- 6.- Activated charcoal treatment also inhibits antifungal activity of FCEG.
- 7.- FCEG significantly inhibits germ tube induction in *C. albicans*.
- 8.- The different species of fungi and bacteria analyzed were inhibited in their growth by the FCEG.

COMPETING INTERESTS

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

- [1] Echániz-Aviles G. Resistir la resistencia: investigación, alianzas y soluciones. Ciencia Cofepris. 2021;3: 9-13. <https://www.gob.mx>. Español.
- [2] González Mendoza J, Maguiña Vargas C, González Ponce FM. La resistencia a los antibióticos: un problema muy serio. Acta Med. Per. 2019;36(2):145-51. DOI: <https://doi.org/10.35663/amp.2019.362.816/>. Español.
- [3] Organización Mundial de la Salud. Resistencia a los antimicrobianos. OMS. 2020:1-10. <https://www.who.int/es/news-room/fact-sheets/detail/antimicrobial-resistance>. Español.
- [4] Bouddine L, Louaste B, Achahbar S, Chami N, Chami F, Remmal A. Comparative study of the antifungal activity of some essential oils and their major phenolic components against *Aspergillus niger* using three different methods. Afric. J. Biotechnol. 2012;11(76):14083-14087. DOI: 10.5897/AJB11.3293
- [5] Okigbo RN, Okorie RE, Putheti RR. *In vitro* effects of garlic (*Allium sativum*) and african basil (*Ocinum Gratissimum* L.) on pathogens isolated from rotted Cassava roots. Interciencia. 2002;34(19):742-747. <http://www.redalyc.org/articulo.oa?id=33913147011>.
- [6] Shentu X, Zhan X, Ma Z, Yu X, Zhang Ch. Antifungal activity of metabolites of the endophytic fungus *Trichoderma brevicompactum* from garlic. Braz. J. Microbiol. 2012;45(1): 248-254. doi: 10.1590/S1517-83822014005000036.
- [7] Gamboa-Leon R, Vera-Ku M, Peraza-Sanchez SR, Ku-Chulim C, Horta-Baas A, Rosado-Vallado M. Antileishmanial activity of a mixture of *Tridax procumbens* and *Allium sativum* in mice. Parasite. 2014;21(15):1-7. DOI: 10.1051/parasite/2014016
- [8] Rodríguez Quintero JA, Méndez Márquez RO, Gutiérrez Hernández R, Reyes Estrada CA. Evaluación del efecto antibacteriano del extracto de cilantro (*Coriandrum sativum*) sobre bacterias patógenas gastrointestinales. Memoria Universitaria. II Seminario de Investigación en Economía: Sustentabilidad y Nanotecnología. 2021;3(2):1-10. <<https://revistas.uaz.edu.mx/index.php/MemUni/article/view/1290>>. Español.
- [9] Ramírez Aristizabal LS, Marín D. Evaluación de la actividad antibacteriana de aceites esenciales y extractos etanólicos utilizando métodos de difusión en agar y dilución en pozo. Scientia et Technica. 2012;XVII(50):152-157. <http://www.redalyc.org/articulo.oa?id=84923878022>. Español.

[10] Souza de Oliveira LF, Meneghello Fuentefria A, da Silva Klein F, Mansur Machado M. Antifungal activity against *Cryptococcus neoformans* strains and genotoxicity assessment in human leukocyte cells of *Euphorbia tirucalli* L. Braz. J. Biotechnol. 2014;45(4):1349-1355. doi: 10.1590/s1517-83822014000400027

UNDER PEER REVIEW

- [11] Valero Galván J, González Díaz CA, González Fernández R. Efecto de los extractos acuosos de hojas de plantas de gobernadora (*Larrea tridentata*), hojaseñ (*Flourensia cernua*) y encino (*Quercus pungens*), sobre el crecimiento micelial in vitro de hongos fitopatógenos. *Acta Universitaria*. 2014;24(5):13-19. doi: 10.15174. au.2014.630. Español.
- [12] Itepu VE, 1, Umezunwa OJ, Ilukho FA, Collins DA, Samuel C. The Effect of Structural Modification on the Blood Brain Barrier Permeation Properties of Allicin. *Asian J. Res. Biochem.* 2019;5(3):1-7. DOI: 10.9734/AJRB/2019/v5i330092.
- [13] Vaijayanthimala P, Amutha K, Anu M, Archana S, Bhuvaneswari S, Balamurugan TS, et. al. In vitro anticoagulant activity of *Allium sativum* plant extract. *World J. Pharm. Pharmac. Scien.* 2017;6(9):1256-1261. DOI: 10.20959/wjpps20179-10017
- [14] Bhattacharya S, Majia U, Khanc GA, Dasd R, Asru K, Sinha AK, et. al. Antidiabetic role of a novel protein from garlic via NO in expression of Glut-4/insulin in liver of alloxan induced diabetic mice. *Blomed. Pharmacot.* 2019;111:1302-1314. <https://doi.org/10.1016/j.biopha.2019.01.036>
- [15] Borlinghaus J, Albrecht F, Gruhlke MCH, Nwachukwu ID, Slusarenko AJ. Allicin: Chemistry and Biological Properties. *Molecules*. 2014;19:12591-12618; doi:10.3390/molecules190812591.
- [16] Robles-Martínez M, Cárdenas González JF, Pérez-Vázquez FJ, Montejano-Carrizales JM, Pérez E, Patiño-Herrera R. Antimycotic Activity Potentiation of *Allium sativum* Extract and Silver Nanoparticles against *Trichophyton rubrum*. *Chem. Biodiver.* 2019;16:1-14. e1800525. DOI: 10.1002/cbdv.2018.
- [17] Medina López C. Efecto antimicótico in vitro del extracto etanólico del bulbo de *Allium sativum* (ajo) en cultivos de *Trichophyton rubrum*. Escuela Profesional de Farmacia y Bioquímica. Facultad de Ciencias de la Salud. Universidad Católica Los Angeles Chimbote. Tesis para optar el Título Profesional de Químico Farmacéutico. 2020. Trujillo, Perú. <http://repositorio.uladech.edu.pe/handle/123456789/22952>. Español.
- [18] Muy-Rangel MD, Osuna-Valle JR, García-Estrada RS, San Martín-Hernández C, Quintana-Obregón EA. In vitro antifungal activity of garlic essential oil (*Allium sativum* L.) against *Alternaria tenuissima*. *Rev. Mex. Fitopatol.* 2018;36(1):162-171. DOI: 10.18781/R.MEX.FIT.1708-3. Bilingual.
- [19] Lora Cahuas C, Luján Velázquez M, Robles Castillo H, Saravia Cuevas V, Cabeza Rodríguez J. Efecto *in vitro* de diferentes concentraciones de *Allium sativum* "ajo" frente a dermatofitos y *Candida albicans*. *UCV-Scientia*. 2010;2(2):23-33. DOI: <https://doi.org/10.18050/revucv-scientia.v2i2.877>. Español.
- [20] Agi VN, Azike CA. Antifungal Action of Garlic (*Allium sativum*) and Ginger (*Zingiber officinale*) on Some Pathogenic Fungi. *Asian J. Res. Biochem.* 2019;4(4):1-6. DOI: 10.9734/AJRB/2019/v4i430075.
- [21] Torres J, Romero H. Actividad antifúngica in vitro del ajoeno en cinco aislamientos clínicos de *Histoplasma capsulatum* var. *Capsulatum*. *Rev. Iberoam. Micol.* 2012;29(1):24–28. doi:10.1016/j.riam.2011.04.001. Español.
- [22] León Ospina IS, Sánchez Giraldo LM. Análisis del efecto antifúngico de los hidrolatos ajo-ají sobre la monilia *Moniliophthora roreri*. Facultad de Ingenierías. Programa de Ingeniería Química. Fundación Universidad de América. Proyecto de Grado para optar al título de Ingeniero Químico. 2021. Bogotá, D.C. Colombia. <https://repositorio.uamerica.edu.co>. Español.
- [23] Juárez-Segovia, KG, Díaz-Darcía EJ, Méndez-López MD, Pina-Canseco, MS, Pérez-Santiago AD, Sánchez-Medina MA. Efecto de extractos crudos de ajo (*Allium sativum*) sobre el desarrollo *in vitro* de *Aspergillus parasiticus* y *Aspergillus niger*. *Polibotánica*. 2019;47:99-111. DOI: 10.18387/polibotanica.47.8. Español.
- [24] Paredes Sandoval JC. Efecto del ajo (*Allium sativum*) en el tratamiento de Saprolegniasis en trucha arcoíris (*Oncorhynchus mykiss*). Carrera de Medicina Veterinaria y Zootecnia. Facultad de Ciencias Agropecuarias. Universidad Técnica de Ambato. Documento Final del Proyecto de Investigación como requisito para obtener el grado de Médico Veterinario Zootecnista. 2021. Cevallos. Ecuador. <https://repositorio.uta.edu.ec>. Español.
- [25] Pereda Aguilar LE, Efecto antibacteriano in vitro de extracto hidroalcohólico de *Allium sativum* (ajo) y *Plantago major* (llantén) frente a cultivos de *Escherichia coli*. Escuela Profesional de Farmacia y Bioquímica. Facultad de Ciencias de la Salud. Universidad Católica Los Ángeles. Chimbote. Tesis para optar el Título Profesional de Químico Farmacéutico. 2020. <http://repositorio.uladech.edu.pe/handle/123456789/23789>. Español.
- [26] Coronel Delgado JD, Grandez Ventura E. Efecto antibacteriano in-vitro de los extractos hidroalcohólicos de *Allium sativum* "Ajo" y *Ocimum basilicum* "Albahaca" sobre *Staphylococcus aureus*. Escuela Profesional de Ciencias Farmacéuticas y Bioquímica. Facultad de Ciencias de la Salud. Universidad Roosevelt. Para optar por el Título Profesional de Químico Farmacéutico. 2021. Huancayo. Perú. <https://repositorio.uoosevelt.edu.pe>. Español.
- [27] Villa Sánchez FE, Moncayo Molina WE, Alvarado Aguilar MCh, Leal Chantong AA, Daza Barcia DD. Síntesis verde de nanopartículas de plata (AgNPs) utilizando ajo (*Allium sativum* L.) estudiando su actividad antimicrobial y catalíticas. *FacSalud*. 2021;5(8):39-50. <https://doi.org/1029076/issn2602-8360vol5iss82021pp39-50p>. Español.
- [28] Requejo Ramos MC, Callao Rodríguez JL. Actividad antibacteriana in vitro del extracto etanólico de *Datura stramonium* (chamico) y *Allium sativum* (ajo) sobre *Staphylococcus aureus*. Escuela Profesional de Farmacia y Bioquímica. Facultad de Ciencias de la Salud. Universidad María Auxiliadora. Tesis para optar el Título Profesional de Químico Farmacéutico. Lima, Perú. 2021. <http://repositorio.uma.edu.pe/handle/UMA/404>. Español.

- [29] Jimenez Macias AF, Zambrano Gutiérrez MI. Efecto antibacteriano del extracto de *Allium sativum* (ajo) blanco, púrpura y Clorhexidina al 0,12% sobre cepas de *Streptococcus mutans*. Dom. Cien. 2017;3(1):234-247. 10.23857/dom.cien.pocaip.2017.3.1.234-247. Español.
- [30] Delzo García SR. Determinación de las características microbiológicas de pasta de *Capsicum chinense* (ají panca) y *Allium sativum* (ajos) expendidos en el mercado de la ciudad de Satipo. Escuela Profesional de Ingeniería en Industrias Alimentarias Tropical. Facultad de Ciencias Agrarias. Universidad Nacional del Centro del Perú. Para optar el Título Profesional de Ingeniera en Ciencias Agrarias. Especialidad en Ingeniería en Industrias Alimentarias. 2021. Satipo. Perú. <http://hdl.handle.net/20.500.12894/7288>. Español.
- [31] Jacha Ayala ZP, Celestino Laberiano MD. Función antibacteriana del extracto puro de bulbo de *Allium sativum* sobre Gram positivos y Gram negativos. Inv. Valdizana. 2017;11(4):255-261. <http://revistas.unheval.edu.pe/index.php/riv/article/view/128>. Español.
- [32] Madrigal Redondo GL, Loria Gutiérrez A, Blanco Barrantes J, Porras Navarro M, Ortega Monge Cerda MC, Cerdas Vargas MJ. Aspectos generales del *Allium sativum*-una revisión. Ars Pharmaceutica. 2021;62(4):471-481. DOI: <https://doi.org/10.30827/ars.v62i4.20843>. Español.
- [33] Lee KL, Buckley HR, Campbell CC. An amino acid liquid synthetic medium for the development of mycelial and yeast forms of *Candida albicans*. Sabouraudia. 1975;13(2):148-153. doi: 10.1080/0036217758519027
- [34] McFarland J. (1907). The Nephelometer: An instrument for estimating the number of bacteria in suspensions used for calculating the opsonic index and for vaccines. JAMA. 1907;XLIX(14):1176-1178. doi:10.1001/jama.1907.25320140022001f
- [35] Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. J. Biol. Chem. 1951;193(19):265-275. PMID: 14907.
- [36] de González MI, Mendoza M, de Albornoz MB, Apitz-Castro R. Efectos del ajoeno sobre dermatofitos, *Candida albicans* y *Malassezia furfur*. Rev. Iberoam. Micol.1998;15:277-281. Hppt://www.reviberoammicol.com>1998-15. Español.
- [37] Nauman K, Iftikhar A, Malik Shah ZL, Tariq R, Sardar Atiq F. Comparison of Antimicrobial Activity, Phytochemical Profile and Minerals Composition of Garlic *Allium sativum* and *Allium tuberosum*. J. Korean Soc. Appl. Biol. Chem.2014;57(3): 311-317. DOI 10.1007/s13765-014-4021-4.
- [38] Bernaldez JLI, Vicencio MCG. Antibacterial Activity of Soap Formulated from Garlic (*Allium sativum* L.). J. Adv. Microbiol. 2021;21(1):63-67. DOI: 10.9734/JAMB/2021/v21i130320.
- [39] Rodríguez Vega JL, Mejía Pinedo DA, Lora Loza MG, Pérez Martinto P. Actividad extracto etanólico de hoja de *Allium sativum* (ajo) sobre *Staphylococcus aureus*. Rev. Epistemia. Número Especial. 2020;4(2)1-7. <https://hdl.handle.net/20.500.12692/56324>. Español.
- [40] Prat P, Henrique CM, de Souza AS, Nunes da Silva VS, Bertoldo Pacheco MT. Evaluation of allicin stability in processed garlic of different cultivars. Food Sci. Technol, Campinas. 2014;34(3):623-628. DOI: <http://dx.doi.org/10.1590/1678-4X.6397>
- [41] Pardi G. Determinantes de Patogenicidad de *Candida Albicans*: Acta Odontol. Venez. 2002;40(2):185-192. Disponible en: http://ve.scielo.org/scielo.php?script=sci_arttext&pid=S0001-63652002000200016&lng=es. Español.
- [42] Hube B, Naglik J. *Candida albicans* proteinases: resolving the mystery of a gene family. Microbiol. 2001;147:1997-2005. doi: 10.1099/00221287-147-8-1997.
- [43] Mitchell AP. Dimorphism and virulence in *Candida albicans*. Curr. Opin. Microbiol. 1998;1:687-92. DOI: [10.1016/s1369-5274\(98\)80116-1](https://doi.org/10.1016/s1369-5274(98)80116-1).
- [44] Castrillón Rivera LE, Palma Ramos A, Padilla Desgarenes C. Factores de virulencia en *Candida* sp. Dermatología Rev. Mex. 2005;49:12-27.

<https://www.researchgate.net/publication/286816593>. Español.

- [45] Sacheli R, Hayette MP. Antifungal Resistance in Dermatophytes: Genetic Considerations, Clinical Presentations and Alternative Therapies. *J. Fungi*. 2021;7(983):1-23. [https://doi.org/ 10.3390/jof7110983](https://doi.org/10.3390/jof7110983)
- [46] Cavallito C, Bailey J. Allicin, the antibacterial principle of *Allium sativum*. I. Isolation, physical properties and antibacterial action. *J. Am. Chem. Soc.* 1944;66:1950–1951. <https://doi.org/10.1021/ja01239a048>
- [47] Josling, P. Allicin—The Heart of Garlic; NWI Publishing: Callahan Florida, FL, USA, 2007. www.allimax.us.
- [48] San-Blas G, Urbina JA, Marchán E, Contreras LM, Sorais F, San-Blas F. Inhibition of *Paracoccidioides brasiliensis* by ajoene is associated with blockade of phosphatidylcholine biosynthesis. *Microbiology*. 1997;143:1583–6. doi: 10.1099/00221287-143-5-1583.
- [49] Urbina JA, Marchan E, Lazard K, Visbal G, Apitz-Castro R, Gil F, et al. Inhibition of phosphatidylcholine biosynthesis and cell proliferation in *Trypanosoma cruzi* by ajoene, an antiplatelet compound isolated from garlic. *Biochem Pharmacol*. 1993;22:2381–7. doi: 10.1016/0006-2952(93)90217-k.
- [50] Li WR, Shi QS, Liang Q, Huang XM, Chen YB. Antifungal effect and mechanism of garlic oil on *Penicillium funiculosum*. *Appl. Microbiol. Biotechnol.* 2014;98:8337–8346 DOI 10.1007/s00253-014-5919-9.
- [51] Eom Y, Won JH, Ryu JY, Lee TG. Biosorption of mercury(II) ions from aqueous solution by garlic (*Allium sativum* L.) powder. *Korean J. Chem. Eng.* 2011;28(6):1439-1443. DOI: 10.1007/s11814-010-0514-y
-