

Minireview Article

Diet and its Relationship with Peripheral Vascular Disease

Abstract

Peripheral artery disease is a major health problem. It affects over 200 million people globally. It results in considerable suffering, curtailed functionality, and markedly reduced quality of life. Its presence prognosticates significantly higher major cardiovascular events. Atherosclerosis is the main disease process underlying its development and progression. Several lifestyle modifications have been shown to stem atherosclerosis. Consumption of a prudent diet is one such beneficial lifestyle. A prudent diet promotes the consumption of fruit, vegetables, whole grains, legumes, healthy fats such as nuts and seeds, low-fat dairy products, fish, and small amounts of lean meat. It limits the consumption of processed red meats, saturated fats, refined carbohydrates, and sugar-sweetened drinks. This manuscript reviews the present understanding of the role of diet in the development and progression of peripheral artery disease.

Keywords: prudent diet, fruits and vegetables, whole grains, processed red meat, saturated fat, peripheral vascular disease

Introduction

Peripheral arterial disease (PAD) affecting the lower extremities is a common clinical condition¹. It is estimated to affect more than 200 million people globally². While it affects only 4.3% in subjects ≥ 40 years, its prevalence rises to 14.5% in those over 70 years³. Since humans are living longer and the aging population is growing, the incidence and prevalence of PAD is on the rise⁴. It is diagnosed by the presence of an ankle-brachial index (ABI) of <0.9 ⁵. This noninvasive method is extremely reliable and has been validated against the gold standard - contrast angiography⁶. Fowkes reported that ABI had a sensitivity of 95% and a specificity of 100%⁶. Common risk factors for PAD, besides aging, include smoking, hypertension (HTN), diabetes mellitus (T2DM), dyslipidemia, and chronic renal failure⁷. It is estimated that the 1.5-fold increase in the incidence of PAD with one risk factor climbs to a 10-fold increase if there are three or more risk factors involved⁸. Smoking is a major risk factor and data indicates that more than 80% of patients with PAD are current or former smokers⁹. The underlying pathology is predominantly atherosclerosis¹⁰. Atherosclerosis is a progressive pathology and usually concurrently affects other parts of the vascular system, such as the coronary arteries and the carotid arteries¹¹. Its pathophysiology includes low-grade inflammation, lipid accumulation, and plaque formation, leading to narrowing of the arteries¹². In the legs this results in ischemia, causing pain on walking and a significant functional loss¹³. The disease may progress to non-healing leg wounds, gangrene, and ultimately result in limb amputation¹⁴. Besides the gradual progression seen in most patients, about 25% develop sudden onset of pain at rest due to acute limb ischemia, often requiring urgent surgical intervention¹⁵. The health-related quality of life is markedly reduced in these patients¹⁶. Treatment is usually pharmacological or surgical¹⁷. Concurrent supervised physical exercise helps reduce morbidity and mortality and leads to a

significant improvement in the health-related quality of life¹⁸. Overall mortality rates are high, and in patients with symptomatic disease, they approach 8.4% over 2 years¹⁹. Kochar et al. determined that in PAD patients, cardiovascular causes were responsible for the majority of deaths²⁰. The economic burden (direct and indirect costs) related to PAD are extremely high²¹. Lifestyle factors, including diet, can play an important role in both primary and secondary prevention of this vascular disease²².

Discussion

Peripheral artery disease is an atherosclerotic vascular disease, especially affecting the lower legs¹. Since coronary artery disease (CAD) is also an atherosclerotic process, CAD and PAD share several common risk factors such as smoking, dyslipidemia, hypertension (HTN), and diabetes mellitus (T2DM)²³. An unhealthy diet is also a major risk factor²⁴. In a prospective cohort study, with a median follow-up of 21.7 years, Kulezic et al. reported an inverse association between adherence to a healthy diet and incident PAD²⁵. A diet rich in processed red meat, saturated fats, refined carbohydrates, and low in micronutrients like magnesium, phosphorus, and calcium as well as vitamins A, D, or E and the vitamin B group, is pro-atherogenic^{26,27}. A plant-based diet rich in fruits, vegetables, whole grains, and nuts helps reduce oxidative stress and endothelial dysfunction, and improve erythrocyte deformability, blood viscosity, oxygen perfusion, and is anti-atherosclerosis²⁸. A prudent diet can therefore play a major role in both the primary (development) and secondary (progression) prevention of PAD²⁹.

Obesity

Diet plays an important role in obesity³⁰. Obesity is usually determined by calculating the body mass index (BMI). A BMI ideally should be 18.5 kg/m² to 24.9 kg/m². A BMI of 25 kg/m² to 29.9 kg/m² is considered overweight. A BMI at or >30 kg/m² is designated as obese³¹. Compared to the BMI calculated obesity, abdominal obesity is more harmful to the cardiovascular system³². Abdominal obesity is objectively ascertained by several anthropometric measurements, such as the waist circumference (WC), waist-hip ratio and the waist height ratio³³. WC should ideally be <102 cm in males and <88 cm in females when measured to the nearest 0.1 cm at the umbilical level in a standing position³⁴. The waist-hip ratio is normally 0.85 or less for women and 0.9 or less for men³⁵. The waist-height ratio is infrequently used – this is calculated by dividing the WC by the patients height. A ratio > 0.5 is consistent with central obesity³⁶. Central or visceral obesity may exist even if the BMI is normal³⁷. Several studies have linked abdominal obesity with PAD³⁸. Wisse found that individuals with central or abdominal obesity had a 4-fold risk of developing PAD (Odds Ratio=4.532)³⁹. Visceral fat is metabolically active and is associated with the development of insulin resistance, inflammation, and oxidative stress⁴⁰. The result is an increase in atherosclerosis and PAD⁴¹. Abdominal obesity is also a known risk factor for CHD and this association has clinical significance as coronary heart disease (CHD) is responsible for many deaths in PAD patients²⁰. Brouwer et al. reported that the presence of abdominal obesity was associated with a 61% increase in the risk of concomitant CHD in these patients⁴². Diet plays an important role in the prevention of excess body weight³⁰. Although caloric intake restriction helps in weight reduction, the addition of exercise not only results in a greater weight loss but also helps to maintain the weight loss⁴³. Weight loss is associated with an improvement

in several atherosclerosis risk biomarkers such as total cholesterol, low-density lipoprotein cholesterol, triglycerides, systolic and diastolic blood pressure, glucose, insulin, and C-reactive protein⁴⁴⁻⁴⁸. Consequently, weight loss in obese patients results in an improvement in peripheral artery disease⁴⁹.

Fruits and Vegetables

It is well known that high consumption of fruits and vegetables is associated with a lower risk of cardiovascular diseases, especially coronary artery disease and stroke⁵⁰. Atherosclerosis is the common underlying pathology in these diseases – and PAD²³. Hung in 2003 found a modest beneficial effect on PAD, with fruit and vegetable consumption, in a study of male health professionals during a 12-year follow-up⁵¹. A Mediterranean dietary pattern (rich in plant-based foods and polyphenol-rich olive oil) is associated with a 56% reduced risk of PAD⁵². In a recent cross-sectional study involving over 3.6 million individuals, Heffron et al. found that the odds of PAD were reduced by 18% when individuals consuming 3 or more servings of fruits and vegetables per day were compared to those consuming >3 servings of fruits and vegetables less than once a month⁵³. In a recent study, Chen et al., in a study of 138,506 US postmenopausal women (1993–1998) in the Women's Health Initiative, with no known PAD at baseline, reported that higher intakes of vegetable protein were associated with a lower risk of developing PAD⁵⁴. Phytochemicals in plants are postulated to exert vascular benefits in these patients⁵⁵.

Whole grains/Fiber/legumes/Nuts

Whole-grain consumption was not found to be associated with reduced risk of PAD in a prospective cohort study of 14,082 individuals in the US with a mean follow-up of 19.9 years⁵⁶. However, whole grains are an important source of dietary fiber, and most studies on fiber intake have shown that fiber consumption beneficially influences PAD⁵⁷. In a cross-sectional study (1,267 subjects aged > or =30 years), lower consumption of fiber from whole grains was found in individuals with PAD⁵⁷. Dietary fiber exerts a protective effect on PAD via several mechanisms. These include reducing obesity, lowering cholesterol, increasing insulin sensitivity, reducing serum glucose levels, producing short-chain fatty acids by gut bacteria, and reducing inflammation^{58,59}. It has also been noted that individuals with a higher intake of fiber often lead healthier lifestyles, including participating in physical activity and avoiding tobacco smoking. Legume and nut intake is also helpful. In the recent study by Chen et al., higher intakes of legumes and dietary fiber were associated with a lower risk of PAD⁶⁰. In a study of adults, Heffron et al. found that increased nut intake was inversely associated with PAD prevalence⁶¹. Inflammation is strongly associated with atherosclerosis and nuts decrease inflammation. In the Multi-Ethnic Study of Atherosclerosis (MESA) (nearly 6000 participants), consumption of nuts and seeds was inversely associated with levels of inflammatory markers such as C-reactive protein, interleukin-6, and fibrinogen⁶².

Coffee/tea/Chocolate

Light-to-moderate coffee intake (0.5-3 cups/day) has been shown to be cardioprotective^{63,64}. Coffee consumption is associated with a decreased risk of developing PAD⁶⁵. In the Atherosclerosis Risk in Communities (ARIC) Study, the authors reported an inverse association

between coffee intake and incident PAD⁵⁶. In a retrospective analysis of 237 women (45-54 years of age), 16.8% of women that drank more than five cups/day of coffee had an ABI lower than 0.9 and as compared to its presence in 27.4% of women who were non-drinkers⁶⁵. Green tea consumption also provides beneficial vascular effects – in a study of 12 healthy men (29 ± 6 yrs), drinking catechin-rich green tea prevented abnormalities of vascular function induced by unhealthy lifestyles⁶⁶. Both coffee and tea (green or black) are high in the beneficial polyphenol antioxidants^{67,68}. Cocoa is also PAD friendly. Dark chocolate improves maximal walking distance and maximal walking time, 2 hours after ingestion in PAD patients⁶⁹. No benefits were noted with milk chocolate. Dark chocolate is richer in polyphenol compared to milk chocolate⁷⁰. As noted with tea and coffee, the polyphenols help in improving endothelial function by reducing oxidative stress and increasing nitric oxide generation^{67,68,71}. As a result, atherosclerosis is retarded.

Alcohol

As noted in coronary artery disease, the risk of PAD is beneficially affected by low to moderate alcohol consumption. In the Cardiovascular Health Study, of older adults (5635 participants) alcohol consumption of 1-13 drinks per week was associated with a lower risk of PAD⁷². In the ARIC (Atherosclerosis Risk in Communities) Study of 14,082 participants, 1–6 drinks/wk, but not ≥7 drinks, was associated with a lower risk of incident PAD⁶⁵. The benefits are postulated to be partly due to an improved lipid profile (especially a higher HDL-cholesterol) and better insulin sensitivity^{73,74}.

Red Meat

Red meat intake, especially that of processed red meat, has been repeatedly shown to increase the risk of cardiovascular diseases (CVD)⁷⁵⁻⁷⁷. A detrimental association with PAD has also been noted^{25,65}. In a recent study, Chen et al. reported that intake of unprocessed red meat, processed meat, and regular soft drinks was associated with a higher risk of PAD⁶⁰. High levels of sodium and nitrate preservatives in processed meats raise blood pressure and increase the incidence of T2DM, both contributing to the increased incidence of PAD⁷⁸⁻⁸⁰.

Saturated and Unsaturated Fat

The deleterious role of saturated fats on atherosclerotic vascular disease is well known⁸¹. In 2008, in a cross-sectional study of 1,267 subjects (aged > or =30 years) individuals with PAD had a higher intake of total (72.8 vs 69.1 g, p=0.016) and saturated fatty acids (17.4 vs 16.3 g, p=0.012)⁸². This harmful relationship was confirmed by Naqvi et al⁸³. In their study of 6352 adults aged 40 years and older participating in the U.S. National Health and Nutrition Examination Survey between 1999 and 2004, the intake of saturated fatty acids was associated with a higher prevalence of PAD (adjusted odds ratio 1.30). Mazidi et al. reported higher plasma trans-fatty acid levels in PAD patients, compared to those without the disease⁸⁴. On the other hand, a higher intake of unsaturated fats, especially eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) appear to be beneficial^{85,86}. Dark fish is a major source of EPA and DHA⁸⁷. However, in a recent cohort of 14,082 participants followed for a mean of 19.9 years, the authors found no association between consumption of fish and the risk of incident PAD⁵⁶.

Another recent study involving 55,248 participants (a median of 13.6 years of follow-up), Cox regression analyses with adjustments for established risk factors showed no statistically significant associations between intake of EPA, DHA, or EPA + DHA and the rate of incident PAD⁸⁸. However, recent data from the Danish Diet, Cancer and Health cohort of 3204 participants provided contrary results. The authors found a statistically significant lower rate of PAD in the highest quintile of EPA (hazard ratio [HR]: 0.55) consumption and a nonsignificant lower rate for DHA (HR: 0.79) consumption. They also reported that a high content of marine n-3 polyunsaturated fatty acids (PUFAs) in adipose tissue, in particular EPA, was associated with a lower risk of incident PAD⁸⁹. This is in line with previous observations that marine n-3 PUFAs protect against atherosclerotic diseases in other beds, such as the coronaries (myocardial infarction)⁹⁰ and the carotid and cerebral arteries (ischemic stroke)⁹¹. Supplementation with omega-3 fatty acids does not improve hemodynamic or clinical outcomes in PAD patients⁹². The data on the benefits of marine unsaturated fatty acids on PAD is therefore unclear.

Vitamins/Micronutrients

PAD has an inverse relationship with vitamin A⁹³, vitamin C⁹⁴, vitamin D⁹⁵, vitamin E⁹⁶, and folic acid⁹⁷ intake. International studies have documented deficiency of these vitamins in PAD. In the USA, it was found that vitamin A was low in 57%, vitamin C in 33%, and vitamin E in 100% of patients with PAD⁹⁸. In a study done in Australia, it was found that among PAD patients 16.7% were deficient in vitamin A, 78.1% in vitamin C, 0% in vitamin E, 58.1% in vitamin D, and 50% in iron and zinc⁹⁹. In another study, individuals without PAD had higher plasma vitamins concentrations compared with PAD patients⁸⁴. Several studies have shown a vascular benefit (improved endothelial function) in patients with atherosclerosis with supplementation with vitamin C¹⁰⁰, vitamin E¹⁰¹, and vitamin D¹⁰². Clinical benefit with oral supplementation of these vitamins in a pill form has however not shown any persuasive clinical benefit¹⁰³. On the other hand, a dietary intake of foods high in these vitamins (and iron and zinc) is associated with a decreased prevalence of PAD^{84,93,104}. Eating a variety of food items usually provides a complex combination of nutrients, and this may result in beneficial interactions leading to a decrease in atherosclerotic vascular disease (as compared to supplementation with individual micronutrients).

Special Diets

In the PREDIMED (Primary Prevention of Cardiovascular Disease with a Mediterranean Diet) randomized clinical trial, individuals randomly assigned to the Mediterranean diet (MedD) demonstrated a significantly lower rate of incident PAD when compared to those in the low-fat diet group¹⁰⁵. In a recent systemic review of 82 studies (30% RCT) Adegbola et al. found that the MedD was associated with a lower incidence of PAD¹⁰⁶. In the Framingham Heart Study, HTN increased the risk of intermittent claudication several fold.¹⁰⁷ Hypertension is common in PAD patients¹⁰⁸. Dash diet is effective in reducing HTN and should therefore lead to a reduction in PAD¹⁰⁹. A vegetarian diet is typically higher in fiber, antioxidants, phytochemicals, and plant protein and lower in saturated fat compared to non-vegetarian dietary patterns¹¹⁰. A plant-based diet has significant health benefits, and studies have shown that a plant-based diet can be an effective treatment for obesity¹¹¹, T2DM¹¹², HTN¹¹³, hyperlipidemia¹¹⁴, and heart disease¹¹⁵. A

vegetarian diet helps in preventing or slowing the development of atherosclerosis in several ways. There is prevention of vascular endothelial cell injury, LDL-cholesterol oxidation, and macrophage activation¹¹⁶. Vegetarians have less atherosclerosis, and this should help decrease the incidence of PAD in this population.

Conclusion

Dietary intervention is an inexpensive way to combat the ever-increasing burden of PAD. Peripheral artery disease is associated with increased risk of major cardiovascular events, limb amputation, and overall mortality. As a vascular disease, it is relatively common and highly indicative of atherosclerotic disease in the coronary and the cerebral circulation. It has been noted repeatedly that low consumption of saturated fat, foods of animal origin and salt, and increased intake of plant-based foods such as whole grains, fruits, vegetables, legumes, and nuts, are linked with reduced atherosclerosis risk¹¹⁷. This finding has also been noted in patients with PAD²⁵. In conclusion, there is ample evidence-based data supporting the consumption of a plant based prudent diet for the primary and secondary prevention of PAD.

References

1. Criqui M.H., V. Aboyans. Epidemiology of peripheral artery disease. *Circ Res*, 116 (2015), pp. 1509-1526.
2. Song P, Rudan D, Zhu Y, Fowkes FJI, Rahimi K, Fowkes FGR, et al. Global, regional, and national prevalence and risk factors for peripheral artery disease in 2015: an updated systematic review and analysis. *Lancet Glob Health*. (2019) 7:e1020–30. 10.1016/S2214-109X(19)30255-4.
3. Selvin E, Erlinger TP. Prevalence of and risk factors for peripheral arterial disease in the United States: Results from the National Health and Nutrition Examination Survey, 1999-2000. *Circulation*. (2004) 110:738–43. 10.1161/01.CIR.0000137913.26087.F0.
4. Bauersachs R, Debus S, Nehler M, Huelsebeck M, Balradj J, Bowrin K, et al. . A Targeted literature review of the disease burden in patients with symptomatic peripheral artery disease. *Angiology*. (2020) 71:303–14. 10.1177/0003319719896477.
5. Vogt MT, Wolfson SK, Kuller LH. Lower extremity arterial disease and the aging process: A review. *J Clin Epidemiol* 1992; 45: 529 –542.
6. Fowkes FG. The measurement of atherosclerotic peripheral arterial disease in epidemiological surveys. *International journal of epidemiology*. 1988;17(2):248–254.
7. Chi Y-W, Jaff MR. Optimal risk factor modification and medical management of the patient with peripheral arterial disease. *Catheterization and cardiovascular interventions: official journal of the Society for Cardiac Angiography and Interventions*. 2008;71(4):475–489.
8. Eraso LH, Fukaya E, Mohler ER III, Xie D, Sha D, Berger JS. Peripheral arterial disease, prevalence and cumulative risk factor profile analysis. *Eur J Prev Cardiol*. 2014;21(6):704–711.

9. Meijer WT, Hoes AW, Rutgers D, Bots ML, Hofman A, Grobbee DE. Peripheral arterial disease in the elderly: the Rotterdam Study. *Arterioscler Thromb Vasc Biol*. 1998;18(2):185–192.
10. Morley R.L., Sharma A., Horsch A.D., Hinchliffe R.J. Peripheral artery disease. *BMJ*. 2018;360 doi: 10.1136/bmj.j5842.
11. Kayashima Y, Maeda-Smithies N. Atherosclerosis in Different Vascular Locations Unbiasedly Approached with Mouse Genetics. *Genes (Basel)*. 2020 Nov 28;11(12):1427. doi: 10.3390/genes11121427.
12. Libby P., Ridker P. M., Hansson G. K. Progress and challenges in translating the biology of atherosclerosis. *Nature*. 2011;473(7347):317–325. doi: 10.1038/nature10146.
13. Marrett E., DiBonaventura M. D., Zhang Q. Burden of peripheral arterial disease in Europe and the United States: a patient survey. *Health and Quality of Life Outcomes*. 2013;11(175).
14. Howard D. P. J., Banerjee A., Fairhead J. F., Hands L., Silver L. E., Rothwell P. M. Population-based study of incidence, risk factors, outcome, and prognosis of ischemic peripheral arterial events: implications for prevention. *Circulation*. 2015;132(19):1805–1815. doi: 10.1161/CIRCULATIONAHA.115.016424.
15. Gerhard-Herman M. D., Gornik H. L., Barrett C., et al. AHA/ACC guideline on the management of patients with lower extremity peripheral artery disease: a report of the American college of cardiology/American heart association task force on clinical practice guidelines. *JACC: Journal of the American College of Cardiology*. 2016;69:e71–e126.
16. Novakovic M, Jug B, Lenasi H. Clinical impact of exercise in patients with peripheral arterial disease. *Vascular*. 2017 Aug;25(4):412–422. doi: 10.1177/1708538116678752.
17. Regensteiner JG, Hiatt WR. Current medical therapies for patients with peripheral arterial disease: a critical review. *The American Journal of Medicine*. 2002;112:49–57.
18. Patterson RB, Pinto B, Marcus B, Colucci A, Braun T, Roberts M. Value of a supervised exercise program for the therapy of arterial claudication. *Journal of Vascular Surgery*. 1997;25:312–319.
19. Stansby G., Mister R., Fowkes G., et al. High risk of peripheral arterial disease in the United Kingdom: 2-year results of a prospective registry. *Angiology*. 2011;62(2):111–118. doi: 10.1177/0003319710387917.
20. Kochar A, Mulder H, Rockhold FW, Baumgartner I, Berger JS, et al. Cause of Death Among Patients With Peripheral Artery Disease: Insights From the EUCLID Trial. *Circ Cardiovasc Qual Outcomes*. 2020 Nov;13(11):e006550. doi: 10.1161/CIRCOUTCOMES.120.006550.
21. Chase M. R., Friedman H. S., Navaratnam P., Heithoff K., Simpson Jr. R. J. Comparative assessment of medical resource use and costs associated with patients with symptomatic peripheral artery disease in the United States. *Journal of Managed Care and Specialty Pharmacy*. 2016;22(6):667–675. doi: 10.18553/jmcp.2016.15010.
22. Lechner K, von Schacky C, McKenzie AL, Worm N, Nixdorff U, et al. Lifestyle factors and high-risk atherosclerosis: Pathways and mechanisms beyond traditional risk factors. *Eur J Prev Cardiol*. 2020 Mar;27(4):394–406. doi: 10.1177/2047487319869400.

23. Bhatt D. L., Steg P. G., Ohman E. M., et al. International prevalence, recognition, and treatment of cardiovascular risk factors in outpatients with atherothrombosis. *Journal of the American Medical Association*. 2006;295(2):180–189. doi: 10.1001/jama.295.2.180.
24. Noflatscher M, Schreinlechner M, Sommer P, et al. Association of Food and Alcohol Consumption with Peripheral Atherosclerotic Plaque Volume as Measured by 3D-Ultrasound. *Nutrients*. 2020;12(12):3711. Published 2020 Nov 30. doi:10.3390/nu12123711.
25. Kulezic A, Bergwall S, Fatemi S, Sonestedt E, Zarrouk M, Gottsäter A, Acosta S. Healthy diet and fiber intake are associated with decreased risk of incident symptomatic peripheral artery disease - A prospective cohort study. *Vasc Med*. 2019 Dec;24(6):511-518. doi: 10.1177/1358863X19867393.
26. Hisamatsu T., Miura K., Fujiyoshi A., Kadota A., Miyagawa N., Satoh A., Zaid M., Yamamoto T., Horie M., Ueshima H., et al. Serum magnesium, phosphorus, and calcium levels and subclinical calcific aortic valve disease: A population-based study. *Atherosclerosis*. 2018;273:145–152. doi: 10.1016/j.atherosclerosis.2018.03.035.
27. Martín-Calvo N., Martínez-González M.Á. Vitamin C Intake is Inversely Associated with Cardiovascular Mortality in a Cohort of Spanish Graduates: The SUN Project. *Nutrients*. 2017;9:954. doi: 10.3390/nu9090954.
28. Carrero JJ, Grimble RF. Does nutrition have a role in peripheral vascular disease? *Br J Nutr* 2006;95:217–29.
29. Klipstein-Grobusch K, den Breeijen JH, Grobbee DE, Boeing H, Hofman A, Witteman JC. Dietary antioxidants and peripheral arterial disease : the Rotterdam Study. *American journal of epidemiology*. 2001 Jul 15;154(2):145–149.
30. Golay A, Bobbioni E. The role of dietary fat in obesity. *Int J Obes Relat Metab Disord*. 1997 Jun;21 Suppl 3:S2-11.
31. Jensen MD, Ryan DH, Apovian CM, Ard JD, Comuzzie AG, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Circulation*. 2014;129(suppl 2):S102–S138. doi: 10.1161/01.cir.0000437739.71477.ee.
32. Després JP. Abdominal obesity and cardiovascular disease: is inflammation the missing link? *Can J Cardiol*. 2012 Nov-Dec;28(6):642-52. doi: 10.1016/j.cjca.2012.06.004.
33. Staynor JMD, Smith MK, Donnelly CJ, Sallam AE, Ackland TR. DXA reference values and anthropometric screening for visceral obesity in Western Australian adults. *Sci Rep*. 2020 Oct 30;10(1):18731. doi: 10.1038/s41598-020-73631-x.
34. Ross R, Neeland IJ, Yamashita S, et al. Waist circumference as a vital sign in clinical practice: a Consensus Statement from the IAS and ICCR Working Group on Visceral Obesity. *Nat Rev Endocrinol*. 2020;16(3):177-189. doi:10.1038/s41574-019-0310-7.
35. <https://www.ncbi.nlm.nih.gov/mesh?Cmd=DetailsSearch&Term=%22Waist-Hip+Ratio%22%5BMeSH+Terms%5D>.
36. Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and

meta-analysis. *Obes Rev.* 2012 Mar;13(3):275-86. doi: 10.1111/j.1467-789X.2011.00952.x.

37. Hu L, Huang X, You C, Li J, Hong K, Li P, Wu Y, Wu Q, Wang Z, Gao R, Bao H, Cheng X. Prevalence of overweight, obesity, abdominal obesity and obesity-related risk factors in southern China. *PLoS One.* 2017 Sep 14;12(9):e0183934. doi: 10.1371/journal.pone.0183934.
38. Planas A, Clara A, Pou JM, Vidal-Barraquer F, Gasol A, et al. (2001) Relationship of obesity distribution and peripheral arterial occlusive disease in elderly men. *Int J Obes Relat Metab Disord* 25: 1068–1070.
39. Wisse BE (2004) The inflammatory syndrome: the role of adipose tissue cytokines in metabolic disorders linked to obesity. *J Am Soc Nephrol* 15: 2792–2800.
40. Zoccali C, Mallamaci F, Tripepi G (2004) Inflammatory proteins as predictors of cardiovascular disease in patients with end-stage renal disease. *Nephrol Dial Transplant* 19: V67–V72.
41. Tzoulaki I, Murray GD, Lee AJ, Rumley A, Lowe GD, et al. (2005) C-reactive protein, interleukin-6, and soluble adhesion molecules as predictors of progressive peripheral atherosclerosis in the general population: Edinburgh Artery Study. *Circulation* 112: 976–983.
42. Brouwer BG, Visseren FL, Stolk RP, van der Graaf Y; SMART Study Group. Abdominal fat and risk of coronary heart disease in patients with peripheral arterial disease. *Obesity (Silver Spring).* 2007 Jun;15(6):1623-30. doi: 10.1038/oby.2007.192.
43. Fock KM, Khoo J. Diet and exercise in management of obesity and overweight. *J Gastroenterol Hepatol.* 2013 Dec;28 Suppl 4:59-63. doi: 10.1111/jgh.12407.
44. Di Buono M., Hannah J.S., Katznel L.I., Jones P.J. Weight loss due to energy restriction suppresses cholesterol biosynthesis in overweight, mildly hypercholesterolemic men. *J. Nutr.* 1999;129:1545–1548.
45. Melanson K., Gootman J., Myrdal A., Kiline G., Rippe J.M. Weight loss and total lipid profile changes in overweight women consuming beef or chicken as the primary protein source. *Nutrition.* 2003;19:409–414. doi: 10.1016/S0899-9007(02)01080-8.
46. Neter J.E., Stam B.E., Kok F.J., Brobbee D.E., Geleijnse J.M. Influence of weight reduction on blood pressure: A meta-analysis of randomized controlled trials. *Hypertension.* 2003;42:878–884. doi: 10.1161/01.HYP.0000094221.86888.AE.
47. Ryan A.S., Nicklas J.B. Reductions in plasma cytokine levels with weight loss improve insulin sensitivity in overweight and obese postmenopausal women. *Diabetes Care.* 2004;27:1699–1705. doi: 10.2337/diacare.27.7.1699.
48. Claessens M., van Baak M.A., Monsheimer S., Saris W.H. The effect of a low-fat, high-protein or high-carbohydrate ad libitum diet on weight loss maintenance and metabolic risk factors. *Int. J. Obes.* 2009;33:296–304. doi: 10.1038/ijo.2008.278.
49. Tamar S. Polonsky, Lu Tian, Dongxue Zhang, Lydia A. Bazzano, Michael H. Criqui, et al. Associations of Weight Change With Changes in Calf Muscle Characteristics and Functional Decline in Peripheral Artery Disease. *Journal of the American Heart Association.* 2019;8:e010890. <https://doi.org/10.1161/JAHA.118.010890/>

50. Satija A, Hu FB. Plant-based diets and cardiovascular health. *Trends Cardiovasc Med*. 2018 Oct;28(7):437-441. doi: 10.1016/j.tcm.2018.02.004.
51. Hung HC, Merchant A, Willett W, Ascherio A, Rosner BA, Rimm E, Joshipura KJ. The association between fruit and vegetable consumption and peripheral arterial disease. *Epidemiology* 2003;14:659–65.
52. Ciccarone E., Di Castelnuovo A., Salcuni M., Siani A., Giacco A., et al. A high-score Mediterranean dietary pattern is associated with a reduced risk of peripheral arterial disease in Italian patients with Type 2 diabetes. *J. Thromb. Haemost.* 2003;1:1744–1752. doi: 10.1046/j.1538-7836.2003.00323.x.
53. Heffron S.P., Rockman C.B., Adelman M.A., Gianos E., Guo Y., Xu J.F., Berger J.S. Greater Frequency of Fruit and Vegetable Consumption Is Associated with Lower Prevalence of Peripheral Artery Disease. *Arterioscl. Throm. Vas.* 2017;37:1234–1240. doi: 10.1161/ATVBAHA.116.308474.
54. Chen GC, Arthur R, Mossavar-Rahmani Y, Xue X, Haring B, et al. Adherence to Recommended Eating Patterns Is Associated With Lower Risk of Peripheral Arterial Disease: Results From the Women's Health Initiative. *Hypertension*. 2021 Aug;78(2):447-455. doi: 10.1161/HYPERTENSIONAHA.121.17432.
55. Howard B.V., Kritchevsky D. Phytochemicals and cardiovascular disease. A statement for healthcare professionals from the American Heart Association. *Circulation*. 1997;95:2591–2593. doi: 10.1161/01.CIR.95.11.2591.
56. Ogilvie, RP, Lutsey, PL, Heiss, G, et al. Dietary intake and peripheral arterial disease incidence in middle-aged adults: The Atherosclerosis Risk in Communities (ARIC) Study. *Am J Clin Nutr* 2017; 105: 651–659.
57. Gimeno SG, Hirai AT, Harima HA, Kikuchi MY, Simony RF, de Barros N Jr, Cardoso MA, Ferreira SR; Japanese-Brazilian Diabetes Study Group. Fat and fiber consumption are associated with peripheral arterial disease in a cross-sectional study of a Japanese-Brazilian population. *Circ J*. 2008 Jan;72(1):44-50. doi: 10.1253/circj.72.44.
58. Hunninghake DB, Miller VT, LaRosa JC, Kinoshian B, Brown V, Howard WJ, et al. Hypocholesterolemic effects of a dietary fiber supplement. *Am J Clin Nutr*. 1994;59(5):1050–1054.
59. Mazidi M, Kengne AP, Mikhailidis DP, Toth PP, Ray KK, Banach M. Dietary food patterns and glucose/insulin homeostasis: a cross-sectional study involving 24,182 adult Americans. *Lipids Health Dis*. 2017;16(1):192. doi: 10.1186/s12944-017-0571-x.
60. Guo-Chong Chen, Rhonda Arthur, Yasmin Mossavar-Rahmani, Xiaonan Xue, Bernhard Haring et al. Adherence to Recommended Eating Patterns Is Associated With Lower Risk of Peripheral Arterial Disease: Results From the Women's Health Initiative. *Hypertension*. 2021;78:447–455. <https://doi.org/10.1161/HYPERTENSIONAHA.121.17432>.
61. Heffron SP, Rockman CB, Gianos E, Guo Y, Berger JS. Greater frequency of nut consumption is associated with lower prevalence of peripheral arterial disease. *Prev Med* 2015;72:15–8.

62. Jiang R., Jacobs D.R. Jr, Mayer-Davis E., Szklo M., Herrington D., Jenny N.S., Kronmal R., Barr R.G. Nut and seed consumption and inflammatory markers in the Multi-Ethnic Study of Atherosclerosis. *Am. J. Epidemiol.* 2006;163:222–231.
63. Simon J, Fung K, Raisi-Estabragh Z, Aung N, Khanji MY, et al. Light to moderate coffee consumption is associated with lower risk of death: a UK Biobank study. *Eur J Prev Cardiol.* 2022 Jan 20, doi: 10.1093/eurjpc/zwac008.
64. Ochiai, R. , Sugiura Y., Shioya Y., Otsuka K., Katsuragi Y., and Hashiguchi T.. 2014. Coffee polyphenols improve peripheral endothelial function after glucose loading in healthy male adults. *Nutr. Res.* 34:155–159.
65. Mattioli AV, Migaldi M, Farinetti A. Coffee in hypertensive women with asymptomatic peripheral arterial disease: a potential nutraceutical effect. *J Cardiovasc Med (Hagerstown).* 2018 Apr;19(4):183-185. doi: 10.2459/JCM.0000000000000626.
66. Roberts KA, Draijer R, Hopkins ND, de Graaf Y, Holder SM, Carter SE, Thijssen DHJ, Low DA. Impact of green tea on the deleterious cardiometabolic effects of 7-days unhealthy lifestyle in young healthy males. *Physiol Rep.* 2021 Mar;9(5):e14720. doi: 10.14814/phy2.14720.
67. Grassi, D., Desideri, G., Croce, G. , Tiberti, S. , Aggio, A. , & Ferri, C. (2009). Flavonoids, vascular function and cardiovascular protection. *Current Pharmaceutical Design*, 15, 1072–1084.
68. Grassi, D., Desideri, G., Di Giosia, P. , De Feo, M. , Fellini, E. , Cheli, P. , Ferri, L. , & Ferri, C. (2013). Tea, flavonoids, and cardiovascular health: Endothelial protection. *American Journal of Clinical Nutrition*, 98, 1660s–1666s.
69. Loffredo L, Perri L, Catasca E, Pignatelli P, Brancorsini M, Nocella C, De Falco E, Bartimoccia S, Frati G, Carnevale R, Violi F. Dark chocolate acutely improves walking autonomy in patients with peripheral artery disease. *J Am Heart Assoc.* 2014 Jul 2;3(4):e001072. doi: 10.1161/JAHA.114.001072.
70. Langer S, Marshall LJ, Day AJ, Morgan MR. Flavanols and methylxanthines in commercially available dark chocolate: a study of the correlation with nonfat cocoa solids. *J Agric Food Chem.* 2011; 59:8435-8441.
71. Heiss C, Kleinbongard P, Dejam A, Perre S, Schroeter H, Sies H, Kelm M. Acute consumption of flavanol-rich cocoa and the reversal of endothelial dysfunction in smokers. *J Am Coll Cardiol.* 2005; 46:1276-1283.
72. Mukamal KJ, Kennedy M, Cushman M, Kuller LH, Newman AB, Polak J, Criqui MH, Siscovick DS. Alcohol consumption and lower extremity arterial disease among older adults: the cardiovascular health study. *Am J Epidemiol.* 2008 Jan 1;167(1):34-41. doi: 10.1093/aje/kwm274,
73. Gaziano JM, Buring JE, Breslow JL, Goldhaber SZ, Rosner B, VanDenburgh M, Willett W, Hennekens CH. Moderate alcohol intake, increased levels of high-density lipoprotein and its subfractions, and decreased risk of myocardial infarction. *N Engl J Med* 1993;329:1829–34.
74. Schrieks IC, Heil AL, Hendriks HF, Mukamal KJ, Beulens JW. The effect of alcohol consumption on insulin sensitivity and glycemic status: a systematic review and meta-analysis of intervention studies. *Diabetes Care* 2015;38:723–32.

75. Micha R, Wallace SK, Mozaffarian D. Red and processed meat consumption and risk of incident coronary heart disease, stroke, and diabetes mellitus: a systematic review and meta-analysis. *Circulation* 2010;121:2271–83.
76. Abete I, Romaguera D, Vieira AR, Lopez de Munain A, Norat T. Association between total, processed, red and white meat consumption and all-cause, CVD and IHD mortality: a meta-analysis of cohort studies. *Br J Nutr* 2014;112:762–75.
77. Haring B, Misialek JR, Rebholz CM, Petruski-Ivleva N, Gottesman RF, Mosley TH, Alonso A. Association of dietary protein consumption with incident silent cerebral infarcts and stroke: the Atherosclerosis Risk in Communities (ARIC) study. *Stroke* 2015;46:3443–50.
78. Karppanen H, Mervaala E. Sodium intake and hypertension. *Prog Cardiovasc Dis.* 2006 Sep-Oct;49(2):59-75. doi: 10.1016/j.pcad.2006.07.001.
79. Micha R, Michas G, Mozaffarian D. Unprocessed red and processed meats and risk of coronary artery disease and type 2 diabetes—an updated review of the evidence. *Curr Atheroscler Rep* 2012;14:515–24.
80. Aune D, Ursin G, Veierod MB. Meat consumption and the risk of type 2 diabetes: a systematic review and meta-analysis of cohort studies. *Diabetologia* 2009;52:2277–87.
81. Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Saturated fatty acids and risk of coronary heart disease: modulation by replacement nutrients. *Curr Atheroscler Rep.* 2010;12(6):384-390. doi:10.1007/s11883-010-0131-6.
82. Gimeno SG, Hirai AT, Harima HA, Kikuchi MY, Simony RF, de Barros N Jr, Cardoso MA, Ferreira SR; Japanese-Brazilian Diabetes Study Group. Fat and fiber consumption are associated with peripheral arterial disease in a cross-sectional study of a Japanese-Brazilian population. *Circ J.* 2008 Jan;72(1):44-50. doi: 10.1253/circj.72.44.
83. Naqvi AZ, Davis RB, Mukamal KJ. Dietary fatty acids and peripheral artery disease in adults. *Atherosclerosis.* 2012 Jun;222(2):545-50. doi: 10.1016/j.atherosclerosis.2012.03.029.
84. Mazidi M, Wong ND, Katsiki N, Mikhailidis DP, Banach M. Dietary patterns, plasma vitamins and Trans fatty acids are associated with peripheral artery disease. *Lipids Health Dis.* 2017;16(1):254. Published 2017 Dec 28. doi:10.1186/s12944-017-0635-y.
85. J.E. Enns, A. Yeganeh, R. Zarychanski, et al. The impact of omega-3 polyunsaturated fatty acid supplementation on the incidence of cardiovascular events and complications in peripheral arterial disease: a systematic review and meta-analysis. *BMC Cardiovasc Disord*, 14 (2014), p. 70.
86. A. Campbell, J. Price, W.R. Hiatt. Omega-3 fatty acids for intermittent claudication. *Cochrane Database Syst Rev*, 7 (2013), p. CD003833.
87. Mann NJ, O'Connell SL, Baldwin KM, Singh I, Meyer BJ. Effects of seal oil and tuna-fish oil on platelet parameters and plasma lipid levels in healthy subjects. *Lipids.* 2010 Aug;45(8):669-81. doi: 10.1007/s11745-010-3450-z.
88. Tram L, Bork CS, Venø SK, Lasota AN, Lundbye-Christensen S, Schmidt EB, Overvad K. Intake of marine n-3 polyunsaturated fatty acids and the risk of incident peripheral artery disease. *Eur J Clin Nutr.* 2021 Oct;75(10):1483-1490. doi: 10.1038/s41430-021-00858-w.

89. Lasota AN, Grønholdt MM, Bork CS, Lundbye-Christensen S, Overvad K, Schmidt EB. Marine n-3 Fatty Acids and the Risk of Peripheral Arterial Disease. *J Am Coll Cardiol*. 2018 Oct 2;72(14):1576-1584. doi: 10.1016/j.jacc.2018.07.045.
90. A. Gammelmark, M.S. Nielsen, C.S. Bork, et al. Adipose tissue content of marine n-3 polyunsaturated fatty acids is inversely associated with myocardial infarction. *J Am Coll Cardiol*, 67 (2016), pp. 1008-1009.
91. H. Saber, M.Y. Yakoob, P. Shi, et al. Omega-3 fatty acids and incident ischemic stroke and its atherothrombotic and cardioembolic subtypes in 3 US cohorts. *Stroke*, 48 (2017), pp. 2678-2685.
92. Schiano V, Laurenzano E, Brevetti G, De Maio JJ, Lanero S, Scopacasa F, Chiariello M. Omega-3 polyunsaturated fatty acid in peripheral arterial disease: effect on lipid pattern, disease severity, inflammation profile, and endothelial function. *Clin Nutr*. 2008(2):241-7. doi:10.1016/j.clnu.2007.11.007.
93. Naqvi AZ, Davis RB, Mukamal KJ. Nutrient intake and peripheral artery disease in adults: key considerations in cross-sectional studies. *Clin Nutr* 2014;33:443–7.
94. Langlois M, Duprez D, Delanghe J, De Buyzere M, Clement DL. Serum vitamin C concentration is low in peripheral arterial disease and is associated with inflammation and severity of atherosclerosis. *Circulation* 2001;103:1863–8.
95. Zagura M, Serg M, Kampus P, Zilmer M, Eha J, Unt E, Lieberg J, Cockcroft JR, Kals J. Aortic stiffness and vitamin D are independent markers of aortic calcification in patients with peripheral arterial disease and in healthy subjects. *Eur J Vasc Endovasc Sur* 2011;42:689–95.
96. Klipstein-Grobusch K, den Breeijen JH, Grobbee DE, Boeing H, Hofman A, Witteman JC. Dietary antioxidants and peripheral arterial disease: the Rotterdam Study. *Am J Epidemiol* 2001;154:145–9.
97. Wilmsink AB, Welch AA, Quick CR, Burns PJ, Hubbard CS, Bradbury AW, Day NE. Dietary folate and vitamin B6 are independent predictors of peripheral arterial occlusive disease. *J Vasc Surg* 2004;39:513–6.
98. Gardner A.W., Bright B.C., Ort K.A., Montgomery P.S. Dietary Intake of Participants with Peripheral Artery Disease and Claudication. *Angiology*. 2011;62:270–275. doi: 10.1177/0003319710384395.
99. Thomas J., Delaney C., Suen J., Miller M. Nutritional status of patients admitted to a metropolitan tertiary care vascular surgery unit. *Asia Pac. J. Clin. Nutr*. 2019;28:64. doi: 10.6133/apjcn.201903_28(1).0010.
100. Ashor A.W., Lara J., Mathers J.C., Siervo M. Effect of vitamin C on endothelial function in health and disease: A systematic review and metaanalysis of randomised controlled trials. *Atherosclerosis*. 2014;235:9–20. doi: 10.1016/j.atherosclerosis.2014.04.004.
101. Hosseini B., Saedisomeolia A., Skilton M.R. Association between Micronutrients Intake/Status and Carotid Intima Media Thickness: A Systematic Review. *J. Acad. Nutr. Diet*. 2017;117:69–82. doi: 10.1016/j.jand.2016.09.031.
102. Beveridge L.A., Khan F., Struthers A.D., Armitage J., Barchetta I., et al. Effect of Vitamin D Supplementation on Markers of Vascular Function: A Systematic Review and

- Individual Participant Meta-Analysis. *J. Am. Heart Assoc.* 2018;7:E008273. doi: 10.1161/JAHA.117.008273.
103. Myung S.K., Ju W., Cho B., Oh S.W., Park S.M., Koo B.K., Park B.J., Korean Meta-Analysis Study Group Efficacy of vitamin and antioxidant supplements in prevention of cardiovascular disease: Systematic review and meta-analysis of randomised controlled trials. *BMJ.* 2013;346:f10. doi: 10.1136/bmj.f10.
 104. Antonelli-Incalzi R, Pedone C, McDermott MM, Bandinelli S, Miniati B, Lova RM, et al. Association between nutrient intake and peripheral artery disease: results from the InCHIANTI study. *Atherosclerosis.* 2006;186(1):200–206. doi: 10.1016/j.atherosclerosis.2005.07.016
 105. Ruiz-Canela M, Estruch R, Corella D, Salas-Salvado J, Martinez-Gonzalez MA. Association of Mediterranean diet with peripheral artery disease: the PREDIMED randomized trial. *JAMA* 2014;311:415–7.
 106. Adegbola A, Behrendt CA, Zyriax BC, Windler E, Kreutzburg T. The impact of nutrition on the development and progression of peripheral artery disease: A systematic review. *Clin Nutr.* 2022 Jan;41(1):49–70. doi: 10.1016/j.clnu.2021.11.005.
 107. Kannel WB, McGee DL. Update on some epidemiologic features of intermittent claudication: the Framingham Study. *Journal of the American Geriatrics Society.* 1985;33(1):13–18.
 108. Wisman PP, Tangelder MJ, van Hattum ES, de Borst GJ, Moll FL. Young women with PAD are at high risk of cardiovascular complications. *European journal of vascular and endovascular surgery: the official journal of the European Society for Vascular Surgery.* 2012;43(4):441–445.
 109. Filippou CD, Tsioufis CP, Thomopoulos CG, Mihas CC, Dimitriadis KS, et al. Dietary Approaches to Stop Hypertension (DASH) Diet and Blood Pressure Reduction in Adults with and without Hypertension: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Adv Nutr.* 2020 Sep 1;11(5):1150–1160. doi: 10.1093/advances/nmaa041.
 110. Satija A, Hu FB. Plant-based diets and cardiovascular health. *Trends Cardiovasc Med.* (2018) 28:437–41. 10.1016/j.tcm.2018.02.004.
 111. Wang Y, Beydoun MA. Meat consumption is associated with obesity and central obesity among US adults. *Int J Obes (Lond)* 2009 Jun;33(6):621–8. DOI: <http://dx.doi.org/10.1038/ijo.2009.45>.
 112. Kim M.S., Hwang S.S., Park E.J., Bae J.W. Strict vegetarian diet improves the risk factors associated with metabolic diseases by modulating gut microbiota and reducing intestinal inflammation. *Environ Microbiol Rep.* 2013;5(5):765–775. doi: 10.1111/1758-2229.12079.
 113. Yokoyama Y., Nishimura K., Barnard N.D. Vegetarian diets and blood pressure: a meta-analysis. *JAMA Intern Med.* 2014;174(4):577–587. doi: 10.1001/jamainternmed.2013.14547.
 114. Yokoyama Y, Levin SM, Barnard ND. Association between plant-based diets and plasma lipids: a systematic review and meta-analysis. *Nutr Rev.* 2017 Sep 1;75(9):683–698. doi: 10.1093/nutrit/nux030.

115. Glenn AJ, Viguioliouk E, Seider M, Boucher BA, Khan TA, et al. Relation of Vegetarian Dietary Patterns With Major Cardiovascular Outcomes: A Systematic Review and Meta-Analysis of Prospective Cohort Studies. *Front Nutr.* 2019 Jun 13;6:80. doi: 10.3389/fnut.2019.00080.
116. Tusso P, Stoll SR, Li WW. A plant-based diet, atherogenesis, and coronary artery disease prevention. *Perm J.* 2015;19(1):62-67. doi:10.7812/TPP/14-036.
117. Riccardi G, Giosuè A, Calabrese I, Vaccaro O. Dietary recommendations for prevention of atherosclerosis. *Cardiovasc Res.* 2021 Jul 6;cvab173. doi: 10.1093/cvr/cvab173.